

-WILDNESS OF SOME CLASSES OF C^ -ALGEBRAS

SERGIO ALBEVERIO, KATE JUSHENKO, DANIIL PROSKURIN, AND YURII SAMOILENKO

Dedicated to the memory of Yuri Daletskii.

ABSTRACT. We consider the complexity of the representation theory of free products of C^* -algebras. Necessary and sufficient conditions for the free product of finite-dimensional C^* -algebras to be $*$ -wild is presented. As a corollary we get criteria for $*$ -wildness of free products of finite groups. It is proved that the free product of a non-commutative nuclear C^* -algebra and the algebra of continuous functions on the one-dimensional sphere is $*$ -wild. This result is applied to estimate the complexity of the representation theory of certain C^* -algebras generated by isometries and partial isometries.

INTRODUCTION

The problem of estimating the complexity of the representation theory of a certain C^* -algebra ($*$ -algebra) is one of important questions of representations theory. In order to compare C^* -algebras according to the complexity of their categories of representations S. A. Kruglyak and Yu. S. Samoilenko proposed the partial order \succ , see [6, 7]. Informally, if \mathcal{A} and \mathcal{B} are C^* -algebras, then $\mathcal{A} \succ \mathcal{B}$ if the problem of classification of $*$ -representations of \mathcal{A} contains, as a subproblem, a classification of $*$ -representations of \mathcal{B} , see Preliminaries below for precise definitions. A C^* -algebra \mathcal{A} is called $*$ -wild if $\mathcal{A} \succ C^*(\mathcal{F}_2)$, where \mathcal{F}_2 is free group with two generators. In fact, if a C^* -algebra is $*$ -wild, then the problem of unitary classification of its irreducible representations contains, as a subproblem, the problem of classification of irreducible representations of any finitely-generated C^* -algebra, see [6].

The notion of $*$ -wildness is closely related to the notion of a free product, see [13] for details on the free products of algebras like C^* -algebras and von Neumann algebras. Indeed, the model example of a $*$ -wild C^* -algebra, the algebra $C^*(\mathcal{F}_2)$, is the C^* -free product of two copies of $C(\mathbf{T})$, the algebra of continuous functions on the circle. A natural problem that arises is to establish when free products of C^* -algebras are $*$ -wild.

In this paper we give a complete answer for free products of finite-dimensional C^* -algebras.

The next step is to consider free products of non-commutative and commutative algebras. In this paper we show that the free product of a non-commutative nuclear algebra having non-trivial projection and $C(\mathbf{T})$ is $*$ -wild. In particular, this fact implies $*$ -wildness of the algebras $M_n(\mathbb{C}) * C(\mathbf{T})$, $n \geq 2$. This result is then applied to prove the $*$ -wildness of C^* -algebras generated by partial isometries with linearly dependent range projections and C^* -algebras generated by families of q -commuting isometries (pairs of q -commuting isometries were also studied in [4]).

The paper is organized as follows. In preliminaries we modify the definition of majorization given in [6, 7], to make it less restrictive but preserving the most important properties of majorization. Then we give a definition of $*$ -wild C^* -algebra and list basic

2000 *Mathematics Subject Classification.* Primary 16G60, 46L05, 46L09; Secondary 22D20.

Key words and phrases. $*$ -Representations, free product, $*$ -wild algebra.

properties of $*$ -wild C^* -algebras. Finally, we recall the definitions and some properties of free products of C^* -algebras.

In Section 1, we study free products of finite-dimensional C^* -algebras. Namely, we give criteria of $*$ -wildness for such free products. As a corollary we obtain necessary and sufficient conditions for $*$ -wildness of free products of finite groups.

In Section 2.1, it is proved that the C^* -free product of a nuclear C^* -algebra with a projection and $C(\mathbf{T})$ is $*$ -wild. In Section 2.2, we apply this result to prove $*$ -wildness of certain C^* -algebras generated by isometries and partial isometries.

0. PRELIMINARIES

0.1. **Enveloping C^* -algebras.** In this subsection we recall the definitions of the enveloping C^* -algebra and the group C^* -algebra, which can be found for example in [3], [5].

Let \mathcal{A} be a $*$ -algebra. The C^* -algebra $\tilde{\mathcal{A}}$ with a $*$ -homomorphism $\varphi : \mathcal{A} \mapsto \tilde{\mathcal{A}}$ is called an enveloping C^* -algebra of the algebra \mathcal{A} if for every representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ of \mathcal{A} there exists a unique representation $\tilde{\pi} : \tilde{\mathcal{A}} \rightarrow \mathcal{B}(\mathcal{H})$ of $\tilde{\mathcal{A}}$ such that the following diagram is commutative:

$$\begin{array}{ccc}
 \mathcal{A} & & \\
 \varphi \downarrow & \searrow \pi & \\
 \tilde{\mathcal{A}} & \xrightarrow{\tilde{\pi}} & \mathcal{B}(\mathcal{H})
 \end{array}$$

A group C^* -algebra of a group G is an enveloping C^* -algebra of the group ring $\mathbb{C}[G]$. More precisely, the group C^* -algebra, $C^*(G)$, is the completion of $\mathbb{C}[G]$ with respect to the C^* -norm

$$\|a\| = \sup\{\pi(a) : \pi \in \text{Rep}(\mathbb{C}[G])\}.$$

0.2. **$*$ -Wild C^* -algebras and their properties.** Recall that the category of representations of a certain C^* -algebra \mathcal{A} , denoted by $\text{Rep } \mathcal{A}$, has representations of \mathcal{A} as objects and intertwining operators as morphisms.

Definition 1. We say that a C^* -algebra \mathcal{A} is majorized by a C^* -algebra \mathcal{B} , $\mathcal{A} \prec \mathcal{B}$, if there exists a homomorphism

$$\varphi : \mathcal{B} \rightarrow \mathcal{C} \otimes \mathcal{A},$$

where \mathcal{C} is a nuclear C^* -algebra, and an irreducible representation

$$\tilde{\pi} : \mathcal{C} \rightarrow \mathcal{B}(\mathcal{H})$$

such that the functor

$$\mathcal{F}_{\mathcal{C},\varphi,\tilde{\pi}} : \text{Rep } \mathcal{A} \rightarrow \text{Rep } \mathcal{B}$$

defined by

$$\begin{aligned}
 \mathcal{F}_{\mathcal{C},\varphi,\tilde{\pi}}(\pi) &= (\tilde{\pi} \otimes \pi) \circ \varphi, \quad \pi \in \text{Ob}(\text{Rep } \mathcal{A}), \\
 \mathcal{F}_{\mathcal{C},\varphi,\tilde{\pi}}(A) &= 1 \otimes A, \quad A \in \text{Mor}(\pi_1, \pi_2)
 \end{aligned}$$

is full.

Remark 1. It can be shown, see [11], that if the functor $\mathcal{F}_{\mathcal{C},\varphi,\tilde{\pi}}$ is full then it is automatically faithful.

Remark 2. Originally in [6, 7], the authors considered the matrix algebras $M_n(\mathbb{C})$ with the identical representation instead of an arbitrary nuclear C^* -algebra \mathcal{C} in the definition of majorization. However that definition was quite restrictive. Namely, if \mathcal{A} has not finite-dimensional irreducible representations, then $\mathcal{A} \not\prec C^*(\mathcal{F}_2)$ according to the original definition. In Section 2 we present an example of such a majorization in our sense.

Let us show that the majorization is a partial order.

Proposition 1. *If $\mathcal{A} \prec \mathcal{B}$ and $\mathcal{B} \prec \mathcal{C}$, then $\mathcal{A} \prec \mathcal{C}$.*

Proof. Let \mathcal{C}_i , $i = 1, 2$, be nuclear C^* -algebras and

$$\tilde{\pi}_i: \mathcal{C}_i \rightarrow B(\mathcal{H}_i), \quad i = 1, 2$$

be their irreducible representations. By Definition 1, there are homomorphisms

$$\varphi_1: \mathcal{B} \rightarrow \mathcal{C}_1 \otimes \mathcal{A}, \quad \varphi_2: \mathcal{C} \rightarrow \mathcal{C}_2 \otimes \mathcal{B}$$

such that the induced functors $\mathcal{F}_{\mathcal{C}, \varphi_i, \tilde{\pi}}$ are full. Consider the homomorphism

$$\varphi := (\text{id}_{\mathcal{C}_2} \otimes \varphi_1) \circ \varphi_2: \mathcal{C} \rightarrow \mathcal{C}_2 \otimes \mathcal{C}_1 \otimes \mathcal{A}$$

and the irreducible representation

$$\tilde{\pi} := \tilde{\pi}_2 \otimes \tilde{\pi}_1: \mathcal{C}_2 \otimes \mathcal{C}_1 \rightarrow B(\mathcal{H}_2 \otimes \mathcal{H}_1).$$

Now we prove that the functor

$$\mathcal{F}_{\mathcal{C}, \varphi, \tilde{\pi}}: \text{Rep } \mathcal{A} \rightarrow \text{Rep } \mathcal{C}$$

is full. Denote $\mathcal{F}_{\mathcal{C}, \varphi, \tilde{\pi}}(\pi) = (\tilde{\pi} \otimes \pi) \circ \varphi$ by $\hat{\pi}$. Then

$$\hat{\pi} = (\tilde{\pi}_2 \otimes \mathcal{F}_{\mathcal{C}_1, \varphi_1, \tilde{\pi}_1}(\pi)) \circ \varphi_2 = \mathcal{F}_{\mathcal{C}_2, \varphi_2, \tilde{\pi}_2}(\mathcal{F}_{\mathcal{C}_1, \varphi_1, \tilde{\pi}_1}(\pi)).$$

Evidently, for any $\Lambda \in \text{Mor}(\pi_1, \pi_2)$, $\pi_i \in \text{Ob}(\text{Rep } \mathcal{A})$, one has

$$\mathcal{F}_{\mathcal{C}, \varphi, \tilde{\pi}}(\Lambda) = \mathbf{1} \otimes \mathbf{1} \otimes \Lambda = \mathcal{F}_{\mathcal{C}_2, \varphi_2, \tilde{\pi}_2}(\mathcal{F}_{\mathcal{C}_1, \varphi_1, \tilde{\pi}_1}(\Lambda)).$$

Hence, $\mathcal{F}_{\mathcal{C}, \varphi, \tilde{\pi}}$ is full as a composition of functors. \square

To give a definition of a $*$ -wild C^* -algebra one has to fix some C^* -algebra with a rather complicated category of representations. The group C^* -algebra of the free group in two generators, $C^*(\mathcal{F}_2)$, is a suitable candidate for this purpose, see [6].

Definition 2. A C^* -algebra \mathcal{A} is $*$ -wild if

$$C^*(\mathcal{F}_2) \prec \mathcal{A}.$$

By Proposition 1, a C^* -algebra majorizing a $*$ -wild C^* -algebra is $*$ -wild.

The following property of $*$ -wild algebras is obvious.

Proposition 2. *An extension of a $*$ -wild C^* -algebra is $*$ -wild.*

Proof. Let \mathcal{A} be a quotient of a $*$ -wild C^* -algebra \mathcal{B} and the majorization $\mathcal{B} \succ C^*(\mathcal{F}_2)$ be given by a nuclear C^* -algebra \mathcal{C} via a homomorphism

$$\varphi: \mathcal{B} \rightarrow \mathcal{C} \otimes C^*(\mathcal{F}_2)$$

and an irreducible representation $\tilde{\pi}: \mathcal{C} \rightarrow B(\mathcal{H})$. Denote by $\psi: \mathcal{A} \rightarrow \mathcal{B}$ the canonical homomorphism. Then the majorization $\mathcal{A} \succ C^*(\mathcal{F}_2)$ is defined by the triple $(\mathcal{C}, \varphi \circ \psi, \tilde{\pi} \circ \psi)$. \square

In the next Proposition we show that $*$ -wild C^* -algebras are not nuclear, see also [4].

Proposition 3. *Let \mathcal{A} be a $*$ -wild C^* -algebra. Then \mathcal{A} is not nuclear.*

Proof. It is sufficient to show that \mathcal{A} has a representation generating a non-hyperfinite factor, since any factor-representation of a nuclear C^* -algebra is hyperfinite. Since $C^*(\mathcal{F}_2) \prec \mathcal{A}$, one has the homomorphism

$$\varphi: \mathcal{A} \rightarrow \mathcal{C} \otimes C^*(\mathcal{F}_2),$$

where \mathcal{C} is a nuclear C^* -algebra, and the irreducible representation

$$\tilde{\pi}: \mathcal{C} \rightarrow B(\mathcal{H})$$

is as in Definition 1.

Consider the representation π of $C^*(\mathcal{F}_2)$ generating a non-hyperfinite factor. Put $(\tilde{\pi} \otimes \pi) \circ \varphi := \pi_1$ and note that

$$(\tilde{\pi} \otimes \pi)(\mathcal{C} \otimes C^*(\mathcal{F}_2))'' = B(\mathcal{H}) \otimes \pi(C^*(\mathcal{F}_2))''$$

is also a non-hyperfinite factor. Since the functor

$$F: \text{Rep } C^*(\mathcal{F}_2) \rightarrow \text{Rep } \mathcal{A}$$

defined in Definition 1 is full, we have

$$\pi_1(\mathcal{A})' = (\tilde{\pi} \otimes \pi)(\mathcal{C} \otimes C^*(\mathcal{F}_2))'.$$

Hence,

$$\pi_1(\mathcal{A})'' = (\tilde{\pi} \otimes \pi)(\mathcal{C} \otimes C^*(\mathcal{F}_2))''$$

is a non-hyperfinite factor. □


Similarly to the definition of a $*$ -wild C^* -algebra, one can define the notion of a $*$ -wild group, see [11].

Definition 3. A group G is called $*$ -wild iff its group C^* -algebra is $*$ -wild.

Note that there exist non-amenable but not $*$ -wild groups, see [11].

0.3. C^* -free products. Here we recall the definition of the free product of a family of C^* -algebras, see [13] for details.

Definition 4. If $(\mathcal{A}_i)_{i \in I}$ is a family of unital C^* -algebras, then their free product is a unique unital C^* -algebra $\mathcal{A} = *_{i \in I} \mathcal{A}_i$ and a unital $*$ -homomorphisms $\psi_i : \mathcal{A}_i \rightarrow \mathcal{A}$ such that, given any unital C^* -algebra \mathcal{B} and a unital $*$ -homomorphisms $\Phi_i : \mathcal{A}_i \rightarrow \mathcal{B}$, there exists a unique unital $*$ -homomorphism $\Phi = *_{i \in I} \Phi_i : \mathcal{A} \rightarrow \mathcal{B}$ making the following diagram commutative,



The definition of the free product of groups (algebras) is absolutely analogous. Moreover, it is easy to see that the free product of the family of group C^* -algebras is isomorphic to the group C^* -algebra of the free product of the corresponding family of groups. Indeed, let $(G_i)_{i \in I}$ be a family of groups. Then a unitary representation of the free product $*_{i \in I} G_i$ is determined uniquely by its restrictions to the groups G_i . Since unitary representations of a group are in one-to-one correspondence with $*$ -representations of its group C^* -algebra, by the universal property of the C^* -free product we get the required isomorphism,

$$C^*(*_{i \in I} G_i) \simeq *_{i \in I} C^*(G_i).$$

1. THE FREE PRODUCTS OF FINITE-DIMENSIONAL C^* -ALGEBRAS

In this section we show that the free product of finite-dimensional C^* -algebras is either of type 1 or is $*$ -wild. In fact, we give a criterion for $*$ -wildness of such free products. Throughout this section we use the old definition of majorization, i.e., $\mathcal{A} \succ \mathcal{B}$ if there exists $n \in \mathbb{N}$ and

$$\varphi: \mathcal{A} \rightarrow M_n(\mathcal{B}) \simeq M_n(\mathbb{C}) \otimes \mathcal{B}$$

such that the corresponding functor $\mathcal{F}_\varphi: \text{Rep} \mathcal{B} \rightarrow \text{Rep} \mathcal{A}$ is full. Here by \mathcal{F}_φ we mean $\mathcal{F}_{M_n(\mathbb{C}), \varphi, \text{id}_n}$ and

$$\text{id}_n: M_n(\mathbb{C}) \rightarrow M_n(\mathbb{C})$$

denotes the identical representation.

Since any finite-dimensional C^* -algebra is $*$ -isomorphic to the direct sum of matrix algebras over \mathbb{C} , see for example [10], our idea is to consider “elementary” free products of the form $\ast_{i \in I} M_{n_i}(\mathbb{C})$ for all possible values of $n_i \in \mathbb{N}$.

In the following, for $A \in M_k(\mathbb{C})$ and $B \in M_n(\mathbb{C})$, by $A \oplus B$ we mean the $n+k \times n+k$ block-diagonal matrix of the form

$$A \oplus B = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}.$$

If $A = (a_{ij})_{i,j=1}^k \in M_k(\mathbb{C})$ and the element b belongs to some $*$ -algebra \mathcal{B} , we denote by $A \otimes b$ the element of $M_k(\mathcal{B})$ of the form $(b a_{ij})_{i,j=1}^k$.

1. Firstly we prove the $*$ -wildness of the C^* -free product of two matrix-algebras with $k > 1$ and $n > 1$.

Proposition 4. *The C^* -algebra $M_k(\mathbb{C}) \ast M_n(\mathbb{C})$ is $*$ -wild for every $k > 1, n > 1$.*

Proof. Let $(e_{ij}^{(k)})_{i,j=1}^k$ and $(e_{ij}^{(n)})_{i,j=1}^n$ be the matrix units of the algebras $M_k(\mathbb{C})$ and $M_n(\mathbb{C})$, respectively. Let u, v be the generators and e be the unit of \mathcal{F}_2 . Set $L := \text{LCM}(n, k)$. We shall construct a $*$ -homomorphism $\psi: M_k(\mathbb{C}) \ast M_n(\mathbb{C}) \rightarrow M_L(C^*(\mathcal{F}_2))$ by the following rule:

$$\begin{aligned} \psi(e_{12}^{(k)}) &= e_{12}^{(k)} \otimes v \oplus \bigoplus_1^{\frac{1}{k}L-1} e_{12}^{(k)} \otimes e, \\ \psi(e_{ii+1}^{(k+1)}) &= \bigoplus_1^{\frac{1}{k}L} e_{ii+1}^{(k)} \otimes e, \quad \text{for } 2 \leq i \leq k; \\ \psi(e_{12}^{(n)}) &= T(e_{12}^{(n)} \otimes u \oplus \bigoplus_1^{\frac{1}{n}L-1} e_{12}^{(n)} \otimes e)T^*, \\ \psi(e_{ii+1}^{(n)}) &= T(\bigoplus_1^{\frac{1}{n}L} e_{ii+1}^{(n)} \otimes e)T^*, \quad \text{for } 2 \leq i \leq n. \end{aligned}$$

Here

$$T = \begin{pmatrix} 1 & -\tan x \\ \tan x & 1 \end{pmatrix} \otimes e \oplus I_{L-2} \otimes e, \quad x \in (0, \frac{\pi}{2}),$$

I_k denotes the $k \times k$ identity matrix. In the right-hand sides of the above formulas, by $e_{ij}^{(k)}, (e_{ij}^{(n)})$ we mean concrete $k \times k$ ($n \times n$ resp.) matrices.

Consider a representation π of \mathcal{F}_2 and any operator $A = (a_{rs})_{r,s=1}^L$ commuting with the operators $(\text{id}_L \otimes \pi)(\psi(e_{ij}^{(k)}))$, $i, j = 1, \dots, k$, and $(\text{id}_L \otimes \pi)(\psi(e_{ij}^{(n)}))$, $i, j = 1, \dots, n$.

It is just a routine to verify that

$$[A, (\text{id}_L \otimes \pi)(\psi(e_{ij}^{(k)}))] = 0, \quad i, j = 1, \dots, k,$$

and

$$[A, (\text{id}_L \otimes \pi)(\psi(e_{ij}^n))] = 0, \quad i, j = 1 \dots, n,$$

implies $A = \text{diag}(c, c, \dots, c)$ with $[c, \pi(u)] = [c, \pi(v)] = 0$, proving the fullness of the functor F_ψ . \square

2. In the following proposition we study the free products of \mathbb{C}^k , $k \geq 2$.

Proposition 5. *The C^* -algebra $\mathbb{C}^n * \mathbb{C}^k$ is $*$ -wild iff $n \geq 2$, $k \geq 3$, $n, k \in \mathbb{N}$. The C^* -algebra $\mathbb{C}^2 * \mathbb{C}^2 * \mathbb{C}^2$ is $*$ -wild.*

Proof. Since for any family of groups $\{G_i, i \in I\}$ one has $C^*(\ast_{i \in I} G_i) = \ast_{i \in I} C^*(G_i)$, we have the following isomorphisms:

$$\begin{aligned} \mathbb{C}^n * \mathbb{C}^k &\simeq C^*(\mathbb{Z}_n * \mathbb{Z}_k), \\ \mathbb{C}^2 * \mathbb{C}^2 * \mathbb{C}^2 &\simeq C^*(\mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2). \end{aligned}$$

As it was proved in [11], $C^*(\mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2)$ is $*$ -wild and $C^*(\mathbb{Z}_n * \mathbb{Z}_k)$ is $*$ -wild if and only if $n \geq 2$, $k \geq 3$. This yields the statement of the proposition. \square

3. It remains to consider the product $\mathbb{C}^k * M_n(\mathbb{C})$.

Proposition 6. *The C^* -algebra $\mathbb{C}^n * M_k(\mathbb{C})$ is $*$ -wild iff $n \geq 2$, $k \geq 2$, $n, k \in \mathbb{N}$.*

Proof. Let $(e_{ij})_{i,j=1}^k$ be matrix units of the algebra $M_k(\mathbb{C})$ and let z be a generator of $C^*(\mathbb{Z}_n) \simeq \mathbb{C}^n$. It is known, see [11], that the group

$$\mathbb{Z}_2 * \mathbb{Z}_3 = \langle u, v | v^2 = u^3 = e \rangle$$

is $*$ -wild. To prove the statement of the proposition we are going to construct a $*$ -homomorphism

$$\varphi : \mathbb{C}^n * M_k(\mathbb{C}) \rightarrow M_l(C^*(\mathbb{Z}_2 * \mathbb{Z}_3)), \quad \text{for sufficient large } l \in \mathbb{N}$$

such that the corresponding functor F_φ is full.

a) Let $n = k = 2$. Define

$$\varphi : \mathbb{C}^2 * M_2(\mathbb{C}) \rightarrow M_4(C^*(\mathbb{Z}_2 * \mathbb{Z}_3))$$

by the following rule:

$$\begin{aligned} \varphi(z) &= \begin{pmatrix} v & 0 & 0 & 0 \\ 0 & 0 & e & 0 \\ 0 & e & 0 & 0 \\ 0 & 0 & 0 & -e \end{pmatrix}, \quad \varphi(e_{12}) = T \begin{pmatrix} 0 & v & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & u \\ 0 & 0 & 0 & 0 \end{pmatrix} T^*, \\ \text{here } T &= \begin{pmatrix} e & 0 & 0 & 0 \\ 0 & e & 0 & 0 \\ 0 & 0 & e & -e \tan x \\ 0 & 0 & e \tan x & e \end{pmatrix}, \quad x \in (0, \pi/2). \end{aligned}$$

One can directly check that the functor F_φ generated by φ is full. Thus, $\mathbb{C}^2 * M_2(\mathbb{C})$ is $*$ -wild.

b) Let $n = 2$, $k \geq 3$. In this case we construct

$$\varphi : \mathbb{C}^2 * M_k(\mathbb{C}) \rightarrow M_k(C^*(\mathbb{Z}_2 * \mathbb{Z}_3))$$

as follows:

$$\varphi(z) = \begin{pmatrix} 0 & u^{-1} & 0 \\ u & 0 & 0 \\ 0 & 0 & v \end{pmatrix} \oplus (I_{k-3} \otimes e), \quad \varphi(e_{ij}) = e_{ij} \otimes e, \quad i, j = 1, \dots, k.$$

To prove that F_φ is full we fix an arbitrary representation π of $\mathbb{Z}_2 * \mathbb{Z}_3$. Let $A = (a_{rs})_{r,s=1}^2$ commute with $(\text{id}_k \otimes \pi)(\varphi(e_{ij}))$, $i, j = 1, \dots, k$, and $(\text{id}_k \otimes \pi)(\varphi(z))$. Then

$$[A, (\text{id}_k \otimes \pi)(\varphi(e_{ij}))] = 0, \quad i, j = 1, \dots, k,$$

implies $A = \text{diag}(a_{11}, \dots, a_{11})$, and

$$[A, (\text{id}_k \otimes \pi)(\varphi(z))] = 0$$

gives $[a_{11}, \pi(u)] = [a_{11}, \pi(v)] = 0$, proving the fullness of the functor F_φ .

c) Let $n \geq 3, k \geq 2$. Define a $*$ -homomorphism

$$\varphi : \mathbb{C}^n * M_k(\mathbb{C}) \rightarrow M_k(C^*(\mathbb{Z}_n * \mathbb{Z}_n))$$

by the following rule:

$$\varphi(z) = \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix} \oplus (I_{k-2} \otimes e), \quad \varphi(e_{ij}) = e_{ij} \otimes e, \quad i, j = 1, \dots, k.$$

One can directly check, using similar arguments as in item **b**), that the functor F_φ generated by φ is full. Since $C^*(\mathbb{Z}_n * \mathbb{Z}_n)$ is $*$ -wild, see [11], $\mathbb{C}^n * M_k(\mathbb{C})$ is also $*$ -wild for $n \geq 3, k \geq 2$. \square

Proposition 7. *Let \mathcal{A} be a $*$ -wild C^* -algebra and \mathcal{B} be a finite-dimensional C^* -algebra. Then $\mathcal{A} * \mathcal{B}$ is $*$ -wild.*

Proof. Evidently, it is sufficient to consider only the case $\mathcal{B} = M_n(\mathbb{C}), n \in \mathbb{N}$. Construct a homomorphism

$$\phi : \mathcal{A} * \mathcal{B} \rightarrow \mathcal{B} \otimes \mathcal{A}$$

defined by $\phi(a) = 1 \otimes a, a \in \mathcal{A}, \phi(b) = b \otimes 1, b \in \mathcal{B}$.

Let the majorization $\mathcal{A} \succ C^*(\mathcal{F}_2)$ be given by the homomorphism

$$\psi : \mathcal{A} \rightarrow \mathcal{C} \otimes C^*(\mathcal{F}_2), \quad \hat{\pi} : \mathcal{C} \rightarrow B(\mathcal{K}),$$

where \mathcal{C} is a nuclear C^* -algebra and $\hat{\pi}$ is its irreducible representation. Then the majorization $\mathcal{A} * \mathcal{B}$ is given by the homomorphism

$$(\text{id}_{\mathcal{B}} \otimes \psi) \circ \phi : \mathcal{A} * \mathcal{B} \rightarrow \mathcal{B} \otimes \mathcal{C} \otimes C^*(\mathcal{F}_2)$$

and the irreducible representation $\text{id}_{\mathcal{B}} \otimes \hat{\pi} : \mathcal{B} \otimes \mathcal{C} \rightarrow B(\mathbb{C}^n \otimes \mathcal{K})$. \square

Remark 3. The statement of the proposition above remains true if we suppose \mathcal{B} to be a nuclear C^* -algebra.

Now we combine the results of the previous propositions to get a criterion for $*$ -wildness of the C^* -free products of finite-dimensional C^* -algebras.

Theorem 1. *The C^* -free product $*_{i \in I} \mathcal{A}_i$ of a family of finite-dimensional C^* -algebras $(\mathcal{A}_i)_{i \in I}$ is $*$ -wild if and only if one of the following conditions holds:*

1. *there exist $i_1, i_2 \in I$ such that \mathcal{A}_{i_1} has $M_k(\mathbb{C})$ and \mathcal{A}_{i_2} has $M_n(\mathbb{C})$ as a direct summand with $k > 1, n > 1$;*
2. *there exist $i_1, i_2 \in I$ such that \mathcal{A}_{i_1} has \mathbb{C}^n and \mathcal{A}_{i_2} has \mathbb{C}^k as a direct summand with $n \geq 2, k \geq 3$;*
3. *there exist $i_1, i_2, i_3 \in I$ such that \mathcal{A}_{i_1} has $\mathbb{C}^2, \mathcal{A}_{i_2}$ has \mathbb{C}^2 , and \mathcal{A}_{i_3} has \mathbb{C}^2 as a direct summand;*

4. there exist $i_1, i_2 \in I$ such that \mathcal{A}_{i_1} has \mathbb{C}^n and \mathcal{A}_{i_2} has $M_k(\mathbb{C})$ as a direct summand with $n \geq 2, k \geq 2$.

Proof. In fact, the $*$ -wildness of the algebras specified in items of the statement follows from Propositions 2,4,5,6,7. The C^* -algebras products which are left aside in the list of concerned C^* -free products are the following:

1. $\mathbb{C}^2 * \mathbb{C}^2$,
2. the C^* -free product of any number of copies of \mathbb{C} and one C^* -algebra $M_n(\mathbb{C})$; evidently, such a product is isomorphic to $M_n(\mathbb{C})$.

The $*$ -representations of the C^* -algebra $\mathbb{C}^2 * \mathbb{C}^2$ are described in [11] and this C^* -algebra is of type **1**. □

Corollary 1. $G_1 * G_2$, where $G_i \neq \langle e \rangle, i = 1, 2$, are finite groups, is $*$ -wild unless $G_i = \mathbb{Z}_2, i = 1, 2$.

2. $*$ -WILDNESS OF THE FREE PRODUCT OF A NUCLEAR NON-COMMUTATIVE C^* -ALGEBRA AND $C(\mathbf{T})$

2.1. Recall that the C^* -algebra generated by a single unitary element is isomorphic to the algebra $C(\mathbf{T})$ of the continuous functions on the one-dimensional torus \mathbf{T} .

Theorem 2. Let \mathcal{A} be non-commutative nuclear C^* -algebra having a non-trivial projection, say p . Then the C^* -algebraic free product $\mathcal{A} * C(\mathbf{T})$ is $*$ -wild.

Proof. In the following we denote by u the standard generator of $C(\mathbf{T})$, and denote by u_1, u_2 the standard free generators of $C^*(\mathcal{F}_2)$. Then the needed majorization is given by the homomorphism

$$\varphi: \mathcal{A} * C(\mathbf{T}) \rightarrow \mathcal{A} \otimes C^*(\mathcal{F}_2)$$

defined by

$$\varphi(x) = x \otimes 1, \quad x \in \mathcal{A}, \quad \varphi(u) = p \otimes u_1 + (1 - p) \otimes u_2,$$

and the irreducible representation $\tilde{\pi}: \mathcal{A} \rightarrow B(\mathcal{H})$, where $\dim \mathcal{H} \geq 2$ such that $\tilde{\pi}(p)$ is non-trivial.

We prove that the functor

$$\mathcal{F}_{\mathcal{A}, \varphi, \tilde{\pi}}: \text{Rep} C^*(\mathcal{F}_2) \rightarrow \text{Rep} \mathcal{A} * C(\mathbf{T})$$

is full.

Put $\mathcal{F}_{\mathcal{A}, \varphi, \tilde{\pi}}(\pi) := \pi_1, \tilde{\pi}(p) := P$, and $\pi(u_i) = U_i, i = 1, 2$. Then

$$\pi_1(u) = P \otimes U_1 P + (1 - P) \otimes U_2.$$

Let $\Lambda \in \pi_1(\mathcal{A} * C(\mathbf{T}))'$, then it follows from the irreducibility of $\tilde{\pi}$ and the inclusion $\mathbf{1} \otimes \tilde{\pi}(\mathcal{A}) \subset \pi_1(\mathcal{A} * C(\mathbf{T}))$ that $\Lambda = \mathbf{1} \otimes \Lambda_1$. We now show that $\Lambda_1 \in \pi(C^*(\mathcal{F}_2))'$. In fact, from

$$\pi_1(up) = P \otimes U_1, \quad \pi_1(u(1 - p)) = (1 - P) \otimes U_2,$$

one has

$$P \otimes \Lambda_1 U_1^\varepsilon = P \otimes U_1^\varepsilon \Lambda_1 \otimes P, \quad (1 - P) \otimes \Lambda_1 U_2^\varepsilon = (1 - P) \otimes U_2^\varepsilon \Lambda_1,$$

hence $\Lambda_1 U_i^\varepsilon = U_i^\varepsilon \Lambda_1, \varepsilon \in \{1, *\}, i = 1, 2$, and $\Lambda_1 \in \pi(C^*(\mathcal{F}_2))'$. □

Remark 4.

1. In fact, the claim of the proposition remains true if we suppose that \mathcal{A} has an irreducible representation $\hat{\pi}$ such that $\hat{\pi}(\mathcal{A})$ is nuclear and contains the non-trivial projection p . Indeed, in this case one can replace

$$\varphi: \mathcal{A} * C(\mathbf{T}) \rightarrow \mathcal{A} \otimes C^*(\mathcal{F}_2)$$

with

$$\varphi: \mathcal{A} * C(\mathbf{T}) \rightarrow \widehat{\pi}(\mathcal{A}) \otimes C^*(\mathcal{F}_2)$$

defined as in the proof of Theorem 2 and put $\tilde{\pi}$ to be the identical representation of $\widehat{\pi}(\mathcal{A})$.

2. It follows from the results of [11, 8] that the C^* -algebra $C(\mathbf{T})$ can be replaced with $C([a, b])$ or $C_0(\mathbb{R})$.

2.2. In the following propositions we present some applications of the above result.

Proposition 8. *Let B be a C^* -algebra generated by partial isometries s_1, \dots, s_d satisfying the relation*

$$\sum_{i=1}^d \alpha_i s_i s_i^* = 1.$$

Suppose that there exist irreducible projections P_i , $i = 1, \dots, d$, acting on a finite-dimensional Hilbert space \mathcal{H} , $\dim \mathcal{H} \geq 2$, satisfying the relation

$$(1) \quad \sum_{i=1}^d \alpha_i P_i = 1.$$

Then B is $$ -wild.*

Proof. Consider the C^* -algebra \mathcal{P} generated by the projections P_i , $i = 1, \dots, d$. Construct the homomorphism

$$\varphi: B \rightarrow \mathcal{P} * C(\mathbf{T})$$

by the formulas

$$\varphi(s_i) = uP_i, \quad i = 1, \dots, d.$$

Evidently, φ is surjective, since $\varphi(s_i^* s_i) = p_i$ and $\varphi(\sum_i \alpha_i s_i) = u$. By Theorem 2 the C^* -algebra $\mathcal{P} * C^*(\mathbf{T})$ is $*$ -wild, therefore Proposition 2 implies that B has the same property. \square

Consider the C^* -algebra \mathcal{A}_n^q generated by the isometries s_i , $i = 1, \dots, n$, satisfying the q -commutation relations of the form

$$s_i s_j = q s_j s_i, \quad |q| = 1, \quad i > j.$$

Below we show that \mathcal{A}_n^q is $*$ -wild for any q specified above.

Remark 5.

1. It is interesting to note that if we consider the C^* -algebra $\widetilde{\mathcal{B}}_n^q$ generated by isometries s_i satisfying relations of the form

$$s_j^* s_i = q s_i s_j^*, \quad |q| = 1, \quad i > j,$$

then relations $s_i s_j = q s_j s_i$, $i > j$, hold automatically, see [12], and \mathcal{B}_q is nuclear, so it is not $*$ -wild.

2. If $|q| \neq 1$, then there are no isometries satisfying the relation $s_i s_j = q s_j s_i$. Indeed, let s_i, s_j act on a Hilbert space \mathcal{H} . Then for any $x \in \mathcal{H}$ one has

$$\|x\| = \|s_i s_j x\| = |q|^2 \|s_j s_i x\| = |q| \|x\|.$$

Below we denote by \mathcal{O}_2 the Cuntz algebra with two generators, see [2]. Namely

$$\mathcal{O}_2 = C^*(t_1, t_2 \mid t_i^* t_i = 1, i = 1, 2, t_1 t_1^* + t_2 t_2^* = 1).$$

Proposition 9. *The C^* -algebra \mathcal{A}_n^q is $*$ -wild for any q , $|q| = 1$.*

Proof. We show that $\mathcal{O}_2 * C([0, 1]) \prec \mathcal{A}_n^q$.

Let us construct the homomorphism

$$\varphi: \mathcal{A}_n^q \rightarrow \mathcal{B}^{\otimes n-1} \otimes (\mathcal{O}_2 * C([0, 1])),$$

where

$$\mathcal{B} = C^*(s, u \mid s^*s = 1, u^*u = uu^* = 1, us = qsu).$$

Evidently \mathcal{B} is nuclear, since it is the crossed product $\mathcal{B} = \mathcal{T}(C(\mathbf{T})) \rtimes \mathbb{Z}$, where $\mathcal{T}(C(\mathbf{T}))$ is the Toeplitz C^* -algebra. Let $c, 0 < c < 1$, denote the generator of $C([0, 1])$ and let t_1, t_2 be generators of \mathcal{O}_2 . Put

$$a_1 := t_1c, \quad a_2 := t_2(1 - c).$$

It is easy to verify that the following relations are satisfied

$$(2) \quad a_1^*a_1 + a_2^*a_2 = 1, \quad a_2^*a_1 = 0.$$

Then φ is defined by following formulas:

$$\begin{aligned} \varphi(s_i) &= \bigotimes_{j < i} u \otimes s \otimes \bigotimes_{i < j \leq n} 1, \quad i = 1, \dots, n - 1, \\ \varphi(s_n) &= \bigotimes_{1 \leq j \leq n-2} u \otimes (u \otimes a_1 + su \otimes a_2). \end{aligned}$$

Further we fix the irreducible representation $\tilde{\pi}$ of \mathcal{B} acting on $\mathcal{K} = l_2(\mathbb{N})$,

$$(3) \quad \tilde{\pi}(s) = S, \quad \tilde{\pi}(u) = D(q),$$

where

$$Se_n = e_{n+1}, \quad D(q)e_n = q^{n-1}e_n, \quad n \in \mathbb{N}.$$

Then the majorization \mathcal{A}_n^q is given by functor the $\mathcal{F}_{\mathcal{B}^{\otimes n-1}, \varphi, \tilde{\pi}}$.

The proof of the fullness of $\mathcal{F}_{\mathcal{B}^{\otimes n-1}, \varphi, \tilde{\pi}}$ is essentially the same as in Theorem 2. For $\pi \in \text{ObRep } \mathcal{O}_2 * C([0, 1])$, put $\mathcal{F}_{\mathcal{B}^{\otimes n-1}, \varphi, \tilde{\pi}}(\pi) = \pi_1$ and denote $\pi(a_i)$ by A_i , $\pi(t_i) = T_i$, $\pi(c) = C$.

Then the equalities $C^2 = A_1^*A_1$ and $T_i = A_iC^{-1}$, $i = 1, 2$, imply that

$$\{A_i, A_i^*, i = 1, 2\}' = \{T_i, T_i^*, C, i = 1, 2\}' = \pi(\mathcal{O}_2 * C([0, 1]))'.$$

So one has to show that any $\Lambda \in \pi_1(\mathcal{A}_n^q)'$ has the form $\Lambda = \mathbf{1}^{\otimes n-1} \otimes \Lambda_1$ with $\Lambda_1 \in \{A_i, A_i^*, i = 1, 2\}'$. □

The following corollary is immediate.

Corollary 2. *The C^* -algebra \mathcal{A}_n^q is not nuclear.*

Acknowledgments. The work on this paper was completed during a visit of the third and the fourth authors to the Institut für Angewandte Mathematik, Universität Bonn, the hospitality and fruitful atmosphere wherein is gratefully acknowledged. The visit was supported by the DFG grant, project no. 436 UKR 113/71/0-1.

REFERENCES

1. C. A. Berger, L. A. Coburn, A. Lebow, *Representations and index theory for C^* -algebras generated by commuting isometries*, J. Funct. Anal. **27** (1978), 51–99.
2. J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57** (1977), 173–185.
3. J. Dixmier, *Les C^* -Algèbres et Leurs Représentations*, Gauthier-Villars Editeur, Paris, 1969.
4. P. E. T. Jørgensen, D. P. Proskurin, Yu. S. Samoilenko, *On C^* -algebras generated by q -commuting isometries*, J. Phys. A **38** (2005), no. 12, 2669–2680.
5. A. Ya. Khelemskii, *Banach Algebras and Poly-Normed Algebras: General Theory, Representations, Homologies*, Nauka, Moscow, 1989. (Russian)
6. S. A. Kruglyak, Yu. S. Samoilenko, *Unitary equivalence of sets of selfadjoint operators*, Funktsional. Anal. i Prilozhen. **14** (1980), no. 1, 60–62. (Russian)

7. S. A. Kruglyak, Yu. S. Samoilenko, *On complexity of description of representation of *-algebras generated by idempotents*, Proc. Amer. Math. Soc. **128** (2002), no. 6, 1655–1664.
8. V. Mazorchuk, L. Turowska, **-representation type of *-doubles of finite-dimensional algebras*, Proc. Edinb. Math. Soc. (2) **47** (2004), no. 3, 669–678.
9. V. Mazorchuk, L. Turowska, *Radical *-doubles of finite-dimensional algebras*, Linear Algebra Appl. **390** (2004), 293–309.
10. G. J. Murphy, *C^* -Algebras and Operator Theory*, Academic Press, London, 1990.
11. V. Ostrovskyi and Yu. Samoilenko, *Introduction to the Theory of Representations of Finitely Presented *-Algebras. I. Representations by Bounded Operators*, The Gordon and Breach Publishing group, London, 1999.
12. D. Proskurin, *Stability of a special class of q_{ij} -CCR and extensions of higher-dimensional noncommutative tori*, Lett. Math. Phys. **52** (2000), no. 2, 165–175.
13. D. V. Voiculescu, K. J. Dykema, A. Nica, *Free Random Variables: A Noncommutative Probability Approach to Free Products With Applications to Random Matrices, Operator Algebras, and Harmonic Analysis on Free Groups*, CRM Monograph Series, vol. 1, Amer. Math. Soc., Providence, R. I., 1992, 1992.

INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, WEGELERSTR. 6, D-53115, BONN, GERMANY; SFB 611, BONN, GERMANY; BiBoS, BIELEFELD, GERMANY; IZKS; CERFIM, LOCARNO, SWITZERLAND; ACCADEMIA DI ARCHITETTURA, MENDRISIO, SWITZERLAND

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY, SE-41296, GÖTEBORG, SWEDEN

DEPARTMENT OF CYBERNETICS, KYIV TARAS SHEVCHENKO NATIONAL UNIVERSITY, 64 VOLODYMYR-S'KA, KYIV, 01033, UKRAINE

E-mail address: `prosk@univ.kiev.ua`

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS'KA, KYIV, 01601, UKRAINE

E-mail address: `yurii.sam@imath.kiev.ua`

Received 26/07/2006