

ANOTHER FORM OF SEPARATION AXIOMS

S. ATHISAYA PONMANI AND M. LELLIS THIVAGAR

ABSTRACT. It is the object of this paper to introduce the $(1, 2)^*$ pre- D_k axioms for $k = 0, 1, 2$.

1. INTRODUCTION

In 1982, Mashhour et al [9] introduced the notion of pre-open sets and the pre-closed sets were defined in [4]. Ashish Kar and Bhattacharya [2], in 1990, continued their work on pre-open sets and offered another set of separation axioms analogous to the semi separation axioms defined by Maheshwari and Prasad [8]. Caldas [10] defined a new class of sets called semi-Difference (briefly sD) sets by using the semi-open sets [5], and introduced the semi- D_i spaces for $i = 0, 1, 2$.

The purpose of this paper is to introduce some pre-separation axioms in bitopological spaces. To define and investigate the axioms we use the $(1, 2)^*$ pre-open sets [6] introduced by Lellis Thivagar and Ravi. We call these axioms as $(1, 2)^*$ pre- T_0 , $(1, 2)^*$ pre- T_1 and $(1, 2)^*$ pre- T_2 . We also define $(1, 2)^*$ pre-Difference sets and utilize them to define the $(1, 2)^*$ pre- D_i , $i = 0, 1, 2$, axioms. We prove a bitopological space is $(1, 2)^*$ pre- D_1 if and only if it is $(1, 2)^*$ pre- D_2 .

We recall some definitions and concepts which are useful in the following sections.

2. PRELIMINARIES

In this section unless it is explicitly stated X is a topological space (X, τ) . For a $A \subset X$, the interior and closure of A in X are denoted by $int(A)$ and $cl(A)$ respectively.

Definition 2.1. A space X is called

- (i). Pre- T_0 [2] iff to each pair of distinct points x, y in X , there exists a pre-open set containing one of the points but not the other.
- (ii). Pre- T_1 [2] iff to each pair of distinct points x, y of X , there exists a pair of pre-open sets one containing x but not y and other containing y but not x .
- (iii). Pre- T_2 [2] iff to each pair of distinct points x, y of X , there exists a pair of disjoint pre-open sets one containing x and the other containing y .

Definition 2.2. A subset A of X is called a semi-Difference set (in short sD -set) if there are two semi-open sets O_1, O_2 in X such that $O_1 \neq X$ and $A = O_1 \setminus O_2$.

Definition 2.3. A space X is called

- (i). Semi- D_0 if for $x, y \in X$, $x \neq y$, there exists a sD -set of X containing one of x and y but not the other.
- (ii). Semi- D_1 if for $x, y \in X$, $x \neq y$, there exists a pair of sD -sets one containing x but not y and the other containing y but not x .

2000 *Mathematics Subject Classification.* 54C55.

Key words and phrases. $(1, 2)^*$ pre- T_k , $(1, 2)^*$ pre- D_k , $k = 0, 1, 2$.

The first author wishes to acknowledge the support of University Grants Commission, New Delhi, India, FIP-X Plan.

- (iii). Semi- D_2 if for $x, y \in X$, $x \neq y$, there exist disjoint sD -sets S_1 and S_2 such that $x \in S_1$ and $y \in S_2$.

A subset A of a bitopological space (X, τ_1, τ_2) is called $\tau_{1,2}$ -open [6] if $A = S_1 \cup S_2$ where $S_1 \in \tau_1$ and $S_2 \in \tau_2$, and $\tau_{1,2}$ -closed if A^c is $\tau_{1,2}$ -open in X . We write $\tau_1\tau_2$ -interior of A and $\tau_1\tau_2$ -closure of A , in short form, $\tau_1\tau_2$ - $int(A)$ and $\tau_1\tau_2$ - $cl(A)$ respectively. $\tau_1\tau_2$ - $int(A)$ is the union of all $\tau_{1,2}$ -open sets contained in A and $\tau_1\tau_2$ - $cl(A)$ is the intersection of all $\tau_{1,2}$ -closed sets containing A .

Definition 2.4. A subset A of X is called $(1, 2)^*$ -pre-open [6] if $A \subset \tau_1\tau_2$ - $int(\tau_1\tau_2$ - $cl(A))$ and $(1, 2)^*$ -pre-closed if its complement in X is $(1, 2)^*$ -pre-open. Or equivalently, $\tau_1\tau_2$ - $cl(\tau_1\tau_2$ - $int(A)) \subset A$.

The family of all $(1, 2)^*$ -pre-open sets of X is denoted by $(1, 2)^*$ - $PO(X)$. $(1, 2)^*$ -pre-closure of A denoted by $(1, 2)^*$ - $pcl(A)$ is the intersection of all $(1, 2)^*$ -pre-closed sets containing A .

A subset A of X is $(1, 2)^*$ -pre-closed if and only if $(1, 2)^*$ - $pcl(A) = A$. Note that every $\tau_{1,2}$ -open set is $(1, 2)^*$ -pre-open.

Definition 2.5. A map $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is called

- (i). $(1, 2)^*$ -pre-irresolute if the inverse image of every $(1, 2)^*$ -pre-open set in Y is $(1, 2)^*$ -pre-open in X .
- (ii). Strongly $(1, 2)^*$ -pre-open [6] if the image of every $(1, 2)^*$ -pre-open set in X is $(1, 2)^*$ -pre-open in Y .

In the next section we introduce the $(1, 2)^*$ -pre- T_k -spaces for $k = 0, 1, 2$. In the following sections by X and Y we mean bitopological spaces (X, τ_1, τ_2) and (Y, σ_1, σ_2) respectively.

3. $(1, 2)^*$ -PRE- T_k -SPACES

Definition 3.1. A bitopological space X is said to be $(1, 2)^*$ -pre- T_0 iff for $x, y \in X$, $x \neq y$, there exists an $(1, 2)^*$ -pre-open set containing only one of x and y but not the other.

Now we proceed to prove that every bitopological space is $(1, 2)^*$ -pre- T_0 .

Lemma 3.2. *If for some $x \in X$, $\{x\}$ is $(1, 2)^*$ -pre-open then $x \notin (1, 2)^*$ - $pcl(\{y\})$ for all $y \neq x$.*

Proof. If $\{x\}$ is $(1, 2)^*$ -pre-open for some $x \in X$, then $X \setminus \{x\}$ is $(1, 2)^*$ -pre-closed and $x \notin X \setminus \{x\}$. If $x \in (1, 2)^*$ - $pcl(\{y\})$ for some $y \neq x$, then x, y both are in all the $(1, 2)^*$ -pre-closed sets containing y which implies that $x \in X \setminus \{x\}$ which is not true. Therefore, $x \notin (1, 2)^*$ - $pcl(\{y\})$. \square

Theorem 3.3. *In a space X , distinct points have distinct $(1, 2)^*$ -pre-closures.*

Proof. Let $x, y \in X$, $x \neq y$. Take $A = \{x\}^c$. Then $\tau_1\tau_2$ - $cl(A) = A$ or X .

Case (a). If $\tau_1\tau_2$ - $cl(A) = A$, then A is $\tau_{1,2}$ -closed and hence $(1, 2)^*$ -pre-closed. Then $X \setminus A = \{x\}$ is $(1, 2)^*$ -pre-open, not containing y . Therefore, by Lemma 3.2, $x \notin (1, 2)^*$ - $pcl(\{y\})$ and $y \in (1, 2)^*$ - $pcl(\{y\})$ which implies that

$$(1, 2)^*pcl(\{x\}) \quad \text{and} \quad (1, 2)^*pcl(\{y\})$$

are distinct.

Case (b). If $\tau_1\tau_2$ - $cl(A) = X$, then A is $(1, 2)^*$ -pre-open and hence $\{x\}$ is $(1, 2)^*$ -pre-closed which shows that $(1, 2)^*$ - $pcl(\{x\}) = \{x\}$ which is not equal to $(1, 2)^*$ - $pcl(\{y\})$. \square

Theorem 3.4. *In a space X , if distinct points have distinct $(1, 2)^*$ pre-closures then X is $(1, 2)^*$ pre- T_0 .*

Proof. Let $x, y \in X$, $x \neq y$. Then $(1, 2)^*pcl(\{x\})$ is not equal to $(1, 2)^*pcl(\{y\})$. Then there exists $z \in X$ such that $z \in (1, 2)^*pcl(\{x\})$ but $z \notin (1, 2)^*pcl(\{y\})$ or $z \in (1, 2)^*pcl(\{y\})$ but $z \notin (1, 2)^*pcl(\{x\})$. Without loss of generality, let $z \in (1, 2)^*pcl(\{x\})$ but $z \notin (1, 2)^*pcl(\{y\})$. If $x \in (1, 2)^*pcl(\{y\})$, then $(1, 2)^*pcl(\{x\})$ is contained in $(1, 2)^*pcl(\{y\})$ and therefore, $z \in (1, 2)^*pcl(\{y\})$, which is a contradiction. Thus we get $x \notin (1, 2)^*pcl(\{y\})$. This implies that $x \in (1, 2)^*pcl(\{y\}^c)$. Therefore, X is $(1, 2)^*$ pre- T_0 . \square

Theorem 3.5. *Every bitopological space is $(1, 2)^*$ pre- T_0 .*

Proof. Follows from Theorem 3.3 and Theorem 3.4. \square

Remark 3.6. It is observed that every T_0 -space is pre- T_0 but not the converse [2]. Here we note that if a space X is T_0 with respect to τ_1 or τ_2 then X is $(1, 2)^*$ pre- T_0 . But if X is $(1, 2)^*$ pre- T_0 , it is not necessary that (X, τ_1) is T_0 or (X, τ_2) is T_0 , as shown in the following example.

Example 3.7. Let $X = \{a, b, c\}$. $\tau_1 = \{\emptyset, \{a, b\}, X\}$, $\tau_2 = \{\emptyset, \{b, c\}, X\}$. Then X is $(1, 2)^*$ pre- T_0 but both (X, τ_1) and (X, τ_2) are T_0 .

Definition 3.8. A space X is called $(1, 2)^*$ pre- T_1 iff for $x, y \in X$, $x \neq y$, there exist $U, V \in (1, 2)^*PO(X)$ such that $x \in U$, $y \notin U$ and $y \in V$, $x \notin V$.

Remark 3.9. It is obvious that every $(1, 2)^*$ pre- T_1 space is $(1, 2)^*$ pre- T_0 but the converse is not true in general as illustrated in the next example.

Example 3.10. Let $X = \{a, b, c\}$. $\tau_1 = \{\emptyset, \{a\}, X\}$, $\tau_2 = \{\emptyset, \{b\}, X\}$. Then X is $(1, 2)^*$ pre- T_0 but not $(1, 2)^*$ pre- T_1 .

Theorem 3.11. *In a space X , the following statements are equivalent.*

- (i). X is $(1, 2)^*$ pre- T_1 .
- (ii). For each $x \in X$, $\{x\}$ is $(1, 2)^*$ pre-closed in X .
- (iii). Each subset of X is the intersection of all $(1, 2)^*$ pre-open sets containing it.
- (iv). The intersection of all $(1, 2)^*$ pre-open sets containing the point $x \in X$ is $\{x\}$.

Proof. (i) \Rightarrow (ii).

Let $x \in X$. If $y \in X$ and $x \neq y$ then there exists an $(1, 2)^*$ pre-open set U_y such that $y \in U_y$. Hence $y \in U_y \subset \{x\}^c$. Therefore, $\{x\}^c = \bigcup \{U_y : y \in \{x\}^c\}$ which is $(1, 2)^*$ pre-open and so $\{x\}$ is $(1, 2)^*$ pre-closed in X .

(ii) \Rightarrow (iii).

Let $A \subset X$ and $y \notin A$. Then $A \subset \{y\}^c$ and $\{y\}^c$ is $(1, 2)^*$ pre-open in X and $A = \bigcap \{\{y\}^c : y \in A^c\}$ which is the intersection of all $(1, 2)^*$ pre-open sets containing A .

(iii) \Rightarrow (iv).

Obvious.

(iv) \Rightarrow (i).

Let $x, y \in X$, $x \neq y$. By our assumption, there exist atleast an $(1, 2)^*$ pre-open set containing x but not y and also an $(1, 2)^*$ pre-open set containing y but not x . Therefore, X is $(1, 2)^*$ pre- T_1 . \square

Definition 3.12. A space X is called $(1, 2)^*$ pre- T_2 iff for $x, y \in X$, $x \neq y$, there exist disjoint $(1, 2)^*$ pre-open sets U, V in X such that $x \in U$ and $y \in V$.

Remark 3.13. $(1, 2)^*$ pre- T_2 ness implies $(1, 2)^*$ pre- T_1 ness but the converse is not true in general. In Example 3.7, X is $(1, 2)^*$ pre- T_1 but not $(1, 2)^*$ pre- T_2 .

Definition 3.14. A subset O of X is said to be $(1, 2)^*$ pre-neighbourhood of a point $x \in X$ iff there exists an $(1, 2)^*$ pre-open set U such that $x \in U \subset O$.

Theorem 3.15. For a space X the following statements are equivalent.

- (i). X is $(1, 2)^*$ pre- T_2
- (ii). If $x \in X$, then for each $y \neq x$, there is an $(1, 2)^*$ pre-neighbourhood $N(x)$ of x such that $y \notin (1, 2)^*$ pcl($N(x)$).
- (iii). For each $x \in \{(1, 2)^*$ pcl(N): N is an $(1, 2)^*$ pre-neighbourhood of $x\} = \{x\}$.

Proof. (i) \Rightarrow (ii).

Let $x \in X$. If $y \in X$ is such that $y \neq x$, there exist disjoint $(1, 2)^*$ pre-open sets U, V such that $x \in U$ and $y \in V$. Then $x \in U \subset X \setminus V$ which implies that $X \setminus V$ is an $(1, 2)^*$ pre-neighbourhood of x . Also $X \setminus V$ is $(1, 2)^*$ pre-closed and $y \notin X \setminus V$. Let $N(x) = X \setminus V$. Then $y \notin (1, 2)^*$ pcl($N(x)$).

(ii) \Rightarrow (iii).

Obvious.

(iii) \Rightarrow (i).

Let $x, y \in X, x \neq y$. By hypothesis, there is atleast an $(1, 2)^*$ pre-neighbourhood N of x such that $y \notin (1, 2)^*$ pcl(N). We have $x \notin X \setminus (1, 2)^*$ pcl(N) is $(1, 2)^*$ pre-open. Since N is an $(1, 2)^*$ pre-neighbourhood of x , there exists $U \in (1, 2)^*$ PO(X) such that $x \in U \subset N$ and $U \cap (X \setminus (1, 2)^*$ pcl(N)) = \emptyset . Hence X is $(1, 2)^*$ pre- T_2 . □

Definition 3.16. A space X is said to be $(1, 2)^*$ pre-regular if for each $(1, 2)^*$ pre-closed set F and each point $x \notin F$ there exist disjoint $(1, 2)^*$ pre-open sets U and V such that $x \in U$ and $F \subset V$.

Theorem 3.17. An $(1, 2)^*$ pre- T_0 space is $(1, 2)^*$ pre- T_2 if it is $(1, 2)^*$ pre-regular.

Proof. Let X be $(1, 2)^*$ pre- T_0 and $(1, 2)^*$ pre-regular. If $x, y \in X, x \neq y$, there exists $U \in (1, 2)^*$ PO(X) such that U contains one of x and y , say x but not y . Then $X \setminus U$ is $(1, 2)^*$ pre-closed and $x \notin X \setminus U$. Since X is $(1, 2)^*$ pre-regular, there exist disjoint $(1, 2)^*$ pre-open sets V_1 and V_2 such that $x \in V_1$ and $X \setminus U \subset V_2$. Thus $x \in V_1$ and $y \in V_2, V_1 \cap V_2 = \emptyset$. Hence X is $(1, 2)^*$ pre- T_2 . □

Theorem 3.18. If $f : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is an injective, $(1, 2)^*$ pre-irresolute map and Y is $(1, 2)^*$ pre- T_2 then X is $(1, 2)^*$ pre- T_2 .

Proof. Let $x, y \in X, x \neq y$. Since f is injective, $f(x) \neq f(y)$ in Y and there exist disjoint $(1, 2)^*$ pre-open sets U, V such that $f(x) \in U$ and $f(y) \in V$. Let $G = f^{-1}(U)$ and $H = f^{-1}(V)$. Then $x \in G, y \in H$ and $G, H \in (1, 2)^*$ PO(X). Also $G \cap H = f^{-1}(U) \cap f^{-1}(V) = f^{-1}(U \cap V) = \emptyset$. Thus X is $(1, 2)^*$ pre- T_2 . □

4. PRE-DIFFERENCE AXIOMS

Definition 4.1. A subset A of X is called $(1, 2)^*$ pre-difference set (briefly $(1, 2)^*$ pD-set) if there are two $(1, 2)^*$ pre-open sets P_1 and P_2 in $X, P_1 \neq X$ such that $A = P_1 \setminus P_2$.

Remark 4.2. It is evident that each $(1, 2)^*$ pre-open set is an $(1, 2)^*$ pD-set.

Now we define another set of separation axioms called $(1, 2)^*$ pre- $D_i, i = 0, 1, 2$ by using the $(1, 2)^*$ pD-sets.

Definition 4.3. A space X is said to be

- (i). $(1, 2)^*$ pre- D_0 if for $x, y \in X, x \neq y$, there exists an $(1, 2)^*$ pD-set containing one of x and y but not the other.
- (ii). $(1, 2)^*$ pre- D_1 if for $x, y \in X, x \neq y$, there exist $(1, 2)^*$ pD-sets U, V in X such that $x \in U, y \notin U$ and $y \in V, x \notin V$.

- (iii). $(1, 2)^*$ pre- D_2 if for $x, y \in X$, $x \neq y$, there exist disjoint $(1, 2)^*$ pD -sets U, V in X such that $x \in U$ and $y \in V$.

Remark 4.4. (i). Every $(1, 2)^*$ pre- T_i space is $(1, 2)^*$ pre- D_i , $i = 0, 1, 2$ respectively.
(ii). If X is $(1, 2)^*$ pre- D_i then it is $(1, 2)^*$ pre- D_{i-1} , $i = 1, 2$.

Remark 4.5. $(1, 2)^*$ pre- D_i ness does not imply $(1, 2)^*$ pre- T_i ness for $i = 1, 2$ respectively. In example 3.10, X is $(1, 2)^*$ pre- D_1 but not $(1, 2)^*$ pre- T_1 and in Example 3.7, X is not $(1, 2)^*$ pre- T_2 but $(1, 2)^*$ pre- D_2 .

The next example shows that $(1, 2)^*$ pre- D_0 does not imply $(1, 2)^*$ pre- D_1 .

Example 4.6. Let $X = \{a, b\}$, $\tau_1 = \{\emptyset, X\}$, $\tau_2 = \{\emptyset, \{a\}, X\}$. Then X is $(1, 2)^*$ pre- D_0 but not $(1, 2)^*$ pre- D_1 .

Theorem 4.7. A space X is $(1, 2)^*$ pre- D_0 if and only if it is $(1, 2)^*$ pre- T_0 .

Proof. Suppose that X is $(1, 2)^*$ pre- D_0 . Let $x, y \in X$, $x \neq y$. Then there exists an $(1, 2)^*$ pD -set U such that U contains x but not y , say. As U is an $(1, 2)^*$ pD -set, it is possible to write $U = P_1 \setminus P_2$ where $P_1 \neq X$ and $P_1, P_2 \in (1, 2)PO(X)$. Now there arises two cases. (i) $y \notin P_1$ (ii) $y \in P_1$ and $y \in P_2$.

Case (i). $y \notin P_1$ and $x \in P_1 \setminus P_2$ implies that $x \in P_1$ and $y \notin P_1$.

Case (ii). $y \in P_1$ and $y \in P_2$. $x \in P_1 \setminus P_2$ implies that $x \notin P_2$. Thus $y \in P_2$ and $x \notin P_2$.

Thus in both the cases, we obtain that X is $(1, 2)^*$ pre- T_0 . Conversely, if X is $(1, 2)^*$ pre- T_0 , by Remark 4.4, X is $(1, 2)^*$ pre- D_0 . \square

It has been showed in section 3, that an $(1, 2)^*$ pre- T_2 -space is $(1, 2)^*$ pre- T_1 but not the converse. But in the case of pre-Difference axioms, we prove that an $(1, 2)^*$ pre- D_1 -space is $(1, 2)^*$ pre- D_2 and so the $(1, 2)^*$ pre- D_1 -space coincides with $(1, 2)^*$ pre- D_2 -space.

Theorem 4.8. A space X is $(1, 2)^*$ pre- D_1 if and only if X is $(1, 2)^*$ pre- D_2 .

Proof. Necessity. Let $x, y \in X$, $x \neq y$. Then there exist $(1, 2)^*$ pD -sets U, V in X such that $x \in U$, $y \notin U$ and $y \in V$, $x \notin V$. Let $U = P_1 \setminus P_2$ and $V = P_3 \setminus P_4$ where $P_i \in (1, 2)^*PO(X)$, $i = 1, 2, 3, 4$ and $P_1 \neq X$, $P_3 \neq X$. It is evident that $x \notin V$ implies the two possibilities, (i) $x \in P_3 \cap P_4$ (ii) $x \notin P_3$.

Case (i). $x \in P_3 \cap P_4$. We have $x \in P_4$ and $y \in P_3 \setminus P_4$ and $P_4 \cap (P_3 \setminus P_4) = \emptyset$ are disjoint.

Case (ii). $x \notin P_3$. $y \notin U$ implies that either $y \in P_1$ and $y \in P_2$ or $y \notin P_1$.

Sub Case (a). $y \in P_1$ and $y \in P_2$ and $x \in P_1 \setminus P_2$. We get $P_1 \setminus P_2$ and P_2 are disjoint $(1, 2)^*$ pD -sets containing x and y respectively.

Sub Case (b). $y \notin P_1$ and $x \in P_1 \setminus P_2$ and $x \notin P_3$ implies that $x \in P_1 \setminus (P_2 \cup P_3)$ and $y \in P_3 \setminus P_4$ and $y \notin P_1$ implies that $y \in P_3 \setminus (P_1 \cup P_4)$ and $P_1 \setminus (P_2 \cup P_3)$ and $P_3 \setminus (P_1 \cup P_4)$ are disjoint. Therefore, X is $(1, 2)^*$ pre- D_2 .

Sufficiency. Follows from Remark 4.4. \square


Theorem 4.9. If X is $(1, 2)^*$ pre- D_1 then it is $(1, 2)^*$ pre- T_0 .

Proof. Follows from (ii) Remark 4.4 and theorem 4.8. \square

Remark 4.10. An $(1, 2)^*$ pre- T_0 -space is not $(1, 2)^*$ pre- D_1 , in general. In Example 3.10, X is $(1, 2)^*$ pre- T_0 but not $(1, 2)^*$ pre- D_1 .

Remark 4.11. From the discussions in Sections three and four, the following implication diagram is drawn. In the diagram,

1. $(1, 2)^*$ pre- T_2
2. $(1, 2)^*$ pre- T_1
3. $(1, 2)^*$ pre- T_0
4. $(1, 2)^*$ pre- D_1



$A \rightarrow B$ (resp. $A \nrightarrow B$) represents that A implies B (resp. A does not imply B).

REFERENCES

1. Abhijit Chattopadhyay, *Pre- T_0 and Pre- T_1 topological spaces*, J. Indian Acad. Math. **17** (1995), no. 2, 156–159.
2. Ashish Kar and Paritosh Bhattacharyya, *Some weak separation axioms*, Bull. Cal. Math. Soc. **82** (1990), 415–422.
3. A. S. Davis, *Indexed system of neighbourhoods for general topological spaces*, Amer. Math. Monthly **68** (1961), 886–893.
4. S. N. El-Deeb, I. A. Hasanein, A. S. Mashhour, and T. Noiri, *On p -regular spaces*, Bull. Math. Soc. Sci. Math. R. S. Roumanie (N. S.) **27(75)** (1983), 311–315.
5. N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly **70** (1963), 36–41.
6. M. Lellis Thivagar and O. Ravi, *A bitopological $(1,2)^*$ semi generalized continuous mappings*, Bull. Malaysian Math. Soc. **29** (2005), no. 1, 1–9.
7. M. Lellis Thivagar, *Generalization of Pairwise α -continuous functions*, Pure and Applied Mathematics Sciences **33** (1991), 55–63.
8. S. N. Maheshwari and R. Prasad, *Some new separation axioms*, Ann. Soc. Sci. Bruxelles **89** (1975), 395–402.
9. A. S. Mashhour, M. E. Abd El-Monsef, and S.N. El-Deeb, *On pre-continuous and weak pre-continuous mappings*, Proc. Math. and Phys. Soc. Egypt (1982), 47–53.
10. Miguel Caldas, *A separation axiom between semi- T_0 and semi- T_1* , Mem. Fac. Sci. Kochi. Univ. (Math.) **18** (1997), 37–42.
11. G. B. Navalagi, *P -normal, almost P -normal and mildly p -normal spaces*, Topology Atlas Preprint 427.
12. G. B. Navalagi, *Pre-neighbourhoods*, Topology Atlas Preprint 422.

DEPARTMENT OF MATHEMATICS, JAYARAJ ANNAPACKIAM COLLEGE FOR WOMEN, PERIYAKULAM, THENI (DT.)-625601, TAMILNADU, INDIA

E-mail address: athisayaponmani@yahoo.co.in

DEPARTMENT OF MATHEMATICS, ARUL ANANDAR COLLEGE, KARUMATHUR, MADURAI (DT.)-625514, TAMILNADU, INDIA

E-mail address: mlthivagar@yahoo.co.in

Received 01/02/2006