

CONSERVATIVE DISCRETE TIME-INVARIANT SYSTEMS AND BLOCK OPERATOR CMV MATRICES

YURY ARLINSKIĬ

Dedicated to the memory of A. Ya. Povzner.

ABSTRACT. It is well known that an operator-valued function Θ from the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$, where \mathfrak{M} and \mathfrak{N} are separable Hilbert spaces, can be realized as a transfer function of a simple conservative discrete time-invariant linear system. The known realizations involve the function Θ itself, the Hardy spaces or the reproducing kernel Hilbert spaces. On the other hand, as in the classical scalar case, the Schur class operator-valued function is uniquely determined by its so-called "Schur parameters". In this paper we construct simple conservative realizations using the Schur parameters only. It turns out that the unitary operators corresponding to the systems take the form of five diagonal block operator matrices, which are analogs of Cantero–Moral–Velázquez (CMV) matrices appeared recently in the theory of scalar orthogonal polynomials on the unit circle. We obtain new models given by truncated block operator CMV matrices for an arbitrary completely non-unitary contraction. It is shown that the minimal unitary dilations of a contraction in a Hilbert space and the minimal Naimark dilations of a semi-spectral operator measure on the unit circle can also be expressed by means of block operator CMV matrices.

1. INTRODUCTION

In what follows the class of all continuous linear operators defined on a complex Hilbert space \mathfrak{H}_1 and taking values in a complex Hilbert space \mathfrak{H}_2 is denoted by $\mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$ and $\mathbf{L}(\mathfrak{H}) := \mathbf{L}(\mathfrak{H}, \mathfrak{H})$. We denote by $I_{\mathcal{H}}$ the identity operator in a Hilbert space \mathcal{H} and by $P_{\mathcal{L}}$ the orthogonal projection onto the subspace (the closed linear manifold) \mathcal{L} . The notation $T|_{\mathcal{L}}$ means the restriction of a linear operator T on the set \mathcal{L} . The range and the null-space of a linear operator T are denoted by $\text{ran } T$ and $\ker T$, respectively. We use the usual symbols $\mathbb{C}, \mathbb{Z}, \mathbb{N}$, and \mathbb{N}_0 for the sets of complex numbers, integers, positive integers, and nonnegative integers, respectively.

Recall that an operator $T \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$ is said to be

- *contractive* if $\|T\| \leq 1$;
- *isometric* if $\|Tf\| = \|f\|$ for all $f \in \mathfrak{H}_1 \iff T^*T = I_{\mathfrak{H}_1}$;
- *co-isometric* if T^* is isometric $\iff TT^* = I_{\mathfrak{H}_2}$;
- *unitary* if it is both isometric and co-isometric.

Given a contraction $T \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$, the operators

$$D_T := (I - T^*T)^{1/2}, \quad D_{T^*} := (I - TT^*)^{1/2}$$

are called the *defect operators* of T , and the subspaces

$$\mathfrak{D}_T = \overline{\text{ran } D_T}, \quad \mathfrak{D}_{T^*} = \overline{\text{ran } D_{T^*}}$$

2000 *Mathematics Subject Classification.* 47A48, 47A56, 47B36, 93B28.

Key words and phrases. Contraction, conservative system, transfer function, realization, Schur class function, Schur parameters, block operator CMV matrices, unitary dilation.

the *defect subspaces* of T . The dimensions $\dim \mathfrak{D}_T$, $\dim \mathfrak{D}_{T^*}$ are known as the *defect numbers* of T . The defect operators satisfy the following intertwining relations

$$(1.1) \quad TD_T = D_{T^*}T, \quad T^*D_{T^*} = D_T T^*.$$

It follows from (1.1) that $T\mathfrak{D}_T \subset \mathfrak{D}_{T^*}$, $T^*\mathfrak{D}_{T^*} \subset \mathfrak{D}_T$, and $T(\ker D_T) = \ker D_{T^*}$, $T^*(\ker D_{T^*}) = \ker D_T$. Moreover, the operators $T \upharpoonright \ker D_T$ and $T^* \upharpoonright \ker D_{T^*}$ are isometries and $T \upharpoonright \mathfrak{D}_T$ and $T^* \upharpoonright \mathfrak{D}_{T^*}$ are *pure* contractions, i.e., $\|Tf\| < \|f\|$ for $f \in \mathfrak{H} \setminus \{0\}$.

The *Schur class* $\mathbf{S}(\mathfrak{H}_1, \mathfrak{H}_2)$ is the set of all holomorphic and contractive $\mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$ -valued functions on the unit disk $\mathbb{D} = \{\lambda \in \mathbb{C} : |\lambda| < 1\}$. This class is a natural generalization of the Schur class \mathbf{S} of scalar analytic functions mapping the unit disk \mathbb{D} into the closed unit disk $\overline{\mathbb{D}}$ [62] and is intimately connected with spectral theory and models for Hilbert space contraction operators [71], [24], [25], [26], [27], [28], the Lax-Phillips scattering theory [55], [1], [21], the theory of scalar and matrix orthogonal polynomials on the unit circle $\mathbb{T} = \{\xi \in \mathbb{C} : |\xi| = 1\}$ [37], [66], [39], [40], the theory of passive (contractive) discrete time-invariant linear systems [51], [52], [12], [13], [14], [20]. One of the characterization of the operator-valued Schur class is that any $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ can be realized as a transfer (characteristic) function of the form

$$\Theta(\lambda) = D + \lambda C(I_{\mathfrak{H}} - \lambda A)^{-1}B, \quad \lambda \in \mathbb{D},$$

of a discrete time-invariant system (colligation)

$$\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}$$

with the input space \mathfrak{M} , the output space \mathfrak{N} , and some state space \mathfrak{H} . Moreover, if the operator U_τ is given by the block operator matrix

$$U_\tau = \begin{bmatrix} D & C \\ B & A \end{bmatrix} : \begin{matrix} \mathfrak{M} & \mathfrak{N} \\ \oplus & \rightarrow \oplus \\ \mathfrak{H} & \mathfrak{H} \end{matrix},$$

then the system τ can be chosen **(a)** passive (U_τ is contractive) and minimal, **(b)** co-isometric (U_τ is co-isometry) and observable, **(c)** isometric (U_τ is isometry) and controllable, **(d)** conservative (U_τ is unitary) and simple (see Section 3). The corresponding models of the systems τ and the state space operators A are well-known. We mention the de Branges–Rovnyak functional model of a co-isometric system [25], [6], [56], the Sz.-Nagy–Foias [71], the Pavlov [58], [59], [60], and the Nikol’skiĭ–Vasyunin [57] functional models of completely non-unitary contractions, the Brodskii [28] functional model of a simple unitary colligation, the Arov–Kaashoek–Pik [14] functional model of a passive minimal and optimal system. All these models involve the Schur class function and/or the Hardy spaces, the de Branges–Rovnyak reproducing kernel Hilbert space.

The main goal of the present paper is constructions of models for simple conservative systems and completely non-unitary contractions by means of the operator analogs of the scalar CMV matrices, which recently appeared in the theory of orthogonal polynomials on the unit circle [30], [66], [68], [37].

In the paper of M. J. Cantero, L. Moral, and L. Velázquez [30] it is established that the semi-infinite matrices of the form

$$(1.2) \quad C = C(\{\alpha_n\}) = \begin{pmatrix} \bar{\alpha}_0 & \bar{\alpha}_1 \rho_0 & \rho_1 \rho_0 & 0 & 0 & \dots \\ \rho_0 & -\bar{\alpha}_1 \alpha_0 & -\rho_1 \alpha_0 & 0 & 0 & \dots \\ 0 & \bar{\alpha}_2 \rho_1 & -\bar{\alpha}_2 \alpha_1 & \bar{\alpha}_3 \rho_2 & \rho_3 \rho_2 & \dots \\ 0 & \rho_2 \rho_1 & -\rho_2 \alpha_1 & -\bar{\alpha}_3 \alpha_2 & -\rho_3 \alpha_2 & \dots \\ 0 & 0 & 0 & \bar{\alpha}_4 \rho_3 & -\bar{\alpha}_4 \alpha_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

and

$$(1.3) \quad \tilde{\mathcal{C}} = \tilde{\mathcal{C}}(\{\alpha_n\}) = \begin{pmatrix} \bar{\alpha}_0 & \rho_0 & 0 & 0 & 0 & \dots \\ \bar{\alpha}_1 \rho_0 & -\bar{\alpha}_1 \alpha_0 & \bar{\alpha}_2 \rho_1 & \rho_2 \rho_1 & 0 & \dots \\ \rho_1 \rho_0 & -\rho_1 \alpha_0 & -\bar{\alpha}_2 \alpha_1 & -\rho_2 \alpha_1 & 0 & \dots \\ 0 & 0 & \bar{\alpha}_3 \rho_2 & -\bar{\alpha}_3 \alpha_2 & \bar{\alpha}_4 \rho_3 & \dots \\ 0 & 0 & \rho_3 \rho_2 & -\rho_3 \alpha_2 & -\bar{\alpha}_4 \alpha_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

give representations of the unitary operator $(Uf)(\zeta) = \zeta f(\zeta)$ in $L_2(\mathbb{T}, d\mu)$, where the $d\mu$ is a nontrivial probability measure on the unite circle, with respect to the orthonormal systems obtained by orthonormalization of the sequences $\{1, \zeta, \zeta^{-1}, \zeta^2, \zeta^{-2}, \dots\}$ and $\{1, \zeta^{-1}, \zeta, \zeta^{-2}, \zeta^2, \dots\}$, respectively. The Verblunsky coefficients $\{\alpha_n\}$, $|\alpha_n| < 1$, arise in the Szegő recurrence formula

$$\zeta \Phi_n(\zeta) = \Phi_{n+1}(\zeta) + \bar{\alpha}_n \zeta^n \overline{\Phi_n(1/\bar{\zeta})}, \quad n = 0, 1, \dots$$

for monic orthogonal with respect to $d\mu$ polynomials $\{\Phi_n\}$, and $\rho_n := \sqrt{1 - |\alpha_n|^2}$. The matrices $\mathcal{C}(\{\alpha_n\})$ and $\tilde{\mathcal{C}}(\{\alpha_n\})$ are called the *CMV matrices*. The matrix $\tilde{\mathcal{C}}$ is transpose to \mathcal{C} . Notice that it has been shown by Berezansky and Dudkin in [22] and [43] that the operator $(Uf)(\zeta) = \zeta f(\zeta)$ admits a three-diagonal block matrix representation.

Given a probability measure μ on \mathbb{T} , define the *Carathéodory function* by

$$F(\lambda) = F(\lambda, \mu) := \int_{\mathbb{T}} \frac{\zeta + \lambda}{\zeta - \lambda} d\mu(\zeta) = 1 + 2 \sum_{n=1}^{\infty} \beta_n \lambda^n, \quad \beta_n = \int_{\mathbb{T}} \zeta^{-n} d\mu$$

the moments of μ . F is an analytic function in \mathbb{D} which obeys $\text{Re } F > 0$, $F(0) = 1$. The Schur class function $f(\lambda)$ is then defined by

$$f(\lambda) = f(\lambda, \mu) := \frac{1}{\lambda} \frac{F(\lambda) - 1}{F(\lambda) + 1}.$$

Given a Schur function $f(\lambda)$, which is not a finite Blaschke product, define inductively

$$f_0(\lambda) = f(\lambda), \quad f_{n+1}(\lambda) = \frac{f_n(\lambda) - f_n(0)}{\lambda(1 - \bar{f}_n(0)f_n(\lambda))}, \quad n \in \mathbb{N}_0.$$

It is clear that $\{f_n\}$ is an *infinite* sequence of Schur functions called the *n*-th *Schur iterates* and neither of its terms is a finite Blaschke product. The numbers $\gamma_n := f_n(0)$ are called the *Schur parameters*

$$\mathcal{S}f = \{\gamma_0, \gamma_1, \dots\}.$$

Note that

$$f_n(\lambda) = \frac{\gamma_n + \lambda f_{n+1}(\lambda)}{1 + \bar{\gamma}_n \lambda f_{n+1}(\lambda)} = \gamma_n + (1 - |\gamma_n|^2) \frac{\lambda f_{n+1}(\lambda)}{1 + \bar{\gamma}_n \lambda f_{n+1}(\lambda)}, \quad n \in \mathbb{N}_0.$$

The method of labeling $f \in \mathbf{S}$ by its Schur parameters is known as the *Schur algorithm* and is due to I. Schur [62]. In the case when

$$f(\lambda) = e^{i\varphi} \prod_{k=1}^N \frac{\lambda - \lambda_k}{1 - \bar{\lambda}_k \lambda}$$

is a finite Blaschke product of order N , the Schur algorithm terminates at the N -th step. The sequence of Schur parameters $\{\gamma_n\}_{n=0}^N$ is finite, $|\gamma_n| < 1$ for $n = 0, 1, \dots, N - 1$, and $|\gamma_N| = 1$.

Due to Geronimus theorem for the function $f(\lambda, \mu)$ the relations $\gamma_n = \alpha_n$ hold true for all $n = 0, 1, \dots$

There is a nice multiplicative structure of the CMV matrices. In the semi-infinite case \mathcal{C} and $\tilde{\mathcal{C}}$ are the products of two matrices: $\mathcal{C} = \mathcal{L}\mathcal{M}$, $\tilde{\mathcal{C}} = \mathcal{M}\mathcal{L}$, where

$$\begin{aligned}\mathcal{L} &= \Psi(\alpha_0) \oplus \Psi(\alpha_2) \oplus \dots \oplus \Psi(\alpha_{2m}) \oplus \dots, \\ \mathcal{M} &= \mathbf{1}_{1 \times 1} \oplus \Psi(\alpha_1) \oplus \Psi(\alpha_3) \oplus \dots \oplus \Psi(\alpha_{2m+1}) \oplus \dots,\end{aligned}$$

and $\Psi(\alpha) = \begin{pmatrix} \bar{\alpha} & \rho \\ \rho & -\alpha \end{pmatrix}$. The finite $(N+1) \times (N+1)$ CMV matrices \mathcal{C} and $\tilde{\mathcal{C}}$ obey $\alpha_0, \alpha_1, \dots, \alpha_{N-1} \in \mathbb{D}$ and $|\alpha_N| = 1$, and also $\mathcal{C} = \mathcal{L}\mathcal{M}$, $\tilde{\mathcal{C}} = \mathcal{M}\mathcal{L}$, where in this case $\Psi(\alpha_N) = (\bar{\alpha}_N)$.

In the paper [11] it is established that the *truncated* CMV matrices

$$\begin{aligned}\mathcal{T}_0 &= \begin{pmatrix} -\bar{\alpha}_1\alpha_0 & -\rho_1\alpha_0 & 0 & 0 & \dots \\ \bar{\alpha}_2\rho_1 & -\bar{\alpha}_2\alpha_1 & \bar{\alpha}_3\rho_2 & \rho_3\rho_2 & \dots \\ \rho_2\rho_1 & -\rho_2\alpha_1 & -\bar{\alpha}_3\alpha_2 & -\rho_3\alpha_2 & \dots \\ 0 & 0 & \bar{\alpha}_4\rho_3 & -\bar{\alpha}_4\alpha_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \\ \tilde{\mathcal{T}}_0 &= \begin{pmatrix} -\bar{\alpha}_1\alpha_0 & \bar{\alpha}_2\rho_1 & \rho_2\rho_1 & 0 & \dots \\ -\rho_1\alpha_0 & -\bar{\alpha}_2\alpha_1 & -\rho_2\alpha_1 & 0 & \dots \\ 0 & \bar{\alpha}_3\rho_2 & -\bar{\alpha}_3\alpha_2 & \bar{\alpha}_4\rho_3 & \dots \\ 0 & \rho_3\rho_2 & -\rho_3\alpha_2 & -\bar{\alpha}_4\alpha_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}\end{aligned}$$

obtained from the “full” CMV matrices

$$\mathcal{C} = \mathcal{C}(\{\alpha_n\}) \quad \text{and} \quad \tilde{\mathcal{C}} = \tilde{\mathcal{C}}(\{\alpha_n\})$$

by deleting the first row and the first column, provide models of completely non-unitary contractions with rank one defect operators.

As pointed out by Simon in [68], the history of CMV matrices is started with the papers of Bunse-Gerstner and Elsner [29] (1991) and Watkins [73] (1993), where unitary semi-infinite five-diagonal matrices were introduced and studied. In [30] Cantero, Moral, and Velazquez (CMV) re-discovered them. In a context different from orthogonal polynomials on the unit circle, Bourget, Howland, and Joye [23] introduced a set of doubly infinite family of matrices with three sets of parameters which for special choices of the parameters reduces to two-sided CMV matrices on $\ell^2(\mathbb{Z})$.

The Schur algorithm for matrix valued Schur class functions and its connection with the matrix orthogonal polynomials on the unit circle have been considered in the paper of Delsarte, Genin, and Kamp [40] and in the book of Dubovoj, Fritzsche, and Kirstein [42]. The CMV matrices, connected with matrix orthogonal polynomials on the unit circle with respect to nontrivial matrix-valued measures are considered in [68], [37]. If the $k \times k$ matrix-valued non-trivial measure μ on \mathbb{T} , $\mu(\mathbb{T}) = I_{k \times k}$ is given, then there are the left and the right orthonormal matrix polynomials. The Szegő recursions take slightly different form than in the scalar case and the Verblunsky $k \times k$ matrix coefficients (the Schur parameters of the corresponding matrix-valued Schur function) $\{\alpha_n\}$ satisfy the inequality $\|\alpha_n\| < 1$ for all n . The latter condition is in fact equivalent to the non-triviality of the measure. The entries of the corresponding CMV matrix have the size $k \times k$ and the numbers ρ_n are replaced by the $k \times k$ defect matrices

$$\rho_n^L = D_{\alpha_n} = (I - \alpha_n^* \alpha_n)^{1/2} \quad \text{and} \quad \rho_n^R = D_{\alpha_n^*} = (I - \alpha_n \alpha_n^*)^{1/2},$$

where α^* is the adjoint matrix. In these notations the CMV matrix is of the form [37]

$$(1.4) \quad \mathcal{C} = \mathcal{C}(\{\alpha_n\}) = \begin{pmatrix} \alpha_0^* & \rho_0^L \alpha_1^* & \rho_0^L \rho_1^L & 0 & 0 & \dots \\ \rho_0^R & -\alpha_0 \alpha_1^* & -\alpha_0 \rho_1^L & 0 & 0 & \dots \\ 0 & \alpha_2^* \rho_1^R & -\alpha_2^* \alpha_1 & \rho_2^L \alpha_3^* & \rho_2^L \rho_3^L & \dots \\ 0 & \rho_2^R \rho_1^R & -\rho_2^R \alpha_1 & -\alpha_2 \alpha_3^* & -\alpha_2 \rho_3^L & \dots \\ 0 & 0 & 0 & \alpha_4^* \rho_3^R & -\alpha_4^* \alpha_3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}.$$

Notice that the spectral problems for CMV matrices with scalar and matrix elements and truncated CMV matrices with scalar elements were considered in [66], [67], [37], [22], [32], [47], [72], [11], [48], [49], [50], [54].

The operator extension of the Schur algorithm was developed by T. Constantinescu in [34] and with numerous applications is presented in the monographs [19], [36], [42], [44]. The next theorem goes back to Shmul'yan [63], [64] and T. Constantinescu [34] (see also [19], [7], [8]) and plays a key role in the operator Schur algorithm.

Theorem 1.1. *Let \mathfrak{M} and \mathfrak{N} be separable Hilbert spaces and let the function Θ be from the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$. Then there exists a function Z from the Schur class $\mathbf{S}(\mathfrak{D}_{\Theta(0)}, \mathfrak{D}_{\Theta^*(0)})$ such that*

$$(1.5) \quad \Theta(\lambda) = \Theta(0) + D_{\Theta^*(0)} Z(\lambda) (I + \Theta^*(0) Z(\lambda))^{-1} D_{\Theta(0)}, \quad \lambda \in \mathbb{D}.$$

The representation (1.5) of a function Θ from the Schur class is called the Möbius representation of Θ and the function Z is called the Möbius parameter of Θ (see [7], [8]). Clearly, $Z(0) = 0$ and by Schwartz's lemma we obtain that

$$\|Z(\lambda)\| \leq |\lambda|, \quad \lambda \in \mathbb{D}.$$

The operator Schur's algorithm [19]. Fix $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$, put $\Theta_0(\lambda) = \Theta(\lambda)$ and let Z_0 be the Möbius parameter of Θ . Define

$$\Gamma_0 = \Theta(0), \quad \Theta_1(\lambda) = \frac{Z_0(\lambda)}{\lambda} \in \mathbf{S}(\mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}), \quad \Gamma_1 = \Theta_1(0) = Z_0'(0).$$

If $\Theta_0, \dots, \Theta_n$ and $\Gamma_0, \dots, \Gamma_n$ have been chosen, then let $Z_{n+1} \in \mathbf{S}(\mathfrak{D}_{\Gamma_n}, \mathfrak{D}_{\Gamma_n^*})$ be the Möbius parameter of Θ_n . Put

$$\Theta_{n+1}(\lambda) = \frac{Z_{n+1}(\lambda)}{\lambda}, \quad \Gamma_{n+1} = \Theta_{n+1}(0).$$

The contractions $\Gamma_0 \in \mathbf{L}(\mathfrak{M}, \mathfrak{N})$, $\Gamma_n \in \mathbf{L}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*})$, $n \in \mathbb{N}$, are called the *Schur parameters* of Θ and the function $\Theta_n \in \mathbf{S}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*})$ we will call the *n-th Schur iterate* of $\Theta(\lambda)$.

Formally we have

$$\Theta_{n+1}(\lambda) \upharpoonright \text{ran } D_{\Gamma_n} = \frac{1}{\lambda} D_{\Gamma_n^*} (I_{\mathfrak{D}_{\Gamma_n^*}} - \Theta_n(\lambda) \Gamma_n^*)^{-1} (\Theta_n(\lambda) - \Gamma_n) D_{\Gamma_n}^{-1} \upharpoonright \text{ran } D_{\Gamma_n}.$$

Clearly, the sequence of Schur parameters $\{\Gamma_n\}$ is infinite if and only if all operators Γ_n are non-unitary. The sequence of Schur parameters consists of a finite number of operators $\Gamma_0, \Gamma_1, \dots, \Gamma_N$ if and only if $\Gamma_N \in \mathbf{L}(\mathfrak{D}_{\Gamma_{N-1}}, \mathfrak{D}_{\Gamma_{N-1}^*})$ is unitary. If Γ_N is isometric (co-isometric) then $\Gamma_n = 0$ for all $n > N$. The following generalization of the classical Schur result is proved in [34] (see also [19]).

Theorem 1.2. *There is a one-to-one correspondence between the Schur class functions $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ and the set of all sequences of contractions $\{\Gamma_n\}_{n \geq 0}$ such that*

$$(1.6) \quad \Gamma_0 \in \mathbf{L}(\mathfrak{M}, \mathfrak{N}), \quad \Gamma_n \in \mathbf{L}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*}), \quad n \in \mathbb{N}.$$

A sequence of contractions of the form (1.6) is called the *choice sequence* [31]. Such objects are used for the indexing of contractive intertwining dilations, of positive Toeplitz

and

$$\tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & D_{\Gamma_1^*} D_{\Gamma_2^*} & 0 & 0 & 0 & \dots \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & -\Gamma_1^* D_{\Gamma_2^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & \Gamma_3 D_{\Gamma_2} & -\Gamma_3 \Gamma_2^* & D_{\Gamma_3^*} \Gamma_4 & D_{\Gamma_3^*} D_{\Gamma_4^*} & 0 & \dots \\ 0 & 0 & D_{\Gamma_3} D_{\Gamma_2} & -D_{\Gamma_3} \Gamma_2^* & -\Gamma_3^* \Gamma_4 & -\Gamma_3^* D_{\Gamma_4^*} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Notice that the relation $\tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0}) = (\mathcal{U}_0(\{\Gamma_n^*\}_{n \geq 0}))^*$ holds true. Hence the CMV matrix (1.4) corresponds to the case

$$\begin{aligned} \mathfrak{M} = \mathfrak{N} = \mathfrak{D}_{\Gamma_0} = \mathfrak{D}_{\Gamma_0^*} = \mathfrak{D}_{\Gamma_1} = \mathfrak{D}_{\Gamma_1^*} = \dots = \mathfrak{D}_{\Gamma_n} = \mathfrak{D}_{\Gamma_n^*} = \dots = \mathbb{C}^k, \\ \alpha_n = \Gamma_n^*, \quad n \in \mathbb{N}_0. \end{aligned}$$

The block operator truncated CMV matrices

$$\mathcal{T}_0 = \mathcal{T}_0(\{\Gamma_n\}_{n \geq 0}) := P_{\mathfrak{H}_0} \mathcal{U}_0 \upharpoonright \mathfrak{H}_0 \quad \text{and} \quad \tilde{\mathcal{T}}_0 = \tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 0}) := P_{\tilde{\mathfrak{H}}_0} \tilde{\mathcal{U}}_0 \upharpoonright \tilde{\mathfrak{H}}_0$$

are given by

$$\begin{aligned} \mathcal{T}_0 &= (-\Gamma_0^* \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2n}} \oplus \dots) (\mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2n-1}} \oplus \dots) : \mathfrak{H}_0 \rightarrow \mathfrak{H}_0, \\ \tilde{\mathcal{T}}_0 &= (\mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2n-1}} \oplus \dots) (-\Gamma_0^* \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2n}} \oplus \dots) : \tilde{\mathfrak{H}}_0 \rightarrow \tilde{\mathfrak{H}}_0 \end{aligned}$$

and can be rewritten in the three diagonal block operator matrix form with 2×2 entries

$$\mathcal{T}_0 = \begin{bmatrix} \mathcal{B}_1 & \mathcal{C}_1 & 0 & 0 & 0 & \cdot \\ \mathcal{A}_1 & \mathcal{B}_2 & \mathcal{C}_2 & 0 & 0 & \cdot \\ 0 & \mathcal{A}_2 & \mathcal{B}_3 & \mathcal{C}_3 & 0 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad \tilde{\mathcal{T}}_0 = \begin{bmatrix} \tilde{\mathcal{B}}_1 & \tilde{\mathcal{C}}_1 & 0 & 0 & 0 & \cdot \\ \tilde{\mathcal{A}}_1 & \tilde{\mathcal{B}}_2 & \tilde{\mathcal{C}}_2 & 0 & 0 & \cdot \\ 0 & \tilde{\mathcal{A}}_2 & \tilde{\mathcal{B}}_3 & \tilde{\mathcal{C}}_3 & 0 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

The constructions above and the corresponding results are presented in Section 5. We essentially rely on the constructions of simple conservative realizations of the Schur iterates $\{\Theta_n\}_{n \geq 1}$ by means of a given simple conservative realization of the function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ [8]. A brief survey of the results in [8] is given in Section 4. The cases when the Schur parameter $\Gamma_m \in \mathbf{L}(\mathfrak{D}_{\Gamma_{m-1}}, \mathfrak{D}_{\Gamma_{m-1}^*})$ of the function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ is isometric, co-isometric, unitary are considered in detail in Section 6. Observe that in fact we give another prove of Theorem 1.2 (the uniqueness of the function from $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ with given its Schur parameters is proved in Section 2). In Section 7 we obtain in the block operator CMV matrix form the minimal unitary dilations of a contraction and the minimal Naimark dilations of a semi-spectral measure on the unit circle. Another and more complicated constructions of the minimal Naimark dilation and a simple conservative realization for a function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ by means of its Schur parameters are given in [33] and in [53], respectively (see also [19]). Simple conservative realizations of scalar Schur functions with operators A, B, C , and D expressed via corresponding Schur parameters have been obtained by V. Dubovoj [41].

We also prove in Section 7 that a unitary operator U in a separable Hilbert space \mathfrak{K} having a cyclic subspace \mathfrak{M} ($\overline{\text{span}}\{U^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathfrak{K}$) is unitarily equivalent to the block operator CMV matrices \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ constructed by means of the Schur parameters of the function $\Theta(\lambda) = \frac{1}{\lambda} (F_{\mathfrak{M}}^*(\bar{\lambda}) - I_{\mathfrak{M}})(F_{\mathfrak{M}}^*(\bar{\lambda}) + I_{\mathfrak{M}})^{-1}$, where $F_{\mathfrak{M}}(\lambda) = P_{\mathfrak{M}}(U + \lambda I_{\mathfrak{K}})(U - \lambda I_{\mathfrak{K}})^{-1} \upharpoonright \mathfrak{M}$, $\lambda \in \mathbb{D}$. In the last Section 8 we prove that the Sz.-Nagy-Foias [71] characteristic functions of truncated block operator CMV matrices \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$, constructing by means of the Schur parameters $\{\Gamma_n\}_{n \geq 0}$ of a purely contractive function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$, coincide with Θ in the sense of [71].

2. THE SCHUR CLASS FUNCTIONS AND THEIR ITERATES

In the sequel we need the well known fact [71], [19] that if $T \in \mathbf{L}(\mathfrak{H}_1, \mathfrak{H}_2)$ is a contraction which is neither isometric nor co-isometric, then the operator (*elementary rotation* [19]) \mathbf{J}_T given by the operator matrix

$$\mathbf{J}_T = \begin{bmatrix} T & D_{T^*} \\ D_T & -T^* \end{bmatrix} : \begin{array}{c} \mathfrak{H}_1 \\ \oplus \\ \mathfrak{D}_{T^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{H}_2 \\ \oplus \\ \mathfrak{D}_T \end{array}$$

is unitary. Clearly, $\mathbf{J}_T^{-1} = \mathbf{J}_T^* = \mathbf{J}_{T^*}$. If T is isometric or co-isometric, then the corresponding unitary elementary rotation takes the row or the column form

$$\mathbf{J}_T^{(r)} = [T \quad I_{\mathfrak{D}_{T^*}}] : \begin{array}{c} \mathfrak{H}_1 \\ \oplus \\ \mathfrak{D}_{T^*} \end{array} \rightarrow \mathfrak{H}_2, \quad \mathbf{J}_T^{(c)} = \begin{bmatrix} T \\ D_T \end{bmatrix} : \mathfrak{H}_1 \rightarrow \begin{array}{c} \oplus \\ \mathfrak{D}_T \end{array},$$

and $(\mathbf{J}_T^{(r)})^* = \mathbf{J}_{T^*}^{(c)}$. In Section 5 we will need the following statement.

Proposition 2.1. [10]. *Let T be a contraction. Then $Th = D_{T^*}g$ if and only if there exists a vector $\varphi \in \mathfrak{D}_T$ such that $h = D_T\varphi$ and $g = T\varphi$.*

Recall that if $\Theta \in \mathbf{S}(\mathfrak{H}_1, \mathfrak{H}_2)$ then there is a uniquely determined decomposition [71, Proposition V.2.1]

$$\Theta(\lambda) = \begin{bmatrix} \Theta_p(\lambda) & 0 \\ 0 & \Theta_u \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Theta(0)} \\ \oplus \\ \ker D_{\Theta(0)} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Theta^*(0)} \\ \oplus \\ \ker D_{\Theta^*(0)} \end{array},$$

where $\Theta_p \in \mathbf{S}(\mathfrak{D}_{\Theta(0)}, \mathfrak{D}_{\Theta^*(0)})$, Θ_p is a pure contraction and Θ_u is a unitary constant. The function Θ_p is called the *pure part* of Θ (see [19]). If $\Theta(0)$ is isometric (respect., co-isometric) then the pure part is of the form $\Theta_p(\lambda) = 0 \in \mathbf{S}(\{0\}, \mathfrak{D}_{\Theta^*(0)})$ (respect., $\Theta_p(\lambda) = 0 \in \mathbf{S}(\mathfrak{D}_{\Theta(0)}, \{0\})$) for all $\lambda \in \mathbb{D}$. The function Θ is called purely contractive if $\ker D_{\Theta(0)} = \{0\}$. Two operator-valued functions $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ and $\Omega \in \mathbf{S}(\mathfrak{K}, \mathfrak{L})$ coincide [71] if there are two unitary operators $V : \mathfrak{N} \rightarrow \mathfrak{L}$ and $U : \mathfrak{K} \rightarrow \mathfrak{M}$ such that

$$(2.1) \quad V\Theta(\lambda)U = \Omega(\lambda), \quad \lambda \in \mathbb{D}.$$

For the corresponding Schur parameters and the Schur iterates relation (2.1) yields the equalities

$$(2.2) \quad \begin{aligned} G_n &= V\Gamma_n U, \\ \mathfrak{D}_{G_n} &= U^*\mathfrak{D}_{\Gamma_n}, \quad \mathfrak{D}_{G_n^*} = V\mathfrak{D}_{\Gamma_n^*}, \quad D_{G_n} = U^*D_{\Gamma_n}U, \quad D_{G_n^*} = VD_{\Gamma_n^*}V^*, \\ V\Theta_n(\lambda)U &= \Omega_n(\lambda), \quad \lambda \in \mathbb{D} \end{aligned}$$

for all $n \in \mathbb{N}_0$.

In what follows we give a proof of Theorem 1.6 different from the original one in [34]. First of all we will prove the uniqueness. The existence will be proved in Section 5.

Theorem 2.2. *Any choice sequence*

$$\Gamma_0 \in \mathbf{L}(\mathfrak{M}, \mathfrak{N}), \quad \Gamma_n \in \mathbf{L}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*}), \quad n \geq 1$$

uniquely determines a function from the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$.

Proof. Let $\Gamma_0 \in \mathbf{L}(\mathfrak{M}, \mathfrak{N})$, $\Gamma_n \in \mathbf{L}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*})$, $n \geq 1$ be a choice sequence. Suppose the functions $\Theta_0(\lambda)$ and $\hat{\Theta}_0(\lambda)$ from the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ have $\{\Gamma_n\}_0^\infty$ as their Schur parameters. Then, for every $n = 0, 1, \dots$, we have the relations

$$\begin{aligned} \Theta_n(\lambda) &= \Gamma_n + \lambda D_{\Gamma_n^*} (I + \lambda \Theta_{n+1}(\lambda) \Gamma_n^*)^{-1} \Theta_{n+1}(\lambda) D_{\Gamma_n}, \\ \hat{\Theta}_n(\lambda) &= \Gamma_n + \lambda D_{\Gamma_n^*} (I + \lambda \hat{\Theta}_{n+1}(\lambda) \Gamma_n^*)^{-1} \hat{\Theta}_{n+1}(\lambda) D_{\Gamma_n}, \quad \lambda \in \mathbb{D}, \end{aligned}$$

where $\{\Theta_n\}$ and $\{\widehat{\Theta}_n\}$ are the Schur iterates of Θ and $\widehat{\Theta}$, respectively. Then one has for every n the equalities

$$(2.3) \quad \begin{aligned} \Theta_n(\lambda) - \widehat{\Theta}_n(\lambda) &= \lambda D_{\Gamma_n^*} (I + \lambda \Theta_{n+1}(\lambda) \Gamma_n^*)^{-1} (\Theta_{n+1}(\lambda) \\ &\quad - \widehat{\Theta}_{n+1}(\lambda)) (I + \lambda \widehat{\Theta}_{n+1}(\lambda) \Gamma_n^*)^{-1} D_{\Gamma_n}, \quad \lambda \in \mathbb{D}. \end{aligned}$$

Since $\|\Theta_{n+1}(\lambda) - \widehat{\Theta}_{n+1}(\lambda)\| \leq 2$ for all $\lambda \in \mathbb{D}$ and $\Theta_{n+1}(0) = \widehat{\Theta}_{n+1}(0) = \Gamma_{n+1}$, by Schwartz's lemma we get $\|\Theta_{n+1}(\lambda) - \widehat{\Theta}_{n+1}(\lambda)\| \leq 2|\lambda|$, $\lambda \in \mathbb{D}$. Further

$$\begin{aligned} \|(I + \lambda \Theta_{n+1}(\lambda) \Gamma_n^*) f\| &\geq (1 - |\lambda|) \|f\|, \\ \|(I + \lambda \widehat{\Theta}_{n+1}(\lambda) \Gamma_n^*) f\| &\geq (1 - |\lambda|) \|f\| \end{aligned}$$

for all $\lambda \in \mathbb{D}$ and for all $f \in \mathfrak{D}_{\Gamma_{n-1}^*}$. These relations imply

$$\|(I + \lambda \Theta_{n+1}(\lambda) \Gamma_n^*)^{-1}\| \leq \frac{1}{1 - |\lambda|}, \quad \|(I + \lambda \widehat{\Theta}_{n+1}(\lambda) \Gamma_n^*)^{-1}\| \leq \frac{1}{1 - |\lambda|}$$

for all $\lambda \in \mathbb{D}$ and for all $n = 0, 1, \dots$. Hence, from (2.3) we have

$$\|\Theta_n(\lambda) - \widehat{\Theta}_n(\lambda)\| \leq 2|\lambda| \frac{|\lambda|}{(1 - |\lambda|)^2}, \quad \lambda \in \mathbb{D}.$$

Then applying (2.3) to Θ_{n-1} and $\widehat{\Theta}_{n-1}$ in the left-hand side, we see that

$$\|\Theta_{n-1}(\lambda) - \widehat{\Theta}_{n-1}(\lambda)\| \leq 2|\lambda| \left(\frac{|\lambda|}{(1 - |\lambda|)^2} \right)^2, \quad \lambda \in \mathbb{D},$$

and finally

$$(2.4) \quad \|\Theta_0(\lambda) - \widehat{\Theta}_0(\lambda)\| \leq 2|\lambda| \left(\frac{|\lambda|}{(1 - |\lambda|)^2} \right)^{n+1}, \quad \lambda \in \mathbb{D},$$

for all $n \in \mathbb{N}_0$. Let $|\lambda| < (3 - \sqrt{5})/2$. Then

$$\frac{|\lambda|}{(1 - |\lambda|)^2} < 1.$$

Letting $n \rightarrow \infty$ in (2.4) we get $\Theta_0(\lambda) = \widehat{\Theta}_0(\lambda)$ for $|\lambda| < (3 - \sqrt{5})/2$. Since Θ_0 and $\widehat{\Theta}_0$ are holomorphic in \mathbb{D} , they are equal on \mathbb{D} . \square

3. CONSERVATIVE DISCRETE-TIME LINEAR SYSTEMS AND THEIR TRANSFER FUNCTIONS

Let \mathfrak{M} , \mathfrak{N} , and \mathfrak{H} be separable Hilbert spaces. A linear system

$$\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}$$

with bounded linear operators A, B, C, D of the form

$$(3.1) \quad \begin{cases} \sigma_k = Ch_k + D\xi_k, \\ h_{k+1} = Ah_k + B\xi_k, \end{cases} \quad k \in \mathbb{N}_0,$$

where $\{\xi_k\} \subset \mathfrak{M}$, $\{\sigma_k\} \subset \mathfrak{N}$, $\{h_k\} \subset \mathfrak{H}$ is called a *discrete time-invariant system*. The Hilbert spaces \mathfrak{M} and \mathfrak{N} are called the input and the output spaces, respectively, and the Hilbert space \mathfrak{H} is called the state space. The operators A, B, C , and D are called the state space operator, the control operator, the observation operator, and the feedthrough operator of τ , respectively. Put

$$U_\tau = \begin{bmatrix} D & C \\ B & A \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array} \rightarrow \begin{array}{c} \mathfrak{N} \\ \oplus \\ \mathfrak{H} \end{array}.$$

If U_τ is contractive, then the corresponding discrete-time system is said to be *passive* [12]. If the operator U_τ is isometric (respect., co-isometric, unitary), then the system is said to be *isometric* (respect., *co-isometric*, *conservative*). Isometric, co-isometric, conservative, and passive discrete time-invariant systems have been studied in [24], [25],

[6], [71], [51], [52], [26], [28], [20], [5], [12], [13], [14], [15], [16], [17], [69], [70], [9], [7], [8], [45]. It is relevant to remark that a brief history of System Theory is presented in the recent preprint of B. Fritzsche, V. Katsnelson, and B. Kirstein [45].

The subspaces

$$(3.2) \quad \mathfrak{H}^c := \overline{\text{span}} \{A^n B \mathfrak{M}, n \in \mathbb{N}_0\} \quad \text{and} \quad \mathfrak{H}^o := \overline{\text{span}} \{A^{*n} C^* \mathfrak{N}, n \in \mathbb{N}_0\}$$

are said to be the *controllable* and *observable* subspaces of the system τ , respectively. The system τ is said to be *controllable* (respect., *observable*) if $\mathfrak{H}^c = \mathfrak{H}$ (respect., $\mathfrak{H}^o = \mathfrak{H}$), and it is called *minimal* if τ is both controllable and observable. The system τ is said to be *simple* if

$$\mathfrak{H} = \text{clos} \{ \mathfrak{H}^c + \mathfrak{H}^o \} = \overline{\text{span}} \{ A^k B \mathfrak{M}, A^{*l} C^* \mathfrak{N}, k, l \in \mathbb{N}_0 \}.$$

It follows from (3.2) that

$$(\mathfrak{H}^c)^\perp = \bigcap_{n=0}^{\infty} \ker(B^* A^{*n}), \quad (\mathfrak{H}^o)^\perp = \bigcap_{n=0}^{\infty} \ker(C A^n),$$

and therefore there are the following alternative characterizations:

- (a) τ is controllable $\iff \bigcap_{n=0}^{\infty} \ker(B^* A^{*n}) = \{0\}$;
- (b) τ is observable $\iff \bigcap_{n=0}^{\infty} \ker(C A^n) = \{0\}$;
- (c) τ is simple $\iff \left(\bigcap_{n=0}^{\infty} \ker(B^* A^{*n}) \right) \cap \left(\bigcap_{n=0}^{\infty} \ker(C A^n) \right) = \{0\}$.

A contraction A acting in a Hilbert space \mathfrak{H} is called *completely non-unitary* [71] if there is no nontrivial reducing subspace of A , on which A generates a unitary operator. Given a contraction A in \mathfrak{H} then there is a canonical orthogonal decomposition [71, Theorem I.3.2]

$$\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}_1, \quad A = A_0 \oplus A_1, \quad A_j = A|_{\mathfrak{H}_j}, \quad j = 0, 1,$$

where \mathfrak{H}_0 and \mathfrak{H}_1 reduce A , the operator A_0 is a completely non-unitary contraction, and A_1 is a unitary operator. Moreover,

$$\mathfrak{H}_1 = \left(\bigcap_{n \geq 1} \ker D_{A^n} \right) \cap \left(\bigcap_{n \geq 1} \ker D_{A^{*n}} \right).$$

Since

$$\bigcap_{k=0}^{n-1} \ker(D_A A^k) = \ker D_{A^n}, \quad \bigcap_{k=0}^{n-1} \ker(D_{A^*} A^{*k}) = \ker D_{A^{*n}},$$

we get

$$(3.3) \quad \begin{aligned} \bigcap_{n \geq 1} \ker D_{A^n} &= \mathfrak{H} \ominus \overline{\text{span}} \{ A^{*n} D_A \mathfrak{H}, n \in \mathbb{N}_0 \}, \\ \bigcap_{n \geq 1} \ker D_{A^{*n}} &= \mathfrak{H} \ominus \overline{\text{span}} \{ A^n D_{A^*} \mathfrak{H}, n \in \mathbb{N}_0 \}. \end{aligned}$$

It follows that

$$(3.4) \quad \begin{aligned} A \text{ is completely non-unitary} &\iff \left(\bigcap_{n \geq 1} \ker D_{A^n} \right) \cap \left(\bigcap_{n \geq 1} \ker D_{A^{*n}} \right) = \{0\} \\ &\iff \overline{\text{span}} \{ A^{*n} D_A, A^m D_{A^*}, n, m \in \mathbb{N}_0 \} = \mathfrak{H}. \end{aligned}$$

If $\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}$ is a conservative system then τ is simple if and only if the state space operator A is a completely non-unitary contraction [28], [20].

The *transfer function*

$$\Theta_\tau(\lambda) := D + \lambda C (I_{\mathfrak{H}} - \lambda A)^{-1} B, \quad \lambda \in \mathbb{D},$$

of a passive system τ belongs to the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ [12]. Conservative systems are also called the unitary colligations and their transfer functions are called the characteristic functions [28].

The examples of conservative systems are given by

$$\Sigma = \left\{ \begin{bmatrix} -A & D_{A^*} \\ D_A & A^* \end{bmatrix}; \mathfrak{D}_A, \mathfrak{D}_{A^*}, \mathfrak{H} \right\}, \quad \Sigma_* = \left\{ \begin{bmatrix} -A^* & D_A \\ D_{A^*} & A \end{bmatrix}; \mathfrak{D}_{A^*}, \mathfrak{D}_A, \mathfrak{H} \right\}.$$

The transfer functions of these systems

$$\Phi_{\Sigma}(\lambda) = (-A + \lambda D_{A^*} (I_{\mathfrak{H}} - \lambda A^*)^{-1} D_A) \upharpoonright \mathfrak{D}_A, \quad \lambda \in \mathbb{D}$$

and

$$\Phi_{\Sigma_*}(\lambda) = (-A^* + \lambda D_A (I_{\mathfrak{H}} - \lambda A)^{-1} D_{A^*}) \upharpoonright \mathfrak{D}_{A^*}, \quad \lambda \in \mathbb{D}$$

are precisely the Sz.-Nagy–Foias characteristic functions [71] of A and A^* , correspondingly.

It is well known that every operator-valued function Θ from the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{N})$ can be realized as a transfer function of some passive system, which can be chosen as controllable isometric (respect., observable co-isometric, simple conservative, minimal passive); cf. [25], [71], [28], [6] [12], [14], [5]. Moreover, two controllable isometric (respect., observable co-isometric, simple conservative) systems with the same transfer function are *unitarily equivalent*: two discrete-time systems

$$\tau_1 = \left\{ \begin{bmatrix} D & C_1 \\ B_1 & A_1 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_1 \right\} \quad \text{and} \quad \tau_2 = \left\{ \begin{bmatrix} D & C_2 \\ B_2 & A_2 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_2 \right\}$$

are said to be unitarily equivalent if there exists a unitary operator V from \mathfrak{H}_1 onto \mathfrak{H}_2 such that

$$(3.5) \quad \begin{aligned} A_1 &= V^{-1} A_2 V, & B_1 &= V^{-1} B_2, & C_1 &= C_2 V \\ &\iff \begin{bmatrix} I_{\mathfrak{M}} & 0 \\ 0 & V \end{bmatrix} \begin{bmatrix} D & C_1 \\ B_1 & A_1 \end{bmatrix} = \begin{bmatrix} D & C_2 \\ B_2 & A_2 \end{bmatrix} \begin{bmatrix} I_{\mathfrak{M}} & 0 \\ 0 & V \end{bmatrix} \end{aligned}$$

cf. [24], [25], [6], [28], [5].

4. CONSERVATIVE REALIZATIONS OF THE SCHUR ITERATES

Let A be a completely non-unitary contraction in a separable Hilbert space \mathfrak{H} . Suppose $\ker D_A \neq \{0\}$. Define the subspaces and operators (see [8])

$$(4.1) \quad \begin{cases} \mathfrak{H}_{0,0} := \mathfrak{H} \\ \mathfrak{H}_{n,0} = \ker D_{A^n}, & \mathfrak{H}_{0,m} := \ker D_{A^{*m}}, \\ \mathfrak{H}_{n,m} := \ker D_{A^n} \cap \ker D_{A^{*m}}, & m, n \in \mathbb{N}, \end{cases}$$

$$(4.2) \quad A_{n,m} := P_{n,m} A \upharpoonright \mathfrak{H}_{n,m} \in \mathbf{L}(\mathfrak{H}_{n,m}),$$

where $P_{n,m}$ are the orthogonal projections in \mathfrak{H} onto $\mathfrak{H}_{n,m}$. The next results have been established in [8].

Theorem 4.1. [8]. *The operators $\{A_{n,m}\}$ are completely non-unitary contractions and the following relations are valid:*

$$(4.3) \quad \begin{aligned} \ker D_{A_{n,m}^k} &= \mathfrak{H}_{n+k,m}, & \ker D_{A_{n,m}^{*k}} &= \mathfrak{H}_{n,m+k}, & m, n \in \mathbb{N}, & k \in \mathbb{N}, \\ \begin{cases} A \mathfrak{H}_{n,m} = \mathfrak{H}_{n-1,m+1}, & n \in \mathbb{N}, & m \in \mathbb{N}_0, \\ A^* \mathfrak{H}_{n,m} = \mathfrak{H}_{n+1,m-1}, & m \in \mathbb{N}, & n \in \mathbb{N}_0, \end{cases} \\ (A_{n,m})_{k,l} &= A_{n+k,m+l}, & n, m \in \mathbb{N}_0, & k, l \in \mathbb{N}, \\ A_{n-1,m+1} A f &= A A_{n,m} f, & f \in \mathfrak{H}_{n,m}, & n \in \mathbb{N}, & n \in \mathbb{N}_0. \end{aligned}$$

Therefore, the operators

$$A_{n,0}, A_{n-1,1}, \dots, A_{n-k,k}, \dots, A_{0,n}, \quad n \in \mathbb{N},$$

are unitarily equivalent.

The relation (4.3) yields the following picture for the creation of the operators $A_{n,m}$:
The process terminates at the N -th step if and only if

$$\begin{aligned} \ker D_{A^N} = \{0\} &\iff \ker D_{A^{N-1}} \cap \ker D_{A^*} = \{0\} \iff \dots \\ \ker D_{A^{N-k}} \cap \ker D_{A^{*k}} = \{0\} &\iff \dots \ker D_{A^{*N}} = \{0\}. \end{aligned}$$

Theorem 4.2. [8]. *Let A be a completely non-unitary contraction in a separable Hilbert space \mathfrak{H} . Assume $\ker D_A \neq \{0\}$ and let the contractions $A_{n,m}$ be defined by (4.1) and (4.2). Then for each $n \in \mathbb{N}$ the characteristic functions of the operators*

$$A_{n,0}, A_{n-1,1}, \dots, A_{n-m,m}, \dots, A_{1,n-1}, A_{0,n}$$

coincide with the pure part of the n -th Schur iterate of the Sz.-Nagy–Foias [71] characteristic function $\Phi_A(\lambda) = (-A + \lambda D_{A^*}(I_{\mathfrak{H}} - \lambda A^*)^{-1} D_A) \upharpoonright \mathfrak{D}_A$ of A . Moreover, each operator from the set $\{A_{n-k,k}\}_{k=0}^n$ is

- (1) a unilateral shift (respect., co-shift) if and only if the n -th Schur parameter Γ_n of Φ is isometric (respect., co-isometric),
- (2) the orthogonal sum of a unilateral shift and co-shift if and only if

$$(4.4) \quad \mathfrak{D}_{\Gamma_{n-1}} \neq \{0\}, \mathfrak{D}_{\Gamma_{n-1}^*} \neq \{0\} \quad \text{and} \quad \Gamma_m = 0 \quad \text{for all} \quad m \geq n.$$

Each subspace from the set $\{\mathfrak{H}_{n-k,k}\}_{k=0}^n$ is trivial if and only if Γ_n is unitary.

Notice that constructions of simple conservative realizations for the Schur iterates of a function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ by means of a simple conservative realization

$$\tau_0 = \left\{ \begin{bmatrix} \Gamma_0 & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}$$

are given in [8]. We mention the following equivalences:

the n -th Schur parameter Γ_n of Θ is isometric (respect., co-isometric) \iff each operator from the set $\{A_{n-k,k}\}_{k=0}^n$ is a co-shift (respect., a shift).

The statement below is established in [8] and is needed in the sequel.

Theorem 4.3. [8]. *Let $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$, $\Gamma_0 = \Theta(0)$, and let*

$$\tau_0 = \left\{ \begin{bmatrix} \Gamma_0 & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}$$

be a simple conservative realization of Θ . Then the systems

$$(4.5) \quad \begin{aligned} \zeta_{0,1} &= \left\{ \begin{bmatrix} D_{\Gamma_0^*}^{-1} C (D_{\Gamma_0}^{-1} B^*)^* & D_{\Gamma_0^*}^{-1} C \\ AP_{1,0} D_{A^*}^{-1} B & A_{0,1} \end{bmatrix}; \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \mathfrak{H}_{0,1} \right\}, \\ \zeta_{1,0} &= \left\{ \begin{bmatrix} D_{\Gamma_0^*}^{-1} C (D_{\Gamma_0}^{-1} B^*)^* & D_{\Gamma_0^*}^{-1} C A \\ P_{1,0} D_{A^*}^{-1} B & A_{1,0} \end{bmatrix}; \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \mathfrak{H}_{1,0} \right\} \end{aligned}$$

are unitarily equivalent, conservative and simple and their transfer function is equal to the first Schur iterate Θ_1 of Θ .

Here the operators $D_{\Gamma_0}^{-1}$, $D_{\Gamma_0^*}^{-1}$, and $D_{A^*}^{-1}$ are the Moore–Penrose pseudo-inverses. In the sequel the transformations of the conservative system

$$\tau \mapsto \zeta_{0,1}, \quad \tau \mapsto \zeta_{1,0}$$

will be denoted by $\Omega_{0,1}(\tau)$ and $\Omega_{1,0}(\tau)$, respectively.

Remark 4.4. The problem of isometric, co-isometric, and conservative realizations of the Schur iterates for a scalar function from the generalized Schur class has been studied in [2], [3], [4]. For a scalar finite Blaschke product the realizations of the Schur iterates are constructed in [45].

5. BLOCK OPERATOR CMV MATRICES AND CONSERVATIVE REALIZATIONS OF THE SCHUR CLASS FUNCTION (THE CASE WHEN THE OPERATOR Γ_n IS NEITHER AN ISOMETRY NOR A CO-ISOMETRY FOR EACH n)

Let

$$\Gamma_0 \in \mathbf{L}(\mathfrak{M}, \mathfrak{N}), \quad \Gamma_n \in \mathbf{L}(\mathfrak{D}_{\Gamma_{n-1}}, \mathfrak{D}_{\Gamma_{n-1}^*}), \quad n \in \mathbb{N},$$

be a choice sequence. In this and next Section 6 we are going to construct by means of $\{\Gamma_n\}_{n \geq 0}$ two unitary equivalent simple conservative systems with such a transfer function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ that $\{\Gamma_n\}_{n \geq 0}$ are its Schur parameters. In particular, this leads to the existence part of Theorem 1.2 and to the well known result that any $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ admits a realization as a transfer function of a simple conservative system. We begin with constructions of block operator CMV matrices for a given choice sequence $\{\Gamma_n\}_{n \geq 0}$ and will suppose that all the operators Γ_n are neither isometries nor co-isometries. We will use the well known constructions of finite and infinite orthogonal sums of Hilbert spaces. Namely, if $\{H_k\}_{k=1}^\infty$ is a given sequence of Hilbert spaces, then

$$\mathfrak{H} = \sum_{k=1}^N \bigoplus H_k$$

is a Hilbert space with the inner product $(f, g) = \sum_{k=0}^N (f_k, g_k)_{H_k}$ for $f = (f_1, \dots, f_N)^T$

and $g = (g_1, \dots, g_N)^T$, $f_k, g_k \in H_k$, $k = 1, \dots, N$ and the norm $\|f\|^2 = \sum_{k=0}^N \|f_k\|_{H_k}^2$. The Hilbert space

$$\mathfrak{H} = \sum_{k=0}^\infty \bigoplus H_k$$

consists of all vectors of the form $f = (f_1, f_2, \dots)^T$, $f_k \in H_k$, $k = 1, 2, \dots$, such that

$$\|f\|^2 = \sum_{k=1}^\infty \|f_k\|_{H_k}^2 < \infty.$$

The inner product is given by $(f, g) = \sum_{k=1}^\infty (f_k, g_k)_{H_k}$.

5.1. Block operator CMV matrices. Define the Hilbert spaces

$$(5.1) \quad \begin{aligned} \mathfrak{H}_0 &= \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 0}) := \sum_{n \geq 0} \bigoplus \begin{matrix} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{matrix}, \\ \tilde{\mathfrak{H}}_0 &= \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 0}) := \sum_{n \geq 0} \bigoplus \begin{matrix} \mathfrak{D}_{\Gamma_{2n}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}} \end{matrix}. \end{aligned}$$

From these definitions it follows that

$$\tilde{\mathfrak{H}}_0(\{\Gamma_n^*\}_{n \geq 0}) = \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 0}), \quad \mathfrak{H}_0(\{\Gamma_n^*\}_{n \geq 0}) = \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 0}).$$

The spaces $\mathfrak{N} \bigoplus \mathfrak{H}_0$ and $\mathfrak{M} \bigoplus \tilde{\mathfrak{H}}_0$ represent in the form

$$\begin{aligned} \mathfrak{N} \bigoplus \mathfrak{H}_0 &= \bigoplus_{\mathfrak{D}_{\Gamma_0}} \bigoplus \sum_{n \geq 1} \bigoplus \begin{matrix} \mathfrak{D}_{\Gamma_{2n-1}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n}} \end{matrix}, \\ \mathfrak{M} \bigoplus \tilde{\mathfrak{H}}_0 &= \bigoplus_{\mathfrak{D}_{\Gamma_0^*}} \bigoplus \sum_{n \geq 1} \bigoplus \begin{matrix} \mathfrak{D}_{\Gamma_{2n-1}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n}^*} \end{matrix}. \end{aligned}$$

Let

$$\mathbf{J}_{\Gamma_0} = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \\ D_{\Gamma_0} & -\Gamma_0^* \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \mathfrak{D}_{\Gamma_0^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{N} \\ \mathfrak{D}_{\Gamma_0} \end{array},$$

$$\mathbf{J}_{\Gamma_k} = \begin{bmatrix} \Gamma_k & D_{\Gamma_k^*} \\ D_{\Gamma_k} & -\Gamma_k^* \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{k-1}} \\ \mathfrak{D}_{\Gamma_k^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{k-1}^*} \\ \mathfrak{D}_{\Gamma_k} \end{array},$$

$$k \in \mathbb{N}.$$

be the elementary rotations. Define the following unitary operators

$$(5.2) \quad \begin{aligned} \mathcal{M}_0 &= \mathcal{M}_0(\{\Gamma_n\}_{n \geq 0}) := I_{\mathfrak{M}} \bigoplus \sum_{n \geq 1} \bigoplus \mathbf{J}_{\Gamma_{2n-1}} : \mathfrak{M} \bigoplus \mathfrak{H}_0 \rightarrow \mathfrak{M} \bigoplus \tilde{\mathfrak{H}}_0, \\ \tilde{\mathcal{M}}_0 &= \tilde{\mathcal{M}}_0(\{\Gamma_n\}_{n \geq 0}) := I_{\mathfrak{N}} \bigoplus \sum_{n \geq 1} \bigoplus \mathbf{J}_{\Gamma_{2n-1}} : \mathfrak{N} \bigoplus \mathfrak{H}_0 \rightarrow \mathfrak{N} \bigoplus \tilde{\mathfrak{H}}_0, \\ \mathcal{L}_0 &= \mathcal{L}_0(\{\Gamma_n\}_{n \geq 0}) := \mathbf{J}_{\Gamma_0} \bigoplus \sum_{n \geq 1} \bigoplus \mathbf{J}_{\Gamma_{2n}} : \mathfrak{M} \bigoplus \tilde{\mathfrak{H}}_0 \rightarrow \mathfrak{N} \bigoplus \mathfrak{H}_0. \end{aligned}$$

Observe that $(\mathcal{L}_0(\{\Gamma_n\}_{n \geq 0}))^* = \mathcal{L}_0(\{\Gamma_n^*\}_{n \geq 0})$. Let

$$(5.3) \quad \mathcal{V}_0 = \mathcal{V}_0(\{\Gamma_n\}_{n \geq 0}) := \sum_{n \geq 1} \bigoplus \mathbf{J}_{\Gamma_{2n-1}} : \mathfrak{H}_0 \rightarrow \tilde{\mathfrak{H}}_0.$$

Clearly, the operator \mathcal{V}_0 is unitary and

$$(5.4) \quad \mathcal{M}_0 = I_{\mathfrak{M}} \bigoplus \mathcal{V}_0, \quad \tilde{\mathcal{M}}_0 = I_{\mathfrak{N}} \bigoplus \mathcal{V}_0.$$

It follows that $(\tilde{\mathcal{M}}_0(\{\Gamma_n\}_{n \geq 0}))^* = \mathcal{M}_0(\{\Gamma_n^*\}_{n \geq 0})$, $(\mathcal{M}_0(\{\Gamma_n\}_{n \geq 0}))^* = \tilde{\mathcal{M}}_0(\{\Gamma_n^*\}_{n \geq 0})$. Finally, define the unitary operators

$$(5.5) \quad \begin{aligned} \mathcal{U}_0 &= \mathcal{U}_0(\{\Gamma_n\}_{n \geq 0}) := \mathcal{L}_0 \mathcal{M}_0 : \mathfrak{M} \bigoplus \mathfrak{H}_0 \rightarrow \mathfrak{N} \bigoplus \mathfrak{H}_0, \\ \tilde{\mathcal{U}}_0 &= \tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0}) := \tilde{\mathcal{M}}_0 \mathcal{L}_0 : \mathfrak{M} \bigoplus \tilde{\mathfrak{H}}_0 \rightarrow \mathfrak{N} \bigoplus \tilde{\mathfrak{H}}_0. \end{aligned}$$

Calculations give

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 & -\Gamma_0^* D_{\Gamma_1^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & \Gamma_2 D_{\Gamma_1} & -\Gamma_2 \Gamma_1^* & D_{\Gamma_2^*} \Gamma_3 & D_{\Gamma_2^*} D_{\Gamma_3^*} & 0 & 0 & 0 & \dots \\ 0 & D_{\Gamma_2} D_{\Gamma_1} & -D_{\Gamma_2} \Gamma_1^* & -\Gamma_2^* \Gamma_3 & -\Gamma_2^* D_{\Gamma_3^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & \Gamma_4 D_{\Gamma_3} & -\Gamma_4 \Gamma_3^* & D_{\Gamma_4^*} \Gamma_5 & D_{\Gamma_4^*} D_{\Gamma_5^*} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

and

$$\tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & D_{\Gamma_1^*} D_{\Gamma_2^*} & 0 & 0 & 0 & \dots \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & -\Gamma_1^* D_{\Gamma_2^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & \Gamma_3 D_{\Gamma_2} & -\Gamma_3 \Gamma_2^* & D_{\Gamma_3^*} \Gamma_4 & D_{\Gamma_3^*} D_{\Gamma_4^*} & 0 & \dots \\ 0 & 0 & D_{\Gamma_3} D_{\Gamma_2} & -D_{\Gamma_3} \Gamma_2^* & -\Gamma_3^* \Gamma_4 & -\Gamma_3^* D_{\Gamma_4^*} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

Let

$$\mathcal{C}_0 = \begin{bmatrix} D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_0} \\ \mathfrak{D}_{\Gamma_1^*} \end{array} \rightarrow \mathfrak{N}, \quad \mathcal{A}_0 = \begin{bmatrix} D_{\Gamma_0} \\ 0 \end{bmatrix} : \mathfrak{M} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_0} \\ \mathfrak{D}_{\Gamma_1^*} \end{array},$$

$$(5.6) \quad \begin{cases} \mathcal{B}_n = \begin{bmatrix} -\Gamma_{2n-2}^* \Gamma_{2n-1} & -\Gamma_{2n-2}^* D_{\Gamma_{2n-1}^*} \\ \Gamma_{2n} D_{\Gamma_{2n-1}} & -\Gamma_{2n} \Gamma_{2n-1}^* \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}^*} \end{array}, \\ \mathcal{C}_n = \begin{bmatrix} 0 & 0 \\ D_{\Gamma_{2n}^*} \Gamma_{2n+1} & D_{\Gamma_{2n}^*} D_{\Gamma_{2n+1}^*} \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}^*} \end{array}, \\ \mathcal{A}_n = \begin{bmatrix} D_{\Gamma_{2n}} D_{\Gamma_{2n-1}} & -D_{\Gamma_{2n}} \Gamma_{2n-1}^* \\ 0 & 0 \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}^*} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \\ \tilde{\mathcal{C}}_0 = [D_{\Gamma_0^*} \ 0] : \begin{array}{c} \mathfrak{D}_{\Gamma_0^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_1} \end{array} \rightarrow \mathfrak{N}, \quad \tilde{\mathcal{A}}_0 = \begin{bmatrix} \Gamma_1 D_{\Gamma_0} \\ D_{\Gamma_1} D_{\Gamma_0} \end{bmatrix} : \mathfrak{M} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_0^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_1} \end{array}, \end{cases}$$

$$(5.7) \quad \begin{cases} \tilde{\mathcal{B}}_n = \begin{bmatrix} -\Gamma_{2n-1} \Gamma_{2n-2}^* & D_{\Gamma_{2n-1}^*} \Gamma_{2n} \\ -D_{\Gamma_{2n-1}} \Gamma_{2n-2}^* & -\Gamma_{2n-1}^* \Gamma_{2n} \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}} \end{array}, \\ \tilde{\mathcal{C}}_n = \begin{bmatrix} D_{\Gamma_{2n-1}^*} D_{\Gamma_{2n}^*} & 0 \\ -\Gamma_{2n-1}^* D_{\Gamma_{2n}^*} & 0 \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}} \end{array}, \\ \tilde{\mathcal{A}}_n = \begin{bmatrix} 0 & \Gamma_{2n+1} D_{\Gamma_{2n}} \\ 0 & D_{\Gamma_{2n+1}} D_{\Gamma_{2n}} \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{2n-2}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n-1}} \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}} \end{array}. \end{cases}$$

It is easy to see that the operators \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ take the following three-diagonal block operator matrix form

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & \mathcal{C}_0 & 0 & 0 & 0 & \cdot & \cdot \\ \mathcal{A}_0 & \mathcal{B}_1 & \mathcal{C}_1 & 0 & 0 & \cdot & \cdot \\ 0 & \mathcal{A}_1 & \mathcal{B}_2 & \mathcal{C}_2 & 0 & \cdot & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots \end{bmatrix}, \quad \tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & \tilde{\mathcal{C}}_0 & 0 & 0 & 0 & \cdot & \cdot \\ \tilde{\mathcal{A}}_0 & \tilde{\mathcal{B}}_1 & \tilde{\mathcal{C}}_1 & 0 & 0 & \cdot & \cdot \\ 0 & \tilde{\mathcal{A}}_1 & \tilde{\mathcal{B}}_2 & \tilde{\mathcal{C}}_2 & 0 & \cdot & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots \end{bmatrix}.$$

The block operator matrices \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ will be called *block operator CMV matrices*. Observe that

$$(5.8) \quad \tilde{\mathcal{M}}_0 \mathcal{U}_0 = \tilde{\mathcal{U}}_0 \mathcal{M}_0,$$

and

$$(5.9) \quad (\mathcal{U}_0(\{\Gamma_n\}_{n \geq 0}))^* = \tilde{\mathcal{U}}_0(\{\Gamma_n^*\}_{n \geq 0}), \quad (\tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0}))^* = \mathcal{U}_0(\{\Gamma_n^*\}_{n \geq 0}).$$

Therefore the matrix $\tilde{\mathcal{U}}_0$ can be obtained from \mathcal{U}_0 by passing to the adjoint \mathcal{U}_0^* and then by replacing Γ_n (respect., Γ_n^*) by Γ_n^* (respect., Γ_n) for all n . In the case when the choice sequence consists of complex numbers from the unit disk the matrix $\tilde{\mathcal{U}}_0$ is the transpose to \mathcal{U}_0 , i.e., $\tilde{\mathcal{U}}_0 = \mathcal{U}_0^t$.

Remark 5.1. The three-diagonal block form of the CMV matrices with scalar entries has been established in [22] (see also [43]).

5.2. Truncated block operator CMV matrices. Define two contractions,

$$(5.10) \quad \mathcal{T}_0 = \mathcal{T}_0(\{\Gamma_n\}_{n \geq 0}) := P_{\mathfrak{H}_0} \mathcal{U}_0 \upharpoonright \mathfrak{H}_0 : \mathfrak{H}_0 \rightarrow \mathfrak{H}_0,$$

$$(5.11) \quad \tilde{\mathcal{T}}_0 = \tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 0}) := P_{\tilde{\mathfrak{H}}_0} \tilde{\mathcal{U}}_0 \upharpoonright \tilde{\mathfrak{H}}_0 : \tilde{\mathfrak{H}}_0 \rightarrow \tilde{\mathfrak{H}}_0.$$

The operators \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$ take on the three-diagonal block operator matrix forms

$$\mathcal{T}_0 = \begin{bmatrix} \mathcal{B}_1 & \mathcal{C}_1 & 0 & 0 & 0 & \cdot \\ \mathcal{A}_1 & \mathcal{B}_2 & \mathcal{C}_2 & 0 & 0 & \cdot \\ 0 & \mathcal{A}_2 & \mathcal{B}_3 & \mathcal{C}_3 & 0 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad \tilde{\mathcal{T}}_0 = \begin{bmatrix} \tilde{\mathcal{B}}_1 & \tilde{\mathcal{C}}_1 & 0 & 0 & 0 & \cdot \\ \tilde{\mathcal{A}}_1 & \tilde{\mathcal{B}}_2 & \tilde{\mathcal{C}}_2 & 0 & 0 & \cdot \\ 0 & \tilde{\mathcal{A}}_2 & \tilde{\mathcal{B}}_3 & \tilde{\mathcal{C}}_3 & 0 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

where $\mathcal{A}_n, \mathcal{B}_n, \mathcal{C}_n, \tilde{\mathcal{A}}_n, \tilde{\mathcal{B}}_n$, and $\tilde{\mathcal{C}}_n$ are given by (5.6) and (5.7). Since the matrices \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$ are obtained from \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ by deleting the first rows and the first columns, we will call them *truncated block operator CMV matrices*. Observe that from the definitions of $\mathcal{L}_0, \mathcal{M}_0, \mathcal{M}_0, \mathcal{T}_0$, and $\tilde{\mathcal{T}}_0$ it follows that \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$ are products of two block-diagonal matrices

$$(5.12) \quad \mathcal{T}_0 = \mathcal{T}_0(\{\Gamma_n\}_{n \geq 0}) = (-\Gamma_0^* \oplus \mathbf{J}_{\Gamma_2} \oplus \cdots \oplus \mathbf{J}_{\Gamma_{2n}} \oplus \cdots) (\mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \cdots \oplus \mathbf{J}_{\Gamma_{2n-1}} \oplus \cdots)$$

(5.13)

$$\tilde{\mathcal{T}}_0 = \tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 0}) = (\mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \cdots \oplus \mathbf{J}_{\Gamma_{2n-1}} \oplus \cdots) (-\Gamma_0^* \oplus \mathbf{J}_{\Gamma_2} \oplus \cdots \oplus \mathbf{J}_{\Gamma_{2n}} \oplus \cdots).$$

In particular, it follows that

$$(5.14) \quad (\mathcal{T}_0(\{\Gamma_n\}_{n \geq 0}))^* = \tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0}).$$

From (5.12) and (5.13) we have $\mathcal{V}_0 \mathcal{T}_0 = \tilde{\mathcal{T}}_0 \mathcal{V}_0$, where the unitary operator \mathcal{V}_0 is defined by (5.3). Therefore, the operators \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$ are unitarily equivalent.

Proposition 5.2. *Let $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ and let $\{\Gamma_n\}_{n \geq 0}$ be the Schur parameters of Θ . Suppose Γ_n is neither isometric nor co-isometric for each n . Let the function $\Omega \in \mathbf{S}(\mathfrak{K}, \mathfrak{L})$ coincide with Θ and let $\{G_n\}_{n \geq 0}$ be the Schur parameters of Ω . Then the truncated block operator CMV matrices $\mathcal{T}_0(\{\Gamma_n\}_{n \geq 0})$ and $\mathcal{T}_0(\{G_n\}_{n \geq 0})$ (respect., $\tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 0})$ and $\tilde{\mathcal{T}}_0(\{G_n\}_{n \geq 0})$) are unitarily equivalent.*

Proof. Since $\Omega(\lambda) = V\Theta(\lambda)U$, where $U \in \mathbf{L}(\mathfrak{K}, \mathfrak{M})$ and $V \in \mathbf{L}(\mathfrak{N}, \mathfrak{L})$ are unitary operators, we get relations (2.2). It follows that $\mathfrak{D}_{G_n} \neq \{0\}$ and $\mathfrak{D}_{G_n^*} \neq \{0\}$ for all n . Hence, we have

$$(5.15) \quad \mathbf{J}_{G_n} \begin{bmatrix} U^* & 0 \\ 0 & V \end{bmatrix} = \begin{bmatrix} V & 0 \\ 0 & U^* \end{bmatrix} \mathbf{J}_{\Gamma_n}, \quad n = 0, 1, \dots$$

Define the Hilbert space

$$\mathfrak{H}_0^\Omega = \mathfrak{H}_0(\{G_n\}_{n \geq 0}) := \sum_{n \geq 0} \bigoplus \begin{matrix} \mathfrak{D}_{G_{2n}} \\ \oplus \\ \mathfrak{D}_{G_{2n+1}^*} \end{matrix}$$

and the truncated block operator CMV matrix

$$\mathcal{T}_0(\{G_n\}_{n \geq 0}) := \begin{bmatrix} -G_0^* & & & & & \\ & \mathbf{J}_{G_2} & & & & \\ & & \mathbf{J}_{G_4} & & & \\ & & & \ddots & & \\ & & & & \mathbf{J}_{G_{2n}} & \\ & & & & & \ddots \end{bmatrix} \begin{bmatrix} \mathbf{J}_{G_1} & & & & & \\ & \mathbf{J}_{G_3} & & & & \\ & & \ddots & & & \\ & & & \mathbf{J}_{G_{2n+1}} & & \\ & & & & \ddots & \end{bmatrix},$$

Define the unitary operator

$$\mathcal{W} = \begin{bmatrix} U^* & & & & \\ & V & & & \\ & & U^* & & \\ & & & V & \\ & & & & \ddots \end{bmatrix} : \mathfrak{H}_0 \rightarrow \mathfrak{H}_0^\Omega.$$

From (5.12) and (5.15) we obtain $\mathcal{W}\mathcal{T}_0(\{\Gamma_n\}_{n \geq 0}) = \mathcal{T}(\{G_n\}_{n \geq 0})\mathcal{W}$. Thus, $\mathcal{T}_0(\{\Gamma_n\}_{n \geq 0})$ and $\mathcal{T}(\{G_n\}_{n \geq 0})$ are unitarily equivalent. \square

Now we are going to find the defect operators and defect subspaces for \mathcal{T}_0 and $\tilde{\mathcal{T}}_0$. Let $\mathbf{f} = (\vec{f}_0, \vec{f}_1, \dots)^T \in \mathfrak{H}_0$, where

$$\vec{f}_n = \begin{bmatrix} h_n \\ g_n \end{bmatrix} \in \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \quad n \in \mathbb{N}_0.$$

Then

$$(5.16) \quad \begin{aligned} \|\mathbf{f}\|^2 - \|\mathcal{T}_0\mathbf{f}\|^2 &= \|P_{\mathfrak{M}}\mathcal{U}_0\mathbf{f}\|^2 = \left\| \mathcal{C}_0 \begin{bmatrix} h_0 \\ g_0 \end{bmatrix} \right\|^2 = \|D_{\Gamma_0^*}(\Gamma_1 h_0 + D_{\Gamma_1^*} g_0)\|^2, \\ \|\mathbf{f}\|^2 - \|\mathcal{T}_0^*\mathbf{f}\|^2 &= \|P_{\mathfrak{M}}\mathcal{U}_0^*\mathbf{f}\|^2 = \left\| \mathcal{A}_0^* \begin{bmatrix} h_0 \\ g_0 \end{bmatrix} \right\|^2 = \|D_{\Gamma_0} h_0\|^2. \end{aligned}$$

Let $\mathbf{x} = (x_0, x_1, \dots)^T \in \tilde{\mathfrak{H}}_0$, where

$$x_n = \begin{bmatrix} h_n \\ g_n \end{bmatrix} \in \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \quad n \in \mathbb{N}_0.$$

Then

$$\begin{aligned} \|\mathbf{x}\|^2 - \|\tilde{\mathcal{T}}_0\mathbf{x}\|^2 &= \|P_{\tilde{\mathfrak{M}}}\tilde{\mathcal{U}}_0\mathbf{x}\|^2 = \left\| \tilde{\mathcal{C}}_0 \begin{bmatrix} h_0 \\ g_0 \end{bmatrix} \right\|^2 = \|D_{\Gamma_0^*} h_0\|^2, \\ \|\mathbf{x}\|^2 - \|\tilde{\mathcal{T}}_0^*\mathbf{x}\|^2 &= \|P_{\tilde{\mathfrak{M}}}\tilde{\mathcal{U}}_0^*\mathbf{x}\|^2 = \left\| \tilde{\mathcal{A}}_0^* \begin{bmatrix} h_0 \\ g_0 \end{bmatrix} \right\|^2 = \|D_{\Gamma_0}(\Gamma_1^* h_0 + D_{\Gamma_1} g_0)\|^2. \end{aligned}$$

Now from Proposition 2.1 it follows that

$$(5.17) \quad \left\{ \begin{array}{l} \ker D_{\mathcal{T}_0} = \left\{ \begin{bmatrix} D_{\Gamma_1}\varphi \\ -\Gamma_1\varphi \end{bmatrix}, \varphi \in \mathfrak{D}_{\Gamma_1} \right\} \oplus \sum_{n \geq 1} \oplus \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \\ \ker D_{\mathcal{T}_0^*} = \mathfrak{D}_{\Gamma_1^*} \oplus \sum_{n \geq 1} \oplus \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \\ \mathfrak{D}_{\mathcal{T}_0} = \left\{ \begin{bmatrix} \Gamma_1^*\psi \\ D_{\Gamma_1^*}\psi \end{bmatrix}, \psi \in \mathfrak{D}_{\Gamma_0^*} \right\} \oplus \vec{0}, \vec{0} \in \sum_{n \geq 1} \oplus \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \\ \mathfrak{D}_{\mathcal{T}_0^*} = \mathfrak{D}_{\Gamma_0} \oplus \vec{0}, \vec{0} \in \mathfrak{D}_{\Gamma_1^*} \oplus \sum_{n \geq 1} \oplus \begin{array}{c} \mathfrak{D}_{\Gamma_{2n}} \\ \oplus \\ \mathfrak{D}_{\Gamma_{2n+1}^*} \end{array}, \end{array} \right.$$

$$(5.18) \quad \left\{ \begin{array}{l} \ker D_{\tilde{\mathcal{T}}_0} = \mathfrak{D}_{\Gamma_1} \oplus \sum_{n \geq 1} \oplus \oplus_{\mathfrak{D}_{\Gamma_{2n+1}^*}} \mathfrak{D}_{\Gamma_{2n}^*}, \\ \ker D_{\tilde{\mathcal{T}}_0^*} = \left\{ \begin{bmatrix} D_{\Gamma_1^*} \varphi \\ -\Gamma_1^* \varphi \end{bmatrix}, \varphi \in \mathfrak{D}_{\Gamma_0^*} \right\} \oplus \sum_{n \geq 1} \oplus \oplus_{\mathfrak{D}_{\Gamma_{2n+1}^*}} \mathfrak{D}_{\Gamma_{2n}^*}, \\ \mathfrak{D}_{\tilde{\mathcal{T}}_0} = \mathfrak{D}_{\Gamma_0} \oplus \vec{0}, \vec{0} \in \mathfrak{D}_{\Gamma_1} \oplus \sum_{n \geq 1} \oplus \oplus_{\mathfrak{D}_{\Gamma_{2n+1}^*}} \mathfrak{D}_{\Gamma_{2n}^*}, \\ \mathfrak{D}_{\tilde{\mathcal{T}}_0^*} = \left\{ \begin{bmatrix} \Gamma_1 \psi \\ D_{\Gamma_1} \psi \end{bmatrix}, \psi \in \mathfrak{D}_{\Gamma_0} \right\} \oplus \vec{0}, \vec{0} \in \sum_{n \geq 1} \oplus \oplus_{\mathfrak{D}_{\Gamma_{2n+1}^*}} \mathfrak{D}_{\Gamma_{2n}^*}. \end{array} \right.$$

5.3. Simple conservative realizations of the Schur class function by means of its Schur parameters. Let

$$\begin{aligned} \mathcal{G}_0 &= \mathcal{G}_0(\{\Gamma_n\}_{n \geq 0}) := [D_{\Gamma_0^*} \Gamma_1 \quad D_{\Gamma_0^*} D_{\Gamma_1^*} \quad 0 \quad 0 \quad \dots] \\ &= D_{\Gamma_0^*} [\Gamma_1 \quad D_{\Gamma_1^*} \quad 0 \quad 0 \quad \dots] : \mathfrak{H}_0 \rightarrow \mathfrak{N}, \\ \tilde{\mathcal{G}}_0 &= \tilde{\mathcal{G}}_0(\{\Gamma_n\}_{n \geq 0}) := [D_{\Gamma_0^*} \quad 0 \quad 0 \quad \dots] = D_{\Gamma_0^*} [I_{\mathfrak{M}} \quad 0 \quad 0 \quad 0 \quad \dots] : \tilde{\mathfrak{H}}_0 \rightarrow \mathfrak{N}, \\ \mathcal{F}_0 &= \mathcal{F}_0(\{\Gamma_n\}_{n \geq 0}) := \begin{bmatrix} D_{\Gamma_0} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{M} \rightarrow \mathfrak{H}_0, \\ \tilde{\mathcal{F}}_0 &= \tilde{\mathcal{F}}_0(\{\Gamma_n\}_{n \geq 0}) := \begin{bmatrix} \Gamma_1 D_{\Gamma_0} \\ D_{\Gamma_1} D_{\Gamma_0} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{M} \rightarrow \tilde{\mathfrak{H}}_0. \end{aligned}$$

The operators \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ can be represented by 2×2 block operator matrices

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & \mathcal{G}_0 \\ \mathcal{F}_0 & \mathcal{T}_0 \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \end{array} \rightarrow \begin{array}{c} \mathfrak{N} \\ \oplus \end{array}, \quad \tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & \tilde{\mathcal{G}}_0 \\ \tilde{\mathcal{F}}_0 & \tilde{\mathcal{T}}_0 \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \end{array} \rightarrow \begin{array}{c} \mathfrak{N} \\ \oplus \end{array}.$$

Define the following conservative systems

$$(5.19) \quad \begin{aligned} \zeta_0 &= \left\{ \begin{bmatrix} \Gamma_0 & \mathcal{G}_0 \\ \mathcal{F}_0 & \mathcal{T}_0 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_0 \right\} = \{\mathcal{U}_0(\{\Gamma_n\}_{n \geq 0}); \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 0})\}, \\ \tilde{\zeta}_0 &= \left\{ \begin{bmatrix} \Gamma_0 & \tilde{\mathcal{G}}_0 \\ \tilde{\mathcal{F}}_0 & \tilde{\mathcal{T}}_0 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \tilde{\mathfrak{H}}_0 \right\} = \{\tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0}); \mathfrak{M}, \mathfrak{N}, \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 0})\}. \end{aligned}$$

Equalities (5.4) and (5.8) yield that systems ζ_0 and $\tilde{\zeta}_0$ are unitarily equivalent. Hence, ζ_0 and $\tilde{\zeta}_0$ have equal transfer functions.

Theorem 5.3. *The unitarily equivalent conservative systems ζ_0 and $\tilde{\zeta}_0$ given by (5.19) are simple and the Schur parameters of the transfer function of ζ_0 and $\tilde{\zeta}_0$ are $\{\Gamma_n\}_{n \geq 0}$.*

Proof. The main step is a proof that the systems $\Omega_{0,1}(\zeta_0)$ and $\Omega_{1,0}(\tilde{\zeta}_0)$ given by (4.5) take the form

$$(5.20) \quad \begin{aligned} \Omega_{0,1}(\zeta_0) &= \left\{ \tilde{U}_0(\{\Gamma_n\}_{n \geq 1}), \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 1}) \right\}, \\ \Omega_{1,0}(\tilde{\zeta}_0) &= \left\{ U_0(\{\Gamma_n\}_{n \geq 1}), \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 1}) \right\}. \end{aligned}$$

First of all we will prove that the systems ζ_0 and $\tilde{\zeta}_0$ are simple.

Define the subspaces

$$\mathfrak{H}_{2k-1} = \sum_{n \geq k} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n-1}^*} \\ \mathfrak{D}_{\Gamma_{2n}}} , \quad \mathfrak{H}_{2k} = \sum_{n \geq k} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n}} \\ \mathfrak{D}_{\Gamma_{2n+1}^*}} , \quad k \in \mathbb{N}.$$

Clearly, $\mathfrak{H}_0 \supset \mathfrak{H}_1 \supset \mathfrak{H}_2 \supset \dots \supset \mathfrak{H}_m \supset \dots$. From (5.1) it follows the equality

$$\bigcap_{m \geq 0} \mathfrak{H}_m = \{0\}.$$

Let $\Gamma_{-1} = 0 : \mathfrak{M} \rightarrow \mathfrak{N}$. Then $\mathfrak{D}_{\Gamma_{-1}} = \mathfrak{M}$, $\mathfrak{D}_{\Gamma_{-1}^*} = \mathfrak{N}$. We can consider \mathcal{U}_0 as acting from $\mathfrak{D}_{\Gamma_{-1}} \oplus \mathfrak{H}_0$ onto $\mathfrak{D}_{\Gamma_{-1}^*} \oplus \mathfrak{H}_0$ and $\tilde{\mathcal{U}}_0$ as acting from $\mathfrak{D}_{\Gamma_{-1}} \oplus \tilde{\mathfrak{H}}_0$ onto $\mathfrak{D}_{\Gamma_{-1}^*} \oplus \tilde{\mathfrak{H}}_0$. Fix $m \in \mathbb{N}$ and define

$$\Gamma_n^{(m)} = \Gamma_{n+m}, \quad n = -1, 0, 1, \dots$$

Then $\{\Gamma_n^{(m)}\}_{n \geq 0} = \{\Gamma_k\}_{k \geq m}$, and

$$\begin{aligned} \mathfrak{H}_{2k-1} &= \tilde{\mathfrak{H}}_0(\{\Gamma_n^{(2k-1)}\}_{n \geq 0}) = \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 2k-1}), \\ \mathfrak{H}_{2k} &= \mathfrak{H}_0(\{\Gamma_n^{(2k)}\}_{n \geq 0}) = \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 2k}). \end{aligned}$$

Let

$$\begin{aligned} \mathcal{W}_{2k-1} &= \tilde{\mathcal{U}}_0(\{\Gamma_n^{(2k-1)}\}_{n \geq 0}) = \tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 2k-1}) : \begin{array}{ccc} \mathfrak{D}_{\Gamma_{2k-2}} & & \mathfrak{D}_{\Gamma_{2k-2}^*} \\ \oplus & \rightarrow & \oplus \\ \mathfrak{H}_{2k-1} & & \mathfrak{H}_{2k-1} \end{array} , \\ \mathcal{W}_{2k} &= \mathcal{U}_0(\{\Gamma_n^{(2k)}\}_{n \geq 0}) = \mathcal{U}_0(\{\Gamma_n\}_{n \geq 2k}) : \begin{array}{ccc} \mathfrak{D}_{\Gamma_{2k-1}} & & \mathfrak{D}_{\Gamma_{2k-1}^*} \\ \oplus & \rightarrow & \oplus \\ \mathfrak{H}_{2k} & & \mathfrak{H}_{2k} \end{array} , \quad k \in \mathbb{N}. \end{aligned}$$

Define the operators

$$(5.21) \quad \mathcal{T}_m = P_{\mathfrak{H}_m} \mathcal{W}_m \upharpoonright \mathfrak{H}_m, \quad m \in \mathbb{N}.$$

Then

$$(5.22) \quad \begin{aligned} \mathcal{T}_{2k-1} &= \tilde{\mathcal{T}}_0(\{\Gamma_n^{(2k-1)}\}_{n \geq 0}) = \tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 2k-1}), \\ \mathcal{T}_{2k} &= \mathcal{T}_0(\{\Gamma_n^{(2k)}\}_{n \geq 0}) = \mathcal{T}_0(\{\Gamma_n\}_{n \geq 2k}). \end{aligned}$$

From (5.17), (5.18), (5.21), and (5.22) we get

$$\ker D_{\mathcal{T}_0^*} = \mathfrak{H}_1, \quad \ker D_{\mathcal{T}_1} = \mathfrak{H}_2, \quad \dots, \quad \ker D_{\mathcal{T}_{2k}^*} = \mathfrak{H}_{2k+1}, \quad \ker D_{\mathcal{T}_{2k-1}} = \mathfrak{H}_{2k}, \quad \dots$$

From (5.12), (5.13), and (5.22) it follows that

$$\begin{aligned} P_{\ker D_{\mathcal{T}_0^*}} \mathcal{T}_0 \upharpoonright \ker D_{\mathcal{T}_0^*} &= \mathcal{T}_1, \quad P_{\ker D_{\mathcal{T}_1}} \mathcal{T}_1 \upharpoonright \ker D_{\mathcal{T}_1} = \mathcal{T}_2, \quad \dots, \\ P_{\ker D_{\mathcal{T}_{2k-1}}} \mathcal{T}_{2k-1} \upharpoonright \ker D_{\mathcal{T}_{2k-1}} &= \mathcal{T}_{2k}, \quad P_{\ker D_{\mathcal{T}_{2k}^*}} \mathcal{T}_{2k} \upharpoonright \ker D_{\mathcal{T}_{2k}^*} = \mathcal{T}_{2k+1}, \quad \dots \end{aligned}$$

Thus,

$$\mathfrak{H}_{2k-1} = \ker D_{\mathcal{T}_0^{*k}} \cap \ker D_{\mathcal{T}_0^{k-1}}, \quad \mathfrak{H}_{2k} = \ker D_{\mathcal{T}_0^{*k}} \cap \ker D_{\mathcal{T}_0^k}.$$

In notations of Section 4 the operators \mathcal{T}_{2k-1} and \mathcal{T}_{2k} coincide with the operators $(\mathcal{T}_0)_{k-1,k}$ and $(\mathcal{T}_0)_{k,k}$, respectively. From the definition of \mathfrak{H}_0 we get

$$\left(\bigcap_{k \geq 1} \ker D_{\mathcal{T}_0^{*k}} \right) \cap \left(\bigcap_{k \geq 1} \ker D_{\mathcal{T}_0^k} \right) = \bigcap_{k \geq 1} \left(\ker D_{\mathcal{T}_0^{*k}} \cap \ker D_{\mathcal{T}_0^k} \right) = \bigcap_{k \geq 1} \mathfrak{H}_{2k} = \{0\}.$$

So, the operators \mathcal{T}_0 , $\tilde{\mathcal{T}}_0$, and $\{\mathcal{T}_k\}_{k \geq 1}$ are completely non-unitary. It follows that the conservative systems

$$\zeta_0 = \left\{ \begin{bmatrix} \Gamma_0 & \mathcal{G}_0 \\ \mathcal{F}_0 & \mathcal{T}_0 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\} \quad \text{and} \quad \tilde{\zeta}_0 = \left\{ \begin{bmatrix} \Gamma_0 & \tilde{\mathcal{G}}_0 \\ \tilde{\mathcal{F}}_0 & \tilde{\mathcal{T}}_0 \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \tilde{\mathfrak{H}}_0 \right\}$$

are simple.

The operators \mathcal{W}_m take the following 2×2 block operator matrix form

$$\mathcal{W}_m = \begin{bmatrix} \Gamma_m & \mathcal{G}_m \\ \mathcal{F}_m & \mathcal{T}_m \end{bmatrix} : \begin{array}{c} \mathfrak{D}_{\Gamma_{m-1}} \\ \oplus \\ \mathfrak{H}_m \end{array} \rightarrow \begin{array}{c} \mathfrak{D}_{\Gamma_{m-1}^*} \\ \oplus \\ \mathfrak{H}_m \end{array},$$

where

$$\begin{aligned} \mathcal{G}_{2k-1} &= \tilde{\mathcal{G}}_0(\{\Gamma_n\}_{n \geq 2k-1}) = [D_{\Gamma_{2k-1}^*} \quad 0 \quad 0 \quad \dots] : \mathfrak{H}_{2k-1} \rightarrow \mathfrak{D}_{\Gamma_{2k-2}^*}, \\ \mathcal{G}_{2k} &= \mathcal{G}_0(\{\Gamma_n\}_{n \geq 2k}) = [D_{\Gamma_{2k}^*} \Gamma_{2k+1} \quad D_{\Gamma_{2k}^*} D_{\Gamma_{2k+1}^*} \quad 0 \quad 0 \quad \dots] : \mathfrak{H}_{2k} \rightarrow \mathfrak{D}_{\Gamma_{2k-1}^*}, \\ \mathcal{F}_{2k-1} &= \tilde{\mathcal{F}}_0(\{\Gamma_n\}_{n \geq 2k-1}) = \begin{bmatrix} \Gamma_{2k} D_{\Gamma_{2k-1}} \\ D_{\Gamma_{2k}} D_{\Gamma_{2k-1}} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{D}_{\Gamma_{2k-2}} \rightarrow \mathfrak{H}_{2k-1}, \\ \mathcal{F}_{2k} &= \mathcal{F}_0(\{\Gamma_n\}_{n \geq 2k}) = \begin{bmatrix} D_{\Gamma_{2k}} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{D}_{\Gamma_{2k-1}} \rightarrow \mathfrak{H}_{2k}. \end{aligned}$$

Suppose that the system ζ_0 has transfer function Ψ , i.e.,

$$\Psi(\lambda) = \Gamma_0 + \lambda \mathcal{G}_0 (I_{\mathfrak{H}_0} - \lambda \mathcal{T}_0)^{-1} \mathcal{F}_0, \quad \lambda \in \mathbb{D}.$$

Then $\Psi(0) = \Gamma_0$. Let Ψ_1 be the first Schur iterate of Ψ . By (4.5) the transfer function of the simple conservative system

$$\Omega_{0,1}(\nu) = \left\{ \begin{bmatrix} D_{\Gamma_0^*}^{-1} C (D_{\Gamma_0}^{-1} B^*)^* & D_{\Gamma_0^*}^{-1} C \upharpoonright \ker D_{A^*} \\ A P_{\ker D_A} D_{A^*}^{-1} B & P_{\ker D_{A^*}} A \upharpoonright \ker D_{A^*} \end{bmatrix}; \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \ker D_{A^*} \right\}$$

is the first Schur iterate of the transfer function of the simple conservative system

$$\nu = \left\{ \begin{bmatrix} \Gamma_0 & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{N}, \mathfrak{H} \right\}.$$

We will construct the system $\zeta_1 = \Omega_{0,1}(\zeta_0)$ from the system ζ_0 . In our case

$$\zeta_1 = \Omega_{0,1}(\zeta_0) = \left\{ \begin{bmatrix} D_{\Gamma_0^*}^{-1} \mathcal{G}_0 (D_{\Gamma_0}^{-1} \mathcal{F}_0^*)^* & D_{\Gamma_0^*}^{-1} \mathcal{G}_0 \upharpoonright \ker D_{\mathcal{T}_0^*} \\ \mathcal{T}_0 P_{\ker D_{\mathcal{T}_0}} D_{\mathcal{T}_0^*}^{-1} \mathcal{F}_0 & P_{\ker D_{\mathcal{T}_0^*}} \mathcal{T}_0 \upharpoonright \ker D_{\mathcal{T}_0^*} \end{bmatrix}; \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \ker D_{\mathcal{T}_0^*} \right\}.$$

Clearly,

$$D_{\Gamma_0^*}^{-1} \mathcal{G}_0 = [\Gamma_1 \quad D_{\Gamma_1^*} \quad 0 \quad 0 \quad \dots] : \mathfrak{H}_0 \rightarrow \mathfrak{D}_{\Gamma_0^*}, \quad (D_{\Gamma_0}^{-1} \mathcal{F}_0^*)^* = \begin{bmatrix} I_{\mathfrak{M}} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{D}_{\Gamma_0} \rightarrow \mathfrak{H}_0.$$

Therefore, $D_{\Gamma_0^*}^{-1} \mathcal{G}_0 (D_{\Gamma_0}^{-1} \mathcal{F}_0^*)^* = \Gamma_1$. Thus, the first Schur parameter of Ψ is equal to Γ_1 . From (5.17) it follows that $\ker D_{\mathcal{T}_0^*} = \mathfrak{H}_1$ and $\mathfrak{D}_{\mathcal{T}_0^*} = D_{\Gamma_0} P_{\mathfrak{D}_{\Gamma_0}}$. Hence,

$$\mathfrak{D}_{\mathcal{T}_0^*}^{-1} \mathcal{F}_0 = \begin{bmatrix} I_{\mathfrak{D}_{\Gamma_0}} \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathfrak{D}_{\Gamma_0} \rightarrow \mathfrak{H}_0.$$

As has been proved above $P_{\ker D_{\mathcal{T}_0^*}} \mathcal{T}_0 \upharpoonright \ker D_{\mathcal{T}_0^*} = \mathcal{T}_1$. Let $h \in \mathfrak{D}_{\Gamma_0}$. Let us find the projection $P_{\ker D_{\mathcal{T}_0}} h$. According to (5.17) we have to find the vectors $\varphi \in \mathfrak{D}_{\Gamma_1}$ and

$\psi \in \mathfrak{D}_{\Gamma_0^*}$ such that

$$\begin{bmatrix} h \\ 0 \end{bmatrix} = \begin{bmatrix} D_{\Gamma_1} \varphi \\ -\Gamma_1 \varphi \end{bmatrix} + \begin{bmatrix} \Gamma_1^* \psi \\ D_{\Gamma_1^*} \psi \end{bmatrix}.$$

We have $h = D_{\Gamma_1} \varphi + \Gamma_1^* \psi$ and $\Gamma_1 \varphi = D_{\Gamma_1^*} \psi$. From the second equation and Proposition 2.1 it follows $\varphi = D_{\Gamma_1} g$, $\psi = \Gamma_1 g$, where $g \in \mathfrak{D}_{\Gamma_0}$. Therefore, $h = D_{\Gamma_1}^2 g + \Gamma_1^* \Gamma_1 g$, i.e., $g = h$. Hence,

$$P_{\ker D_{\mathcal{T}_0}} \mathfrak{D}_{\mathcal{T}_0^*}^{-1} \mathcal{F}_0 h = P_{\ker D_{\mathcal{T}_0}} h = \begin{bmatrix} D_{\Gamma_1}^2 h \\ -\Gamma_1 D_{\Gamma_1} h \\ 0 \\ 0 \\ \vdots \end{bmatrix}, \quad \mathcal{T}_0 P_{\ker D_{\mathcal{T}_0}} \mathfrak{D}_{\mathcal{T}_0^*}^{-1} \mathcal{F}_0 h = \begin{bmatrix} 0 \\ \Gamma_2 D_{\Gamma_1} h \\ D_{\Gamma_2} D_{\Gamma_1} h \\ 0 \\ 0 \\ \vdots \end{bmatrix} \in \mathfrak{H}_1.$$

Thus,

$$\begin{aligned} \zeta_1 &= \Omega_{0,1}(\zeta_0) \\ &= \left\{ \begin{bmatrix} \Gamma_1 & \mathcal{G}_1 \\ \mathcal{F}_1 & \mathcal{T}_1 \end{bmatrix}; \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \mathfrak{H}_1 \right\} = \left\{ \tilde{U}_0(\{\Gamma_n\}_{n \geq 1}), \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 1}) \right\}. \end{aligned}$$

Similarly

$$\Omega_{1,0}(\tilde{\zeta}_0) = \{U_0(\{\Gamma_n\}_{n \geq 1}), \mathfrak{D}_{\Gamma_0}, \mathfrak{D}_{\Gamma_0^*}, \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 1})\}.$$

The transfer functions of these systems are equal to Ψ_1 (see Section 4), and Γ_1 is exactly is the first Schur parameter of $\Psi(\lambda)$. Let Ψ_2 be the second Schur iterate of Ψ . Constructing the simple conservative system $\zeta_2 = \Omega_{1,0}(\zeta_1)$ of the form (4.5) with the transfer function Ψ_2 we will get the system

$$\zeta_2 = \left\{ \begin{bmatrix} \Gamma_2 & \mathcal{G}_2 \\ \mathcal{F}_2 & \mathcal{T}_2 \end{bmatrix}; \mathfrak{D}_{\Gamma_1}, \mathfrak{D}_{\Gamma_1^*}, \mathfrak{H}_2 \right\} = \left\{ \mathcal{U}_0(\{\Gamma_n\}_{n \geq 2}); \mathfrak{D}_{\Gamma_1}, \mathfrak{D}_{\Gamma_1^*}, \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 2}) \right\}.$$

Let $\Psi_m(\lambda)$ be the m -th Schur iterate of Ψ . Arguing by induction we get that $\Psi_m(\lambda)$ is transfer function of the system

$$\begin{aligned} \zeta_m &= \left\{ \begin{bmatrix} \Gamma_m & \mathcal{G}_m \\ \mathcal{F}_m & \mathcal{T}_m \end{bmatrix}; \mathfrak{D}_{\Gamma_{m-1}}, \mathfrak{D}_{\Gamma_{m-1}^*}, \mathfrak{H}_m \right\} \\ &= \begin{cases} \left\{ \tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 2k-1}); \mathfrak{D}_{\Gamma_{2k-2}}, \mathfrak{D}_{\Gamma_{2k-2}^*}, \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 2k-1}) \right\}, & m = 2k-1, \\ \left\{ \mathcal{U}_0(\{\Gamma_n\}_{n \geq 2k}); \mathfrak{D}_{\Gamma_{2k-1}}, \mathfrak{D}_{\Gamma_{2k-1}^*}, \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 2k}) \right\}, & m = 2k, \end{cases} \end{aligned}$$

for all m . Observe that

$$\zeta_{2k-1} = \Omega_{0,1}(\zeta_{2k-2}), \quad \zeta_{2k} = \Omega_{1,0}(\zeta_{2k-1}), \quad k \in \mathbb{N}.$$

Thus, $\{\Gamma_n\}_{n \geq 0}$ are the Schur parameters of Ψ . \square

From Theorem 5.3 and Theorem 2.2 we immediately arrive at the following result.

Theorem 5.4. *Let $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ and let $\{\Gamma_n\}_{n \geq 0}$ be the Schur parameters of Θ . Then the systems (5.19) are simple conservative realizations of Θ .*

Observe that in fact we have proved Theorem 1.2 differently than in [34] and [19].

Remark 5.5. A more complicated construction of the state Hilbert space and a simple conservative realization for a Schur function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ by means of a block operator matrix are given in [53] (see [19]). These constructions also involve Schur parameters of Θ and some additional Hilbert spaces and operators. One more model based on the Schur parameters of a scalar Schur class function Θ is obtained in [41]. In terms of this model there we given in [41] necessary and sufficient conditions in order for Θ to have a meromorphic pseudocontinuation of bounded type to the exterior of the unit disk. In

- co-shifts of the form

$$\mathcal{T}_m = \tilde{\mathcal{T}}_m = \begin{bmatrix} 0 & I_{\mathfrak{D}_{\Gamma_m^*}} & 0 & 0 & \dots \\ 0 & 0 & I_{\mathfrak{D}_{\Gamma_m^*}} & 0 & \dots \\ 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_m^*}} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} : \begin{matrix} \mathfrak{D}_{\Gamma_m^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_m^*} \\ \oplus \\ \vdots \end{matrix} \rightarrow \begin{matrix} \mathfrak{D}_{\Gamma_m^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_m^*} \\ \oplus \\ \vdots \end{matrix},$$

when Γ_m is isometry,

- the unilateral shifts of the form

$$\mathcal{T}_m = \tilde{\mathcal{T}}_m = \begin{bmatrix} 0 & 0 & 0 & 0 & \dots \\ I_{\mathfrak{D}_{\Gamma_m}} & 0 & 0 & 0 & \dots \\ 0 & I_{\mathfrak{D}_{\Gamma_m}} & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} : \begin{matrix} \mathfrak{D}_{\Gamma_m} \\ \oplus \\ \mathfrak{D}_{\Gamma_m} \\ \oplus \\ \vdots \end{matrix} \rightarrow \begin{matrix} \mathfrak{D}_{\Gamma_m} \\ \oplus \\ \mathfrak{D}_{\Gamma_m} \\ \oplus \\ \vdots \end{matrix},$$

when Γ_m is co-isometry.

One can see that Proposition 5.2 remains true.

Similarly to (5.19) let us consider the conservative systems

$$\zeta_0 = \{\mathcal{U}_0; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_0\}, \quad \tilde{\zeta}_0 = \{\tilde{\mathcal{U}}_0; \mathfrak{M}, \mathfrak{N}, \tilde{\mathfrak{H}}_0\}.$$

One can check that the systems ζ_0 and $\tilde{\zeta}_0$ are simple and unitarily equivalent. Moreover, relations (5.20) and, therefore, Theorem 5.3 and Theorem 5.4 remain valid for the situations considered here.

In order to obtain precise forms of \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ one can consider the following cases:

- (1) Γ_{2N} is isometric (co-isometric) for some N ,
- (2) Γ_{2N+1} is isometric (co-isometric) for some N ,
- (3) the operator Γ_{2N} is unitary for some N ,
- (4) the operator Γ_{2N+1} is unitary for some N .

We shall give several examples.

Example 6.1. The operator Γ_4 is isometric. Define the state spaces

$$\begin{aligned} \mathfrak{H}_0 &:= \begin{matrix} \mathfrak{D}_{\Gamma_0} \\ \oplus \\ \mathfrak{D}_{\Gamma_1^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_2} \\ \oplus \\ \mathfrak{D}_{\Gamma_3^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_4} \\ \oplus \\ \mathfrak{D}_{\Gamma_4^*} \\ \oplus \\ \dots \end{matrix}, \\ \tilde{\mathfrak{H}}_0 &:= \begin{matrix} \mathfrak{D}_{\Gamma_0^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_1} \\ \oplus \\ \mathfrak{D}_{\Gamma_2^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_3} \\ \oplus \\ \mathfrak{D}_{\Gamma_4^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_4} \\ \oplus \\ \dots \end{matrix}. \end{aligned}$$

Then the spaces $\mathfrak{M} \oplus \tilde{\mathfrak{H}}_0$ and $\mathfrak{N} \oplus \mathfrak{H}_0$ can be represented as follows:

$$\begin{aligned} \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0 &= \begin{matrix} \mathfrak{M} \\ \oplus \\ \mathfrak{D}_{\Gamma_0^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_1} \\ \oplus \\ \mathfrak{D}_{\Gamma_2} \\ \oplus \\ \mathfrak{D}_{\Gamma_3} \\ \oplus \\ \mathfrak{D}_{\Gamma_4} \\ \oplus \\ \dots \end{matrix}, \\ \mathfrak{N} \oplus \mathfrak{H}_0 &= \begin{matrix} \mathfrak{N} \\ \oplus \\ \mathfrak{D}_{\Gamma_0} \\ \oplus \\ \mathfrak{D}_{\Gamma_1^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_2} \\ \oplus \\ \mathfrak{D}_{\Gamma_3^*} \\ \oplus \\ \mathfrak{D}_{\Gamma_4} \\ \oplus \\ \dots \end{matrix}. \end{aligned}$$

Define the unitary operators

$$\begin{aligned} \mathcal{M}_0 &= I_{\mathfrak{M}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus I_{\mathfrak{D}_{\Gamma_4^*}} \oplus I_{\mathfrak{D}_{\Gamma_4}} \oplus \dots : \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0 \rightarrow \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0, \\ \tilde{\mathcal{M}}_0 &= I_{\mathfrak{N}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus I_{\mathfrak{D}_{\Gamma_4^*}} \oplus I_{\mathfrak{D}_{\Gamma_4}} \oplus \dots : \mathfrak{N} \oplus \mathfrak{H}_0 \rightarrow \mathfrak{N} \oplus \tilde{\mathfrak{H}}_0, \\ \mathcal{L}_0 &= \mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2} \oplus \mathbf{J}_{\Gamma_4}^{(r)} \oplus I_{\mathfrak{D}_{\Gamma_4^*}} \oplus I_{\mathfrak{D}_{\Gamma_4}} \oplus \dots : \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0 \rightarrow \mathfrak{N} \oplus \mathfrak{H}_0. \end{aligned}$$

Then

$$\mathcal{U}_0 = \mathcal{L}_0 \mathcal{M}_0$$

$$= \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 & -\Gamma_0^* D_{\Gamma_1^*} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & \Gamma_2 D_{\Gamma_1} & -\Gamma_2 \Gamma_1^* & D_{\Gamma_2^*} \Gamma_3 & D_{\Gamma_2^*} D_{\Gamma_3^*} & 0 & 0 & 0 & 0 & \dots \\ 0 & D_{\Gamma_2} D_{\Gamma_1} & -D_{\Gamma_2} \Gamma_1^* & -\Gamma_2^* \Gamma_3 & -\Gamma_2^* D_{\Gamma_3^*} & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & \Gamma_4 D_{\Gamma_3} & -\Gamma_4 \Gamma_3^* & I_{\mathfrak{D}_{\Gamma_4^*}} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_4^*}} & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_4^*}} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix},$$

$$\tilde{\mathcal{U}}_0 = \tilde{\mathcal{M}}_0 \mathcal{L}_0$$

$$= \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & D_{\Gamma_1^*} D_{\Gamma_2^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & -\Gamma_1^* D_{\Gamma_2^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & \Gamma_3 D_{\Gamma_2} & -\Gamma_3 \Gamma_2^* & D_{\Gamma_3^*} \Gamma_4 & D_{\Gamma_3^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & D_{\Gamma_3} D_{\Gamma_2} & -D_{\Gamma_3} \Gamma_2^* & -\Gamma_3^* \Gamma_4 & -\Gamma_3^* & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_4^*}} & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_4^*}} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

Example 6.2. The operator Γ_2 is co-isometric. In this case,

$$\begin{aligned} \mathfrak{H}_0 &= \bigoplus_{\mathfrak{D}_{\Gamma_0}} \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \dots \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \dots, \\ \tilde{\mathfrak{H}}_0 &= \bigoplus_{\mathfrak{D}_{\Gamma_1}} \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \dots \bigoplus \mathfrak{D}_{\Gamma_2} \bigoplus \dots, \end{aligned}$$

$$\mathcal{U}_0 = \mathcal{L}_0 \mathcal{M}_0 = \left(\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2}^{(c)} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus \dots \right) \left(I_{\mathfrak{H}} \oplus \mathbf{J}_{\Gamma_1} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus \dots \right)$$

$$= \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} & 0 & 0 & 0 & \dots \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 & -\Gamma_0^* D_{\Gamma_1^*} & 0 & 0 & 0 & \dots \\ 0 & \Gamma_2 D_{\Gamma_1} & -\Gamma_2 \Gamma_1^* & 0 & 0 & 0 & \dots \\ 0 & D_{\Gamma_2} D_{\Gamma_1} & -D_{\Gamma_2} \Gamma_1^* & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_2}} & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_2}} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix},$$

$$\tilde{\mathcal{U}}_0 = \tilde{\mathcal{M}}_0 \mathcal{L}_0 = \left(I_{\mathfrak{H}} \oplus \mathbf{J}_{\Gamma_1} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus \dots \right) \left(\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2}^{(c)} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus I_{\mathfrak{D}_{\Gamma_2}} \oplus \dots \right)$$

$$= \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 & 0 & 0 & \dots \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & 0 & 0 & 0 & \dots \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & 0 & 0 & 0 & \dots \\ 0 & 0 & D_{\Gamma_2} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_2}} & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_2}} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

$$\tilde{\mathcal{U}}_0 = \tilde{\mathcal{M}}_0 \mathcal{L}_0$$

$$= \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & D_{\Gamma_1^*} D_{\Gamma_2^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & -\Gamma_1^* D_{\Gamma_2^*} & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & \Gamma_3 D_{\Gamma_2} & -\Gamma_3 \Gamma_2^* & D_{\Gamma_3^*} \Gamma_4 & D_{\Gamma_3^*} D_{\Gamma_4^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & D_{\Gamma_3} D_{\Gamma_2} & -D_{\Gamma_3} \Gamma_2^* & -\Gamma_3^* \Gamma_4 & -\Gamma_3^* D_{\Gamma_4^*} & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \Gamma_5 D_{\Gamma_4} & -\Gamma_5 \Gamma_4^* & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & D_{\Gamma_5} D_{\Gamma_4} & -D_{\Gamma_5} \Gamma_4^* & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_5}} & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{\Gamma_5}} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

Example 6.5. The operator Γ_{2N} is unitary. In this case,

$$\mathfrak{H}_0 = \sum_{n=0}^{N-1} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n}} \\ \mathfrak{D}_{\Gamma_{2n+1}^*}}} \oplus, \quad \tilde{\mathfrak{H}}_0 = \sum_{n=0}^{N-1} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n}^*} \\ \mathfrak{D}_{\Gamma_{2n+1}}} \oplus,$$

$$\mathcal{U}_0 = (\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2(N-1)}} \oplus \Gamma_{2N}) (I_{\mathfrak{M}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N-1}}),$$

$$\tilde{\mathcal{U}}_0 = (I_{\mathfrak{M}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N-1}}) (\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2(N-1)}} \oplus \Gamma_{2N}).$$

If $N = 1$ (Γ_2 is unitary) then we have

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 & -\Gamma_0^* D_{\Gamma_1^*} \\ 0 & \Gamma_2 D_{\Gamma_1} & -\Gamma_2 \Gamma_1^* \end{bmatrix}, \quad \tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 \end{bmatrix}.$$

Example 6.6. The operator Γ_{2N+1} is unitary.

$$\mathfrak{H}_0 = \mathfrak{D}_{\Gamma_0}, \quad \tilde{\mathfrak{H}}_0 = \mathfrak{D}_{\Gamma_0^*} \text{ if } N = 0,$$

$$\mathfrak{H}_0 = \sum_{n=0}^{N-1} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n}} \\ \mathfrak{D}_{\Gamma_{2n+1}^*}}} \oplus \mathfrak{D}_{\Gamma_{2N}}, \quad \tilde{\mathfrak{H}}_0 = \sum_{n=0}^{N-1} \bigoplus_{\substack{\mathfrak{D}_{\Gamma_{2n}^*} \\ \mathfrak{D}_{\Gamma_{2n+1}}} \oplus \mathfrak{D}_{\Gamma_{2N}^*} \text{ if } N \geq 1,$$

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 \end{bmatrix} \begin{bmatrix} I_{\mathfrak{M}} & 0 \\ 0 & \Gamma_1 \end{bmatrix} = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 \end{bmatrix},$$

$$\tilde{\mathcal{U}}_0 = \begin{bmatrix} I_{\mathfrak{M}} & 0 \\ 0 & \Gamma_1 \end{bmatrix} \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \\ D_{\Gamma_0} & -\Gamma_0^* \end{bmatrix} = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* \end{bmatrix} \text{ if } N = 0,$$

$$\mathcal{U}_0 = (\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N}}) (I_{\mathfrak{M}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N-1}} \oplus \Gamma_{2N+1}),$$

$$\tilde{\mathcal{U}}_0 = (I_{\mathfrak{M}} \oplus \mathbf{J}_{\Gamma_1} \oplus \mathbf{J}_{\Gamma_3} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N-1}} \oplus \Gamma_{2N+1}) (\mathbf{J}_{\Gamma_0} \oplus \mathbf{J}_{\Gamma_2} \oplus \dots \oplus \mathbf{J}_{\Gamma_{2N}}) \text{ if } N \geq 1.$$

If $N = 1$ (Γ_3 is unitary), then

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} \Gamma_1 & D_{\Gamma_0^*} D_{\Gamma_1^*} & 0 \\ D_{\Gamma_0} & -\Gamma_0^* \Gamma_1 & -\Gamma_0^* D_{\Gamma_1^*} & 0 \\ 0 & \Gamma_2 D_{\Gamma_1} & -\Gamma_2 \Gamma_1^* & D_{\Gamma_2^*} \Gamma_3 \\ 0 & D_{\Gamma_2} D_{\Gamma_1} & -D_{\Gamma_2} \Gamma_1^* & -\Gamma_2^* \Gamma_3 \end{bmatrix},$$

$$\tilde{\mathcal{U}}_0 = \begin{bmatrix} \Gamma_0 & D_{\Gamma_0^*} & 0 & 0 \\ \Gamma_1 D_{\Gamma_0} & -\Gamma_1 \Gamma_0^* & D_{\Gamma_1^*} \Gamma_2 & D_{\Gamma_1^*} D_{\Gamma_2^*} \\ D_{\Gamma_1} D_{\Gamma_0} & -D_{\Gamma_1} \Gamma_0^* & -\Gamma_1^* \Gamma_2 & -\Gamma_1^* D_{\Gamma_2^*} \\ 0 & 0 & \Gamma_3 D_{\Gamma_2} & -\Gamma_3 \Gamma_2^* \end{bmatrix}.$$

7. UNITARY OPERATORS WITH CYCLIC SUBSPACES, DILATIONS, AND BLOCK OPERATOR CMV MATRICES

7.1. Carathéodory class functions associated with conservative systems.

Definition 7.1. Let \mathfrak{M} be a separable Hilbert space. The class $\mathbf{C}(\mathfrak{M})$ of $\mathbf{L}(\mathfrak{M})$ -valued functions holomorphic on the unit disk \mathbb{D} and having positive real part for all $\lambda \in \mathbb{D}$ is called the Carathéodory class.

Consider a conservative systems $\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{M}, \mathfrak{H} \right\}$ whose input and output spaces coincide. Put

$$\mathcal{H} = \mathfrak{M} \oplus \mathfrak{H}$$

and let the function $F_\tau(z)$ be defined as follows

$$(7.1) \quad F_\tau(\lambda) = P_{\mathfrak{M}}(U_\tau + \lambda I_{\mathcal{H}})(U_\tau - \lambda I_{\mathcal{H}})^{-1} \upharpoonright \mathfrak{M}, \quad \lambda \in \mathbb{D},$$

where

$$U_\tau = \begin{bmatrix} D & C \\ B & A \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array} \rightarrow \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array}$$

is the unitary operator in \mathcal{H} associated with the system τ . The function $F_\tau(z)$ is holomorphic in \mathbb{D} and

$$F_\tau(\lambda) + F_\tau^*(\lambda) = 2(1 - |\lambda|^2)P_{\mathfrak{M}}(U_\tau^* - \bar{\lambda}I_{\mathcal{H}})^{-1}(U_\tau - \lambda I_{\mathcal{H}})^{-1} \upharpoonright \mathfrak{M}.$$

It follows that $F_\tau(\lambda) + F_\tau^*(\lambda) \geq 0$ for all $\lambda \in \mathbb{D}$.

The function $F_\tau(\lambda)$ defined by (7.1) belongs to the Carathéodory class $\mathbf{C}(\mathfrak{M})$ and, in addition, $F_\tau(0) = I_{\mathfrak{M}}$. We also shall consider the function

$$\tilde{F}_\tau(\lambda) := F_\tau^*(\bar{\lambda}) = P_{\mathfrak{M}}(I_{\mathcal{H}} + \lambda U_\tau)(I_{\mathcal{H}} - \lambda U_\tau)^{-1}, \quad \lambda \in \mathbb{D}.$$

The functions F_τ and \tilde{F}_τ will be called the Carathéodory functions associated with the conservative system $\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{M}, \mathfrak{H} \right\}$.

Proposition 7.2. Let

$$\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{M}, \mathfrak{H} \right\}$$

be a conservative system. Then the transfer function Θ_τ and the Carathéodory function F_τ are connected by the following relations

$$(7.2) \quad \begin{aligned} \Theta_\tau^*(\bar{\lambda}) &= \frac{1}{\lambda}(F_\tau(\lambda) - I_{\mathfrak{M}})(F_\tau(\lambda) + I_{\mathfrak{M}})^{-1}, \\ F_\tau(\lambda) &= (I_{\mathfrak{M}} + \lambda\Theta_\tau^*(\bar{\lambda}))(I_{\mathfrak{M}} - \lambda\Theta_\tau^*(\bar{\lambda}))^{-1}, \quad \lambda \in \mathbb{D}. \end{aligned}$$

Proof. We use the well known Schur–Frobenius formula for the inverse of block operators. Let Φ be a bounded linear operator given by the block operator matrix

$$\Phi = \begin{pmatrix} X & Y \\ Z & W \end{pmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array} \rightarrow \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array}.$$

Suppose that $W^{-1} \in \mathbf{L}(\mathfrak{H})$ and $(X - YW^{-1}Z)^{-1} \in \mathbf{L}(\mathfrak{M})$. Then $\Phi^{-1} \in \mathbf{L}(\mathfrak{M} \oplus \mathfrak{H}, \mathfrak{M} \oplus \mathfrak{H})$ and

$$\Phi^{-1} = \begin{pmatrix} K^{-1} & -K^{-1}YW^{-1} \\ -W^{-1}ZK^{-1} & W^{-1} + W^{-1}ZK^{-1}YW^{-1} \end{pmatrix},$$

where $K = X - YW^{-1}Z$. Applying this formula for

$$\Phi = I_{\mathcal{H}} - \lambda U_\tau = \begin{pmatrix} I_{\mathfrak{M}} - \lambda D & -\lambda C \\ -\lambda B & I_{\mathfrak{H}} - \lambda A \end{pmatrix}, \quad \lambda \in \mathbb{D},$$

we get $K = I_{\mathfrak{M}} - \lambda D - \lambda^2 C(I_{\mathfrak{H}} - \lambda A)^{-1} B = I_{\mathfrak{M}} - \lambda \Theta_{\tau}(\lambda)$. Therefore,

$$P_{\mathfrak{M}}(I_{\mathcal{H}} - \lambda U_{\tau})^{-1} \upharpoonright \mathfrak{M} = (I_{\mathfrak{M}} - \lambda \Theta_{\tau}(\lambda))^{-1}, \quad \lambda \in \mathbb{D}.$$

Hence

$$P_{\mathfrak{M}}(I_{\mathcal{H}} - \lambda U_{\tau}^*)^{-1} \upharpoonright \mathfrak{M} = (I_{\mathfrak{M}} - \lambda \Theta_{\tau}^*(\bar{\lambda}))^{-1}, \quad \lambda \in \mathbb{D}.$$

Since U_{τ} is unitary, from (7.1) we get

$$\begin{aligned} F_{\tau}(\lambda) &= P_{\mathfrak{M}}(I_{\mathcal{H}} + \lambda U_{\tau}^*)(I_{\mathcal{H}} - \lambda U_{\tau})^{-1} \upharpoonright \mathfrak{M} \\ &= -I_{\mathfrak{M}} + 2P_{\mathfrak{M}}(I_{\mathcal{H}} - \lambda U_{\tau}^*)^{-1} \upharpoonright \mathfrak{M} = -I_{\mathfrak{M}} + 2(I_{\mathfrak{M}} - \lambda \Theta_{\tau}^*(\bar{\lambda}))^{-1} \\ &= (I_{\mathfrak{M}} + \lambda \Theta_{\tau}^*(\bar{\lambda}))(I_{\mathfrak{M}} - \lambda \Theta_{\tau}^*(\bar{\lambda}))^{-1}, \quad \lambda \in \mathbb{D}. \end{aligned}$$

□

The following theorem is well known (see [28]).

Theorem 7.3. *Let \mathfrak{M} be a separable Hilbert space and let $F \in \mathbf{C}(\mathfrak{M})$. Then*

(1) *F admits the integral representation*

$$F(\lambda) = \frac{1}{2}(F(0) - F^*(0)) + \int_0^{2\pi} \frac{e^{it} + \lambda}{e^{it} - \lambda} d\Sigma(t), \quad \lambda \in \mathbb{D},$$

where $\Sigma(t)$ is a non-decreasing and nonnegative $\mathbf{L}(\mathfrak{M})$ -valued function on $[0, 2\pi]$;

(2) *under the condition $F(0) = I_{\mathfrak{M}}$ there exists a Hilbert space \mathcal{H} containing \mathfrak{M} as a subspace, and a unitary operator U in \mathcal{H} such that*

$$F(\lambda) = P_{\mathfrak{M}}(U + \lambda I_{\mathcal{H}})(U - \lambda I_{\mathcal{H}})^{-1} \upharpoonright \mathfrak{M}, \quad \lambda \in \mathbb{D};$$

moreover, the pair $\{\mathcal{H}, U\}$ can be chosen minimal in the sense

$$\overline{\text{span}} \{U^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathcal{H}.$$

Proposition 7.4. [41]. *The conservative system*

$$\tau = \left\{ \begin{bmatrix} D & C \\ B & A \end{bmatrix}; \mathfrak{M}, \mathfrak{M}, \mathfrak{H} \right\}$$

is simple if and only if $\overline{\text{span}} \{U_{\tau}^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathcal{H}$.

Proof. Let τ be a simple conservative system. Suppose $h \in \mathcal{H}$ and h is orthogonal to $U_{\tau}^n \mathfrak{M}$ for all $n \in \mathbb{Z}$. Then the vectors $U_{\tau}^{*n} h$ are orthogonal to \mathfrak{M} in \mathcal{H} for all $n \in \mathbb{Z}$. It follows that $h \in \mathfrak{H}$ and

$$(7.3) \quad \begin{aligned} Ch &= CAh = CA^2h = \dots = CA^n h = \dots = 0, \\ B^* h &= B^* A^* h = B^* A^{*2} h = \dots = B^* A^{*n} h = \dots = 0. \end{aligned}$$

Hence, $h \in \left(\bigcap_{n \geq 0} \ker(CA^n) \right) \cap \left(\bigcap_{n \geq 0} \ker(B^* A^{*n}) \right)$. Since τ is simple we get $h = 0$, i.e.,

$$\overline{\text{span}} \{U_{\tau}^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathcal{H}.$$

Conversely, let $\overline{\text{span}} \{U_{\tau}^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathcal{H}$. Suppose that relations (7.3) hold for some $h \in \mathfrak{H}$. Then $h \perp U_{\tau}^n \mathfrak{M}$ for all $n \in \mathbb{Z}$. Hence, $h = 0$ and τ is simple. □

7.2. Unitary operators with cyclic subspaces. Let U be a unitary operator in a separable Hilbert space \mathfrak{K} and let \mathfrak{M} be a subspace of \mathfrak{K} . Put $\mathfrak{H} = \mathfrak{K} \ominus \mathfrak{M}$. Then U takes the block operator matrix form

$$U = \begin{bmatrix} D & C \\ B & A \end{bmatrix} : \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array} \rightarrow \begin{array}{c} \mathfrak{M} \\ \oplus \\ \mathfrak{H} \end{array}.$$

Since U is unitary, the system $\eta = \{U; \mathfrak{M}, \mathfrak{M}, \mathfrak{H}\}$ is conservative. By Proposition 7.4 the system η is simple if and only if

$$(7.4) \quad \overline{\text{span}} \{U^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathfrak{K}.$$

A subspace \mathfrak{M} of \mathfrak{K} is called *cyclic* for U if the condition (7.4) is satisfied.

Define the Carathéodory function

$$F_{\mathfrak{M}}(\lambda) = P_{\mathfrak{M}}(U + \lambda I_{\mathcal{H}})(U - \lambda I_{\mathcal{H}})^{-1} \upharpoonright \mathfrak{M}, \quad \lambda \in \mathbb{D},$$

and a Schur function

$$E_{\mathfrak{M}}(\lambda) = \frac{1}{\lambda} (F_{\mathfrak{M}}(\lambda) - I_{\mathfrak{M}})(F_{\mathfrak{M}}(\lambda) + I_{\mathfrak{M}})^{-1}, \quad \lambda \in \mathbb{D}.$$

According to Proposition 7.2 the transfer function $\Theta(\lambda)$ of the system η and the function $E_{\mathfrak{M}}(\lambda)$ are connected by the relation

$$\Theta(\lambda) = E_{\mathfrak{M}}^*(\bar{\lambda}), \quad \lambda \in \mathbb{D}.$$

Theorem 7.5. *Let U be a unitary operator in a separable Hilbert space and let \mathfrak{M} be a cyclic subspace for U . Then U is unitarily equivalent to the block operator CMV matrices $\mathcal{U}_0(\{\Gamma_n\}_{n \geq 0})$ and $\tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0})$ in the Hilbert spaces $\mathcal{H} = \mathfrak{M} \oplus \mathfrak{H}_0(\{\Gamma_n\}_{n \geq 0})$ and $\tilde{\mathcal{H}} = \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0(\{\Gamma_n\}_{n \geq 0})$, respectively, where $\{\Gamma_n\}_{n \geq 0}$ are the Schur parameters of the function*

$$\Theta(\lambda) = \frac{1}{\lambda} (F_{\mathfrak{M}}^*(\bar{\lambda}) - I_{\mathfrak{M}})(F_{\mathfrak{M}}^*(\bar{\lambda}) + I_{\mathfrak{M}})^{-1}, \quad \lambda \in \mathbb{D}.$$

Proof. Because \mathfrak{M} is a cyclic subspace for U , the conservative system η is simple. By Theorem 5.4 the system η is unitarily equivalent to the systems ζ_0 and $\tilde{\zeta}_0$ given by (5.19). From (3.5) it follows that U is unitarily equivalent to $\mathcal{U}_0(\{\Gamma_n\}_{n \geq 0})$ and $\tilde{\mathcal{U}}_0(\{\Gamma_n\}_{n \geq 0})$. \square

Suppose that the cyclic subspace \mathfrak{M} for unitary operator U in \mathfrak{K} is one-dimensional. Let $\varphi \in \mathfrak{M}$, $\|\varphi\| = 1$, and let $\mu(\zeta) = (\mathcal{E}(\zeta)\varphi, \varphi)_{\mathfrak{K}}$, where $\mathcal{E}(\zeta)$, $\zeta \in \mathbb{T}$, is the resolution of the identity for U . Then the scalar Carathéodory function $F(\lambda)$ is of the form

$$F(\lambda) = ((U + \lambda I_{\mathcal{H}})(U - \lambda I_{\mathcal{H}})^{-1} \varphi, \varphi)_{\mathfrak{K}} = \int_{\mathbb{T}} \frac{\zeta + \lambda}{\zeta - \lambda} d\mu(\zeta), \quad \lambda \in \mathbb{D}.$$

Thus, the function F is associated with the probability measure μ on \mathbb{T} . The Schur function associated with μ [66] is the function

$$E(\lambda) = \frac{1}{\lambda} \frac{F(\lambda) - 1}{F(\lambda) + 1}, \quad \lambda \in \mathbb{D}.$$

By Geronimus theorem [46] the Schur parameters of the function E coincide with Verblunsky coefficient $\{\alpha_n\}_{n \geq 0}$ of the measure μ (see [66]). Let $\Theta(\lambda) := \overline{E(\bar{\lambda})}$, $\lambda \in \mathbb{D}$, and let $\{\gamma_n\}_{n \geq 0}$ be the Schur parameters of Θ . Then $\bar{\alpha}_n = \gamma_n$ for all n and the CMV matrices $\mathcal{U}_0 = \mathcal{U}_0(\{\gamma_n\}_{n \geq 0})$ and $\tilde{\mathcal{U}}_0 = \tilde{\mathcal{U}}_0(\{\gamma_n\}_{n \geq 0})$ coincide with the CMV matrices \mathcal{C} and $\tilde{\mathcal{C}}$ given by (1.2) and (1.3), correspondingly. Observe that $\dim \mathfrak{K} = m \iff$ the function E is the Blaschke product of the form

$$E(\lambda) = e^{i\varphi} \prod_{k=1}^m \frac{\lambda - \lambda_k}{1 - \bar{\lambda}_k \lambda}.$$

7.3. Unitary dilations of a contraction. Let T be a contraction acting in a Hilbert space H . A unitary operator U in a Hilbert space \mathcal{H} containing H as a subspace is called a unitary dilation of T if $T^n = P_H U^n \upharpoonright \mathcal{H}$ for all $n \in \mathbb{N}$ [71]. Two unitary dilations U in \mathcal{H} and U' in \mathcal{H}' of T are called isomorphic if there exists a unitary operator $W \in \mathbf{L}(\mathcal{H}, \mathcal{H}')$ such that $W \upharpoonright H = I_H$ and $WU = U'W$. It is established in [71] that for every contraction

T in the Hilbert space H there exists a unitary dilation U in a space H such that U is minimal [71], i.e.,

$$\overline{\text{span}} \{U^n H, n \in \mathbb{Z}\} = \mathcal{H}.$$

Moreover, two minimal unitary dilations of T are isomorphic [71]. The minimal unitary dilation by means of the infinite matrix form is constructed in [71] on the base of Schäffer paper [61]. Below we show that the minimal unitary dilations can be given by the operator CMV matrices.

Theorem 7.6. *Let T be a contraction in a Hilbert space H . Define the Hilbert spaces*

$$(7.5) \quad \mathfrak{H}_0 := \begin{array}{cccc} \mathfrak{D}_T & & \mathfrak{D}_T & \\ \oplus & \oplus & \oplus & \oplus \cdots \\ \mathfrak{D}_{T^*} & & \mathfrak{D}_{T^*} & \end{array}, \quad \tilde{\mathfrak{H}}_0 := \begin{array}{cccc} \mathfrak{D}_{T^*} & & \mathfrak{D}_{T^*} & \\ \oplus & \oplus & \oplus & \oplus \cdots \\ \mathfrak{D}_T & & \mathfrak{D}_T & \end{array},$$

and the Hilbert spaces $\mathcal{H}_0 := H \oplus \mathfrak{H}_0$, and $\tilde{\mathcal{H}}_0 := H \oplus \tilde{\mathfrak{H}}_0$. Let

$$\mathbf{J}_0 = \begin{bmatrix} 0 & I_{\mathfrak{D}_{T^*}} \\ I_{\mathfrak{D}_T} & 0 \end{bmatrix} : \begin{array}{cc} \mathfrak{D}_T & \mathfrak{D}_{T^*} \\ \oplus & \oplus \\ \mathfrak{D}_{T^*} & \mathfrak{D}_T \end{array} \rightarrow \begin{array}{cc} \mathfrak{D}_{T^*} & \mathfrak{D}_T \\ \oplus & \oplus \\ \mathfrak{D}_T & \mathfrak{D}_{T^*} \end{array}.$$

Define operators

$$(7.6) \quad \begin{aligned} \mathcal{M}_0 &= I_H \bigoplus \mathbf{J}_0 \bigoplus \mathbf{J}_0 \bigoplus \cdots : \mathcal{H}_0 \rightarrow \tilde{\mathcal{H}}_0, \\ \mathcal{L}_0 &= \mathbf{J}_T \bigoplus \mathbf{J}_0 \bigoplus \mathbf{J}_0 \bigoplus \cdots : \tilde{\mathcal{H}}_0 \rightarrow \mathcal{H}_0, \end{aligned}$$

and

$$(7.7) \quad \mathcal{U}_0 = \mathcal{L}_0 \mathcal{M}_0 : \mathcal{H}_0 \rightarrow \mathcal{H}_0, \quad \tilde{\mathcal{U}}_0 = \mathcal{M}_0 \mathcal{L}_0 : \tilde{\mathcal{H}}_0 \rightarrow \tilde{\mathcal{H}}_0.$$

Then $\{\mathcal{H}_0, \mathcal{U}_0\}$ and $\{\tilde{\mathcal{H}}_0, \tilde{\mathcal{U}}_0\}$ are unitarily equivalent minimal unitary dilations of the operator T .

Proof. Define the $\mathbf{L}(H)$ -valued function

$$\tilde{F}(\lambda) = (I_H + \lambda T)(I_H - \lambda T)^{-1}, \quad \lambda \in \mathbb{D}.$$

Then the function \tilde{F} belongs to the Carathéodory class $\mathbf{C}(H)$ and

$$\Theta(\lambda) := \frac{1}{\lambda} (\tilde{F}(\lambda) - I_H)(\tilde{F}(\lambda) + I_H)^{-1} = T, \quad \lambda \in \mathbb{D},$$

belongs to the Schur class $\mathbf{S}(H, H)$. The Schur parameters of Θ is the sequence

$$\Gamma_0 = T, \quad \Gamma_n = 0 \in \mathbf{S}(\mathfrak{D}_T, \mathfrak{D}_{T^*}), \quad n \in \mathbb{N}.$$

Let \mathfrak{H}_0 and $\tilde{\mathfrak{H}}_0$ be defined by (7.5), $\mathcal{H}_0 = H \oplus \mathfrak{H}_0$, $\tilde{\mathcal{H}}_0 = H \oplus \tilde{\mathfrak{H}}_0$. Then the operators \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$ defined by (7.7) are the block operator CMV matrices constructed by means of the Schur parameters of Θ . Let $\zeta_0 = \{\mathcal{U}_0, \mathfrak{M}, \mathfrak{M}, \mathfrak{H}_0\}$ and $\tilde{\zeta}_0 = \{\tilde{\mathcal{U}}_0, \mathfrak{M}, \mathfrak{M}, \tilde{\mathfrak{H}}_0\}$ be the corresponding conservative systems. By Theorem 5.4 the systems ζ_0 and $\tilde{\zeta}_0$ are simple, unitary equivalent, and their transfer functions are equal Θ . By Proposition 7.2 we have

$$\begin{aligned} (I_H + \lambda T)(I_H - \lambda T)^{-1} &= \tilde{F}(\lambda) = (I_H + \lambda \Theta(\lambda))(I_H - \lambda \Theta(\lambda))^{-1} \\ &= P_H(I_{\mathcal{H}_0} + \lambda \mathcal{U}_0)(I_{\mathcal{H}_0} - \lambda \mathcal{U}_0)^{-1} \upharpoonright H, \quad \lambda \in \mathbb{D}. \end{aligned}$$

Hence $T^n = P_H \mathcal{U}_0^n \upharpoonright H$, $n \in \mathbb{N}$. Therefore \mathcal{U}_0 is a unitary dilation of T in \mathcal{H}_0 . By Proposition 7.4 this dilation is minimal. Similarly, the operator $\tilde{\mathcal{U}}_0$ is a minimal unitary dilation of T in $\tilde{\mathcal{H}}_0$. \square

Taking into account (7.6), (5.6), and (5.7) we obtain the following operator matrix forms for the minimal unitary dilations \mathcal{U}_0 and $\tilde{\mathcal{U}}_0$:

$$\mathcal{U}_0 = \begin{bmatrix} T & 0 & D_{T^*} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ D_T & 0 & -T^* & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & 0 & 0 & 0 & 0 & \dots \\ 0 & I_{\mathfrak{D}_T} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & 0 & 0 & \dots \\ 0 & 0 & 0 & I_{\mathfrak{D}_T} & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix},$$

$$\tilde{\mathcal{U}}_0 = \begin{bmatrix} T & D_{T^*} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & 0 & 0 & 0 & 0 & 0 & \dots \\ D_T & -T^* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & 0 & 0 & 0 & \dots \\ 0 & 0 & I_{\mathfrak{D}_T} & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{D}_{T^*}} & 0 & \dots \\ 0 & 0 & 0 & 0 & I_{\mathfrak{D}_T} & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

7.4. The Naimark dilation. Let \mathfrak{M} be a separable Hilbert space. Denote by $\mathfrak{B}(\mathbb{T})$ the σ -algebra of Borelian subsets of the unit circle $\mathbb{T} = \{\xi \in \mathbb{C} : |\xi| = 1\}$. Let μ be a $\mathbf{L}(\mathfrak{M})$ -valued Borel measure on $\mathfrak{B}(\mathbb{T})$, i.e.,

- (a) for any $\delta \in \mathfrak{B}(\mathbb{T})$ the operator $\mu(\delta)$ is nonnegative,
- (b) $\mu(\emptyset) = 0$,
- (c) μ is σ -additive with respect to the strong operator convergence.

Denote by $\mathbf{M}(\mathbb{T}, \mathfrak{M})$ the set of all $\mathbf{L}(\mathfrak{M})$ -valued Borel measures.

Definition 7.7. [33], [44], [19], [41]. Let $\mu \in \mathbf{M}(\mathbb{T}, \mathfrak{M})$ be a probability measure ($\mu(\mathbb{T}) = I_{\mathfrak{M}}$) and let the operators $\{S_n\}_{n \in \mathbb{Z}}$ be the sequence of Fourier coefficients of μ , i.e.,

$$S_n = \int_{\mathbb{T}} \xi^{-n} \mu(d\xi), \quad n \in \mathbb{Z}.$$

A Naimark dilation of μ is a pair $\{\mathcal{H}, \mathcal{U}\}$, where \mathcal{H} is a separable Hilbert space containing \mathfrak{M} as a subspace, \mathcal{U} is unitary operator in \mathcal{H} such that

$$S_n = P_{\mathfrak{M}} \mathcal{U}^n \upharpoonright \mathfrak{M}, \quad n \in \mathbb{Z}.$$

A Naimark dilation is called minimal if $\overline{\text{span}} \{\mathcal{U}^n \mathfrak{M}, n \in \mathbb{Z}\} = \mathcal{H}$.

Proposition 7.8. [33], [44], [19], [41]. Let $\{\mathcal{H}_1, \mathcal{U}_1\}$ and $\{\mathcal{H}_2, \mathcal{U}_2\}$ be two minimal Naimark dilations of a probability measure $\mu \in \mathbf{M}(\mathbb{T}, \mathfrak{M})$. Then there exists a unitary operator $\mathcal{W} \in \mathbf{L}(\mathcal{H}_1, \mathcal{H}_2)$ such that $\mathcal{W} \mathcal{U}_1 = \mathcal{U}_2 \mathcal{W}$ and $\mathcal{W} \upharpoonright \mathfrak{M} = I_{\mathfrak{M}}$.

Notice that $S_0 = I_{\mathfrak{M}}$, $S_n = S_n^*$ for all $n \in \mathbb{Z}$ and the sequence $\{S_n\}_{n \in \mathbb{Z}}$ is positive [33], [35], [19], [44], i.e.,

$$\sum_{i,j=0}^{\infty} (S_{i-j} h_i, h_j)_{\mathfrak{M}} \geq 0$$

for all sequences $\{h_i\}_{i \geq 0} \subset \mathfrak{M}$ with finite support. If ν is the spectral measure of a minimal Naimark dilation of μ , then $\mu = P_{\mathfrak{M}} \nu \upharpoonright \mathfrak{M}$. The minimal Naimark dilation was constructed by T. Constantinescu in [33] (see also ([44]) by means of an infinite in both sides block operator matrix whose entries depend on the choice sequence determined by $\{S_n\}_{n \in \mathbb{Z}}$ (see [35]), the Hessenberg matrix representation of a minimal isometric dilation

of $\{S_n\}_{n \geq 0}$ is obtained in [44]. Here we construct the minimal Naimark dilations in the form of block operator CMV matrices.

Theorem 7.9. *Let \mathfrak{M} be a separable Hilbert space and let $\mu \in \mathbf{M}(\mathbb{T}, \mathfrak{M})$ be a probability measure. Define the functions*

$$F(\lambda) = \int_{\mathbb{T}} \frac{\xi + \lambda}{\xi - \lambda} \mu(d\xi), \quad E(\lambda) = \frac{1}{\lambda} (F(\lambda) - I_{\mathfrak{M}})(F(\lambda) + I_{\mathfrak{M}})^{-1}, \quad \lambda \in \mathbb{D}.$$

Then $E \in \mathbf{S}(\mathfrak{M}, \mathfrak{M})$. Let $\{G_n\}_{n \geq 0}$ be the Schur parameters of E . Define the Hilbert spaces $\mathfrak{H}_0 = \mathfrak{H}_0(\{G_n\}_{n \geq 0})$, $\tilde{\mathfrak{H}}_0 = \tilde{\mathfrak{H}}_0(\{G_n\}_{n \geq 0})$ and the Hilbert spaces $\mathcal{H}_0 = \mathfrak{M} \oplus \mathfrak{H}_0$, $\tilde{\mathcal{H}}_0 = \mathfrak{M} \oplus \tilde{\mathfrak{H}}_0$. Let

$$\mathcal{U}_0 = \mathcal{U}_0(\{G_n\}_{n \geq 0}), \quad \tilde{\mathcal{U}}_0 = \tilde{\mathcal{U}}_0(\{G_n\}_{n \geq 0})$$

be the block operator CMV matrices constructed by means of $\{G_n\}$. Then the pairs $\{\mathcal{H}_0, \mathcal{U}_0\}$ and $\{\tilde{\mathcal{H}}_0, \tilde{\mathcal{U}}_0\}$ are unitarily equivalent minimal Naimark dilations of the measure μ .

Proof. Clearly

$$F(\lambda) = I_{\mathfrak{M}} + 2 \sum_{n=1}^{\infty} \lambda^n \int_{\mathbb{T}} \xi^{-n} \mu(d\xi) = I_{\mathfrak{M}} + 2 \sum_{n=1}^{\infty} \lambda^n S_n, \quad \lambda \in \mathbb{D}.$$

Then

$$F^*(\bar{\lambda}) = I_{\mathfrak{M}} + 2 \sum_{n=1}^{\infty} \lambda^n S_{-n}.$$

Because $F(\lambda) + F^*(\lambda) \geq 0$ for $\lambda \in \mathbb{D}$, the $\mathbf{L}(\mathfrak{M})$ -valued function E belongs to the Schur class $\mathbf{S}(\mathfrak{M}, \mathfrak{M})$. Construct the Hilbert spaces $\mathfrak{H}_0 = \mathfrak{H}_0(\{G_n\}_{n \geq 0})$, $\mathcal{H}_0 = \mathfrak{M} \oplus \mathfrak{H}_0$ and let $\mathcal{U}_0 = \mathcal{U}_0(\{G_n\}_{n \geq 0}) = (\mathcal{U}_0(\{G_n\}_{n \geq 0}))^*$ be the block operator CMV matrix. Then \mathcal{U}_0 is a unitary operator in the Hilbert space \mathcal{H}_0 . The system $\zeta_0 = \{\mathcal{U}_0; \mathfrak{M}, \mathfrak{M}, \mathfrak{H}_0\}$ is conservative and simple, and its transfer function is equal to $E(\lambda)$ (see Subsection 5.3, (5.19), Theorem 5.4). Hence the transfer of the adjoint system $\zeta_0^* = \{\mathcal{U}_0^*; \mathfrak{M}, \mathfrak{M}, \mathfrak{H}_0\}$ is equal to $\Theta(\lambda) = E^*(\bar{\lambda})$. By definition of $F(\lambda)$ and $E(\lambda)$, and by Proposition 7.2, and (7.2) we have

$$\begin{aligned} F(\lambda) &= (I_{\mathfrak{M}} + \lambda E(\lambda))(I_{\mathfrak{M}} - \lambda E(\lambda))^{-1} = (I_{\mathfrak{M}} + \lambda \Theta^*(\bar{\lambda}))(I_{\mathfrak{M}} - \lambda \Theta^*(\bar{\lambda}))^{-1} \\ &= P_{\mathfrak{M}}(\mathcal{U}_0^* + \lambda I_{\mathcal{H}_0})(\mathcal{U}_0^* - \lambda I_{\mathcal{H}_0})^{-1} \upharpoonright \mathfrak{M}. \end{aligned}$$

Hence,

$$\begin{aligned} F(\lambda) &= I_{\mathfrak{M}} + 2 \sum_{n=1}^{\infty} \lambda^n P_{\mathfrak{M}} \mathcal{U}_0^n \upharpoonright \mathfrak{M}, \\ F^*(\bar{\lambda}) &= I_{\mathfrak{M}} + 2 \sum_{n=1}^{\infty} \lambda^n P_{\mathfrak{M}} \mathcal{U}_0^{-n} \upharpoonright \mathfrak{M}. \end{aligned}$$

Thus, the pair $\{\mathcal{H}_0, \mathcal{U}_0\}$ is the minimal Naimark dilation of the measure μ . The same is true for the pair $\{\tilde{\mathcal{H}}_0, \tilde{\mathcal{U}}_0\}$. □

8. THE BLOCK OPERATOR CMV MATRIX MODELS FOR COMPLETELY NON-UNITARY CONTRACTIONS

Theorem 8.1. *Let T be a completely non-unitary contraction in a separable Hilbert space H . Let*

$$\Phi_T(\lambda) = (-T + \lambda D_{T^*}(I_H - \lambda T^*)^{-1} D_T) \upharpoonright \mathfrak{D}_T, \quad \lambda \in \mathbb{D},$$

be the Sz.-Nagy–Foias characteristic function of T [71]. If $\{\Gamma_n\}_{n \geq 0}$ are the Schur parameters of Φ_T , then the operator T is unitarily equivalent to the truncated block operator CMV matrices $\mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0})$ and $\tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0})$.

Proof. The transfer function of the simple conservative system

$$\eta = \left\{ \begin{bmatrix} -T^* & D_T \\ D_{T^*} & T \end{bmatrix}; \mathfrak{D}_{T^*}, \mathfrak{D}_T, H \right\}$$

is given by

$$\Theta_\eta(\lambda) = (-T^* + \lambda D_T(I_H - \lambda T)^{-1} D_{T^*}) \upharpoonright \mathfrak{D}_{T^*}, \quad \lambda \in \mathbb{D}.$$

It follows that $\Phi_T(\lambda) = \Theta_\eta^*(\bar{\lambda})$, $\lambda \in \mathbb{D}$. Hence, if $\{\Gamma_n\}_{n \geq 0}$ are the Schur parameters of $\Phi_T(\lambda)$, then $\{\Gamma_n^*\}_{n \geq 0}$ are the Schur parameters of $\Theta_\eta(\lambda)$. Construct the Hilbert spaces $\mathfrak{H}_0 = \mathfrak{H}_0(\{\Gamma_n^*\}_{n \geq 0})$, $\tilde{\mathfrak{H}}_0 = \tilde{\mathfrak{H}}_0(\{\Gamma_n^*\}_{n \geq 0})$, the block operator CMV matrices $\mathcal{U}_0 = \mathcal{U}_0(\{\Gamma_n^*\}_{n \geq 0})$, $\tilde{\mathcal{U}}_0 = \tilde{\mathcal{U}}_0(\{\Gamma_n^*\}_{n \geq 0})$, truncated block CMV matrices $\mathcal{T}_0 = \mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0})$ and $\tilde{\mathcal{T}}_0 = \tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0})$. Consider the corresponding conservative systems

$$\zeta_0 = \{\mathcal{U}_0; \mathfrak{D}_{T^*}, \mathfrak{D}_T, \mathfrak{H}_0\}, \quad \tilde{\zeta}_0 = \{\tilde{\mathcal{U}}_0; \mathfrak{D}_{T^*}, \mathfrak{D}_T, \tilde{\mathfrak{H}}_0\}.$$

By Theorem 5.4 the systems ζ_0 and $\tilde{\zeta}_0$ are simple conservative realizations of the function Θ . It follows that the operator T is unitarily equivalent to the operators $\mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0})$ and $\tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0})$. \square

Observe that $\mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0}) = \left(\tilde{\mathcal{T}}_0(\{\Gamma_n\}_{n \geq 0})\right)^*$ and $\tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0}) = \left(\mathcal{T}_0(\{\Gamma_n\}_{n \geq 0})\right)^*$.

The results of Sz.-Nagy–Foias [71, Theorem VI.3.1] states that if a function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ is purely contractive ($\|\Theta(0)f\| < \|f\|$ for all $f \in \mathfrak{M} \setminus \{0\}$), then there exists a completely non-unitary contraction T whose characteristic function coincides with Θ . Here we give another proof of this result.

Theorem 8.2. *Let a function $\Theta \in \mathbf{S}(\mathfrak{M}, \mathfrak{N})$ be purely contractive. If $\{\Gamma_n\}_{n \geq 0}$ are the Schur parameters of Θ then the characteristic functions of completely non-unitary contractions given by the truncated block operator CMV matrices $\mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0})$ and $\tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0})$ coincide with Θ .*

Proof. Let $\tilde{\Theta}(\lambda) := \Theta^*(\bar{\lambda})$. Then $\{\Gamma_n^*\}_{n \geq 0}$ are the Schur parameters of $\tilde{\Theta}$. Construct the Hilbert spaces $\mathfrak{H}_0 = \mathfrak{H}_0(\{\Gamma_n^*\}_{n \geq 0})$, $\tilde{\mathfrak{H}}_0 = \tilde{\mathfrak{H}}_0(\{\Gamma_n^*\}_{n \geq 0})$, the block operator CMV matrices $\mathcal{U}_0 = \mathcal{U}_0(\{\Gamma_n^*\}_{n \geq 0})$, $\tilde{\mathcal{U}}_0 = \tilde{\mathcal{U}}_0(\{\Gamma_n^*\}_{n \geq 0})$, the truncated block CMV matrices $\mathcal{T}_0 = \mathcal{T}_0(\{\Gamma_n^*\}_{n \geq 0})$, $\tilde{\mathcal{T}}_0 = \tilde{\mathcal{T}}_0(\{\Gamma_n^*\}_{n \geq 0})$, and consider the corresponding conservative systems

$$\zeta_0 = \{\mathcal{U}_0; \mathfrak{M}, \mathfrak{N}, \mathfrak{H}_0\}, \quad \tilde{\zeta}_0 = \{\tilde{\mathcal{U}}_0; \mathfrak{M}, \mathfrak{N}, \tilde{\mathfrak{H}}_0\}.$$

Then the transfer functions of ζ_0 and $\tilde{\zeta}_0$ are equal to $\tilde{\Theta}(\lambda)$. Since the operator

$$\mathcal{U}_0 = \begin{bmatrix} \Gamma_0^* & \mathcal{G}_0 \\ \mathcal{F}_0 & \mathcal{T}_0 \end{bmatrix} : \begin{array}{cc} \mathfrak{M} & \mathfrak{N} \\ \oplus & \rightarrow \oplus \\ \mathfrak{H}_0 & \mathfrak{H}_0 \end{array}$$

is a contraction, there exist contractions (see [18], [38], [65]) $\mathcal{K} \in \mathbf{L}(\mathfrak{D}_{\mathcal{T}_0}, \mathfrak{N})$, $\mathcal{M} \in \mathbf{L}(\mathfrak{M}, \mathfrak{H}_0)$, $\mathcal{X} \in \mathbf{L}(\mathfrak{D}_{\mathcal{M}}, \mathfrak{D}_{\mathcal{K}^*})$ such that

$$\mathcal{G}_0 = \mathcal{K}D_{\mathcal{T}_0}, \quad \mathcal{F}_0 = D_{\mathcal{T}_0^*}\mathcal{M}, \quad \Gamma_0^* = -\mathcal{K}T_0^*\mathcal{M} + D_{\mathcal{K}^*}\mathcal{X}D_{\mathcal{M}}.$$

Because \mathcal{U}_0 is unitary, the operators \mathcal{K} , \mathcal{M}^* are isometries and \mathcal{X} is unitary (see [7],[8]), the characteristic function of \mathcal{T}_0^* and the transfer function of the system ζ_0 are connected by the relation (see [9], [8])

$$\tilde{\Theta}(\lambda) = \mathcal{K}\Phi_{\mathcal{T}_0^*}(\lambda)\mathcal{M} + \mathcal{X}D_{\mathcal{M}}.$$

Since $D_{\mathcal{M}}$ is an orthogonal projection in \mathfrak{M} onto $\ker \mathcal{M}$, and $\tilde{\Theta}(\lambda) \upharpoonright \ker \mathcal{M} = \mathcal{X}$, $\lambda \in \mathbb{D}$, we have, for $f \in \ker \mathcal{M}$,

$$\|\Gamma_0^*f\| = \|\tilde{\Theta}(0)f\| = \|\mathcal{X}f\| = \|f\|.$$

Since Γ_0^* is a pure contraction, we obtain $\ker \mathcal{M} = \{0\}$. Similarly, $\ker \mathcal{K}^* = \{0\}$, i.e., \mathcal{K} and \mathcal{M} are unitary operators, and $\tilde{\Theta}(\lambda) = \mathcal{K}\Phi_{\mathcal{T}_0^*}(\lambda)\mathcal{M}$, $\lambda \in \mathbb{D}$. Thus, the characteristic function $\Phi_{\mathcal{T}_0}$ of \mathcal{T}_0 coincides with Θ . Similarly, the characteristic function $\Phi_{\tilde{\mathcal{T}}_0}$ of $\tilde{\mathcal{T}}_0$ coincides with Θ . \square

Remark 8.3. For completely non-unitary contractions with one-dimensional defect operators and for a scalar Schur class functions, Theorem 8.1 and Theorem 8.2 have been established in [11].

REFERENCES

1. V. M. Adamjan and D. Z. Arov, *On unitary coupling of semi-unitary operators*, Math. Issled., Kishinev I(2) (1966), 3–65. (Russian); English transl. Amer. Math. Soc. Transl., Ser. 2, **95** (1970), 75–129.
2. D. Alpay, T. Azizov, A. Dijksma, and H. Langer, *The Schur algorithm for generalized Schur functions. I. Coisometric realizations*, Oper. Theory Adv. Appl. **129** (2001), 1–36.
3. D. Alpay, T. Azizov, A. Dijksma, H. Langer, and G. Wanjala, *The Schur algorithm for generalized Schur functions. IV. Unitary realizations*, Oper. Theory Adv. Appl. **149** (2004), 23–45.
4. D. Alpay, A. Dijksma, and H. Langer, *The transformation of Issai Schur and related topics in indefinite setting*, Oper. Theory Adv. Appl. **176** (2007), 1–98.
5. D. Alpay, A. Dijksma, J. Rovnyak, and H.S.V. de Snoo, *Schur Functions, Operator Collocations, and Reproducing Kernel Pontryagin Spaces*, Oper. Theory Adv. Appl., Vol. 96, Birkhäuser Verlag, Basel, 1997.
6. T. Ando, *De Branges Spaces and Analytic Operator Functions*, Research Institute of Applied Electricity, Hokkaido University, Sapporo, Japan, 1990.
7. Yu. Arlinskiĭ, *The Kalman-Yakubovich-Popov inequality for passive discrete time-invariant systems*, Operators and Matrices **2** (2008), no. 1, 15–51.
8. Yu. M. Arlinskiĭ, *Conservative realizations of the functions associated with Schur’s algorithm for the Schur class operator-valued function*, Operators and Matrices **3** (2009), no. 1, 59–96.
9. Yu. M. Arlinskiĭ, S. Hassi, and H.S.V. de Snoo, *Parametrization of contractive block-operator matrices and passive discrete-time systems*, Complex Analysis and Operator Theory **1** (2007), no. 2, 211–233.
10. Yu. M. Arlinskiĭ, S. Hassi, and H.S.V. de Snoo, *Q-functions of Hermitian contractions of Kreĭn-Ovcharenko type*, Integral Equations and Operator Theory **53** (2005), no. 2, 153–189.
11. Yu. Arlinskiĭ, L. Golinskiĭ, and E. Tsekanovskiĭ, *Contractions with rank one defect operators and truncated CMV matrices*, J. Funct. Anal. **254** (2008), 154–195.
12. D. Z. Arov, *Passive linear stationary dynamical systems*, Sib. Mat. Zh. **20** (1979), no. 2, 211–228. (Russian); English transl. Siberian Math. J. **20** (1979), 149–162.
13. D. Z. Arov, *Stable dissipative linear stationary dynamical scattering systems*, J. Operator Theory **1** (1979), 95–126. (Russian); English transl. Interpolation Theory, Systems Theory and Related Topics, The Harry Dym Anniversary Volume, Oper. Theory Adv. Appl. **134** (2002), 99–136.
14. D. Z. Arov, M. A. Kaashoek, and D. P. Pik, *Minimal and optimal linear discrete time-invariant dissipative scattering systems*, Integral Equations and Operator Theory **29** (1997), 127–154.
15. D. Z. Arov and M. A. Nudel’man, *Criterion of unitary similarity of minimal passive scattering systems with a given transfer function*, Ukrain. Mat. Zh. **52**, no. 2 (2000), 147–156. (Russian); English transl. Ukrainian Math. J. **52** (2000), no. 2, 161–172.
16. D. Z. Arov and M. A. Nudel’man, *Tests for the similarity of all minimal passive realizations of a fixed transfer function (scattering and resistance matrix)*, Mat. Sb. **193** (2002), no. 6, 3–24. (Russian); English transl. Sb. Math. **193** (2002), no. 6, 791–810.
17. D. Z. Arov and O. Staffans, *State/signal linear time-invariant systems theory: passive discrete time systems*, Internat. J. Robust Nonlinear Control **17** (2007), 497–548.
18. Gr. Arsene and A. Gheondea, *Completing matrix contractions*, J. Operator Theory **7** (1982), 179–189.
19. M. Bakonyi and T. Constantinescu, *Schur’s Algorithm and Several Applications*, Pitman Research Notes in Mathematics Series, Vol. 261, Longman Scientific and Technical, Harlow, 1992.
20. J. A. Ball and N. Cohen, *De Branges-Rovnyak operator models and systems theory: a survey*. Topics in Matrix and Operator Theory (Rotterdam, 1989), Oper. Theory Adv. Appl., vol. 50, Birkhäuser Verlag, Basel, 1991, pp. 93–136.

21. J. A. Ball, Ph. T. Carrol, and Yo. Uetake, *Lax-Phillips scattering theory and well-posed linear systems: a coordinate-free approach*, Mathematics of Control, Signals, and Systems **20** (2008), no. 1, 37–79.
22. Yu. M. Berezansky and M. E. Dudkin, *The direct and inverse spectral problems for the block Jacobi matrices*, Methods Funct. Anal. Topology **11** (2005), no. 4, 327–345.
23. O. Bourget, J. S. Howland, and A. Joye, *Spectral analysis of unitary band matrices*, Comm. Math. Phys. **234** (2003), 191–227.
24. L. de Branges and J. Rovnyak, *Square Summable Power Series*, Holt, Rinehart and Winston, New-York, 1966.
25. L. de Branges and J. Rovnyak, *Appendix on square summable power series. Canonical models in quantum scattering theory*, Perturbation theory and its applications in quantum mechanics (C. H. Wilcox, ed.), New-York, 1966, pp. 295–392.
26. V. M. Brodskii, *On operator colligations and their characteristic functions*, Soviet Math. Dokl. **12** (1971), 696–700.
27. M. S. Brodskii, *Triangular and Jordan Representations of Linear Operators*, Nauka, Moscow, 1969. (Russian); English transl. Transl. Math. Monographs, Vol. 32, Amer. Math. Soc., Providence, RI, 1971.
28. M. S. Brodskii, *Unitary operator colligations and their characteristic functions*, Uspekhi Mat. Nauk **33** (1978), no. 4, 141–168. (Russian); English transl. Russian Math. Surveys **33** (1978), no. 4, 159–191.
29. A. Bunse-Gerstner and L. Elsner, *Schur parameter pencils for the solution of unitary eigenproblem*, Linear Algebra Appl. **154/156** (1991), 741–778.
30. M. J. Cantero, L. Moral, and L. Velázquez, *Five-diagonal matrices and zeros of orthogonal polynomials on the unit circle*, Linear Algebra Appl. **362** (2003), 29–56.
31. Z. Ceausescu, C. Foias, *On intertwining dilations. V*, Acta Sci. Math. (Szeged) **40** (1978), 9–32; *Corrections* **41** (1979).
32. S. Clark, F. Gesztesy, and M. Zinchenko, *Weyl-Titchmarsh theory and Borg-Marchenko type uniqueness results for CMV operators with matrix valued Verblunsky coefficients*, Operators and Matrices **1** (2007), no. 4, 535–592.
33. T. Constantinescu, *On the structure of Naimark dilation*, J. Operator Theory **12** (1984), 159–175.
34. T. Constantinescu, *Operator Schur algorithm and associated functions*, Math. Balcanica **2** (1988), 244–252.
35. T. Constantinescu, *On the structure of positive Toeplitz forms*, Dilation theory, Toeplitz operators, and other topics, (Timisoara/Herculane, 1982), Oper. Theory Adv. Appl. **11** (1983), 127–149.
36. T. Constantinescu, *Schur Parameters, Factorization and Dilation Problems*, Oper. Theory Adv. Appl., Vol. 82, Birkhäuser Verlag, Basel, 1996.
37. D. Damanik, A. Pushnitskii, and B. Simon, *The analytical theory of matrix orthogonal polynomials*, Surv. Approx. Theory **4** (2008), 1–85.
38. Ch. Davis, W. M. Kahan, and H. F. Weinberger, *Norm preserving dilations and their applications to optimal error bounds*, SIAM J. Numer. Anal. **19** (1982), 445–469.
39. Ph. Delsarte, Y. Genin, and Y. Kamp, *Orthogonal polynomial matrices on the unit circle*, IEEE Trans. Circuits and Systems **25** (1978), no. 3, 149–160.
40. Ph. Delsarte, Y. Genin, and Y. Kamp, *Schur parametrization of positive definite block-Toeplitz systems*, SIAM J. Appl. Math. **36** (1979), no. 1, 34–46.
41. V. K. Dubovoy, *Shift operators contained in contractions, Schur parameters and pseudocontinuable Schur functions*, Oper. Theory Adv. Appl. **165** (2006), 175–250.
42. V. K. Dubovoj, B. Fritzsche, and B. Kirstein, *Matricial Version of the Classical Schur Problem*, Teubner-Texte zur Mathematik, Vol. 129, B. G. Teubner, Verlagsgesellschaft, Stuttgart, 1992.
43. M. E. Dudkin, *The inner structure of the block Jacobi type unitary matrices connected with the corresponding direct and inverse spectral problems*, Methods Funct. Anal. Topology **14** (2008), no. 2, 168–176.
44. C. Foias and A. E. Frazho, *The Commutant Lifting Approach to Interpolation Problems*, Oper. Theory Adv. Appl., Vol. 44, Birkhäuser Verlag, Basel, 1990.
45. B. Fritzsche, V. Katsnelson, and B. Kirstein, *The Schur algorithm in terms of system realizations*, arXiv: 0805.4732 [math. CV] v 1, 30 May 2008.
46. Ya. L. Geronimus, *On polynomials orthogonal on the circle, on trigonometric moment problem, and on allied Carathéodory and Schur functions*, Mat. Sb. **15** (1944), 99–130. (Russian)
47. F. Gesztesy and M. Zinchenko, *A Borg-type theorem associated with orthogonal polynomials on the unit circle*, J. London Math. Soc. **74** (2006), 757–777.

48. L. Golinskii and M. Kudryavtsev, *An inverse spectral theory for finite CMV matrices*, arXiv:0705.4353 [math. SP] v 1, 30 May 2007.
49. L. Golinskii and M. Kudryavtsev, *Rational interpolation and mixed inverse spectral problem for finite CMV matrices*, arXiv:0709.2214 [math. SP] v 1, 14 Sep 2007.
50. L. Golinskii, A. Kheifets, F. Peherstorfer, and P. Yuditskii, *Faddeev-Marchenko scattering for CMV matrices and the strong Szego theorem*, arXiv:0807.4017 [math. SP] v 1, 25 Jul 2008.
51. J. W. Helton, *The characteristic functions of operator theory and electrical network realization*, Indiana Univ. Math. J. **22** (1972/73), 403–414.
52. J. W. Helton, *Discrete time systems, operator models, and scattering theory*, J. Funct. Anal. **16** (1974), 15–38.
53. T. Kailath and A. M. Bruckstein, *Naimark dilations, state-space generators and transmission lines*, Oper. Theory Adv. Appl. **17** (1986), 173–186.
54. A. Kheifets, F. Peherstorfer, and P. Yuditskii, *On scattering for CMV matrices*, arXiv:0706.2970 [math. SP] v 1, 20 Jun 2007.
55. P. D. Lax and R. S. Phillips, *Scattering Theory*. Academic Press, New York, 1967.
56. N. K. Nikolskiĭ and V. I. Vasyunin, *Notes on two function models*, Proc. Conf. in honour of de Branges' proof of the Bieberbach conjecture (West Lafayette, Ind., 1985), Math. Surveys Monogr., vol. 21, Amer. Math. Soc., Providence, RI, 1986, pp. 113–141.
57. N. K. Nikolskiĭ and V. I. Vasyunin, *A unified approach to function models, and the transcription problem*, Oper. Theory Adv. Appl. **41** (1988), 405–434.
58. B. S. Pavlov, *The continuous spectrum of resonances on a nonphysical sheet*, Dokl. Akad. Nauk SSSR **206** (1972), 1301–1304. (Russian)
59. B. S. Pavlov, *Conditions for separation of the spectral components of a dissipative operator*, Izv. Akad. Nauk SSSR, Ser. Mat. **39** (1975), 123–148. (Russian)
60. B. S. Pavlov, *A remark on spectral meaning of the symmetric functional model*, Oper. Theory Adv. Appl. **154** (2004), 163–177.
61. J. J. Schäffer, *On unitary dilations of contractions*, Proc. Amer. Math. Soc. **6** (1955), 322.
62. I. Schur, *Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind*, I, II, J. Reine Angew. Math. **147** (1917), 205–232; **148** (1918), 122–145.
63. Yu. L. Shmul'yan, *Generalized fractional-linear transformations of operator balls*, Sib. Mat. Zh. **21** (1980), no. 5, 114–131. (Russian); English transl. Siberian Math. J. **21** (1980), no. 5, 728–740.
64. Yu. L. Shmul'yan, *Certain stability properties for analytic operator-valued functions*, Mat. Zametki **20** (1976), no. 4, 511–520. (Russian); English transl. Math. Notes **20** (1976), no. 4, 843–848.
65. Yu. L. Shmul'yan and R. N. Yanovskaya, *Blocks of a contractive operator matrix*, Izv. Vyssh. Uchebn. Zaved., Mat. **7** (1981), 72–75. (Russian)
66. B. Simon, *Orthogonal Polynomials on the Unit Circle*. Part 1. *Classical Theory*, Part 2. *Spectral Theory*, Amer. Math. Soc. Colloquium Publications, Providence, RI, 2005.
67. B. Simon, *Analogues of the m -function in the theory of orthogonal polynomials on the unit circle*, J. Comput. Appl. Math. **171** (2004), no. 1–2, 411–424.
68. B. Simon, *CMV matrices: five years after*, J. Comput. Appl. Math. **208** (2007), no. 1, 120–154.
69. O. Staffans, *Well-Posed Linear Systems*, Encyclopedia of Mathematics and its Applications, 103, Cambridge University Press, Cambridge, 2005.
70. O. Staffans, *Passive linear discrete time-invariant systems*, Proceedings of the International Congress of Mathematicians, vol. III, Eur. Math. Soc., Zurich, 2006, pp. 1367–1388.
71. B. Sz. Nagy and C. Foias, *Harmonic Analysis of Operators on Hilbert Space*, North-Holland, New York, 1970.
72. M. Zinchenko, *Trace formulas and a Borg-type theorem for CMV operators with matrix-valued coefficients*, arXiv:0808.0382 [math-ph] v 1, 4 Aug 2008.
73. D. S. Watkins, *Some perspectives on the eigenvalue problem*, SIAM Rev. **35** (1993), 430–471.

DEPARTMENT OF MATHEMATICAL ANALYSIS, EAST UKRAINIAN NATIONAL UNIVERSITY, 20-A KVARTAL MOLODIZHNY, LUGANSK, 91034, UKRAINE

E-mail address: yury_arlinskii@yahoo.com

E-mail address: yma@snu.edu.ua

Received 16/02/2009