

CONNECTED COMPONENTS OF PARTITION PRESERVING DIFFEOMORPHISMS

SERGIY MAKSYMENKO

ABSTRACT. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a homogeneous polynomial and $\mathcal{S}(f)$ be the group of diffeomorphisms h of \mathbb{R}^2 preserving f , i.e. $f \circ h = f$. Denote by $\mathcal{S}_{\text{id}}(f)^r$, ($0 \leq r \leq \infty$), the identity component of $\mathcal{S}(f)$ with respect to the weak Whitney C_W^r -topology. We prove that $\mathcal{S}_{\text{id}}(f)^\infty = \dots = \mathcal{S}_{\text{id}}(f)^1$ for all f and that $\mathcal{S}_{\text{id}}(f)^1 \neq \mathcal{S}_{\text{id}}(f)^0$ if and only if f is a product of at least two distinct irreducible over \mathbb{R} quadratic forms.

1. INTRODUCTION

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a homogeneous polynomial of degree $p \geq 1$. Thus up to sign we can write

$$(1.1) \quad f(x, y) = \pm \prod_{i=1}^l L_i^{\alpha_i}(x, y) \cdot \prod_{j=1}^k Q_j^{\beta_j}(x, y),$$

where every L_i is a linear function, Q_j is a positive definite quadratic form, $\alpha_i, \beta_j \geq 1$, and

$$\frac{L_i}{L_{i'}} \neq \text{const for } i \neq i', \quad \frac{Q_j}{Q_{j'}} \neq \text{const for } j \neq j'.$$

Denote by $\mathcal{S}(f) = \{h \in \mathcal{D}(\mathbb{R}^2) : f \circ h = f\}$ the stabilizer of f with respect to the right action of the group $\mathcal{D}(\mathbb{R}^2)$ of C^∞ -diffeomorphisms of \mathbb{R}^2 on the space $C^\infty(\mathbb{R}^2, \mathbb{R})$. It consists of diffeomorphisms of \mathbb{R}^2 preserving every level-set $f^{-1}(c)$ of f , $c \in \mathbb{R}$.

Let $\mathcal{S}_{\text{id}}(f)^r$, $0 \leq r \leq \infty$, be the identity component of $\mathcal{S}(f)$ with respect to the weak Whitney C_W^r -topology. Thus $\mathcal{S}_{\text{id}}(f)^r$ consists of diffeomorphisms $h \in \mathcal{S}(f)$ isotopic in $\mathcal{S}(f)$ to $\text{id}_{\mathbb{R}^2}$ via (an f -preserving isotopy) $H : \mathbb{R}^2 \times I \rightarrow \mathbb{R}^2$ whose partial derivatives in $(x, y) \in \mathbb{R}^2$ up to order r continuously depend on (x, y, t) , see Section 2 for a precise definition. Then it is easy to see that

$$\mathcal{S}_{\text{id}}(f)^\infty \subset \dots \subset \mathcal{S}_{\text{id}}(f)^r \subset \dots \subset \mathcal{S}_{\text{id}}(f)^1 \subset \mathcal{S}_{\text{id}}(f)^0.$$

It follows from results in [12, 13] that $\mathcal{S}_{\text{id}}(f)^\infty = \mathcal{S}_{\text{id}}(f)^1$. Moreover, it is actually proved in [9] that $\mathcal{S}_{\text{id}}(f)^\infty = \mathcal{S}_{\text{id}}(f)^0$ for $p \leq 2$, see also [11]. The aim of this note is to prove the following theorem describing the relation between $\mathcal{S}_{\text{id}}(f)^r$ for all $p \geq 1$.

Theorem 1.1. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a homogeneous polynomial of degree $p \geq 1$. Then $\mathcal{S}_{\text{id}}(f)^\infty = \dots = \mathcal{S}_{\text{id}}(f)^1$. Moreover, $\mathcal{S}_{\text{id}}(f)^1 \neq \mathcal{S}_{\text{id}}(f)^0$ if and only if f is a product of at least two distinct definite quadratic forms, i.e. $f = Q_1^{\beta_1} \dots Q_k^{\beta_k}$ for $k \geq 2$.*

This theorem is based on a rather general result about partition preserving diffeomorphisms, see Theorem 4.7. The applications of Theorem 1.1 will be given in another paper concerning smooth functions on surfaces with isolated singularities.

2000 *Mathematics Subject Classification.* 57S05.

Key words and phrases. Diffeomorphisms, weak Whitney topologies.

This research is partially supported by Grant of President of Ukraine Φ -26/418-2008.

Structure of the paper. In Section 2 we describe homotopies which induce continuous paths into functional spaces with the weak Whitney C_W^r -topologies. Section 3 introduces the so-called *singular partitions* of manifolds being the main object of the paper. Section 4 contains the main result, Theorem 4.7, about invariant contractions of singular partitions. In Section 5 an application of this theorem to local extremes of smooth functions is given. Section 6 contains a description of the group of linear symmetries of f . Finally in Section 7 we prove Theorem 1.1.

2. r -HOMOTOPIES

Denote $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and $\overline{\mathbb{N}}_0 = \mathbb{N} \cup \{0, \infty\}$.

Let M, N be two smooth manifolds of dimensions m and n respectively. Then for every $r \in \overline{\mathbb{N}}_0$ the space $C^r(M, N)$ admits the so-called *weak* Whitney topology denoted by C_W^r , see e.g. [8, 5].

Recall, e.g. [7, § 44.IV] that there exists a homeomorphism

$$C^0(I, C^0(M, N)) \approx C^0(M \times I, N)$$

with respect to the corresponding C_W^0 -topologies (also called *compact open* ones) associating to every (continuous) path $w : I \rightarrow C^0(M, N)$ a homotopy $H : M \times I \rightarrow N$ defined by $H(x, t) = w(t)(x)$.

We will now describe homotopies inducing continuous paths $w : I \rightarrow C^r(M, N)$ with respect to C_W^r -topologies.

Definition 2.1. Let $H : M \times I \rightarrow N$ be a homotopy and $r \in \overline{\mathbb{N}}_0$. We say that H is an *r -homotopy* if

- (1) $H_t : M \rightarrow N$ is C^r for every $t \in I$;
- (2) partial derivatives of $H(x, t)$ with respect to x , up to order r , continuously depend on (x, t) .

More precisely, let $z \in M \times I$. Then in some local coordinates at z we can regard H as a map

$$H = (H_1, \dots, H_n) : \mathbb{R}^m \times I \rightarrow \mathbb{R}^n$$

such that for every fixed t and i the function $H_i(x, t)$ is C^r . Condition (2) requires that for every $i = 1, \dots, n$ and every non-negative integer vector $k = (k_1, \dots, k_m)$ of norm $|k| = \sum_{j=1}^m k_j \leq r$ the function

$$\frac{\partial^{|k|} H_i}{\partial x_1^{k_1} \dots \partial x_m^{k_m}}(x_1, \dots, x_m, t)$$

continuously depend on (x, t) .

Thus a 0-homotopy H is just the usual homotopy.

It easily follows from the definition of C_W^r -topologies that a path $w : I \rightarrow C^r(M, N)$ is continuous in the standard topology of I and C_W^r -topology of $C^r(M, N)$ if and only if the corresponding homotopy $H : M \times I \rightarrow N$ is an r -homotopy.

We can also define a *C^r -homotopy* as a C^r -map $M \times I \rightarrow N$. Evidently, every C^r -homotopy is an r -homotopy as well, but the converse is not true.

Example 2.2. Let $H : \mathbb{R} \times I \rightarrow \mathbb{R}$ be given by

$$H(x, t) = \begin{cases} t \ln(x^2 + t^2), & (x, t) \neq (0, 0), \\ 0, & (x, t) = (0, 0). \end{cases}$$

Then H is continuous, while $\frac{\partial H}{\partial x} = \frac{2tx}{x^2+y^2}$ is C^∞ for every fixed t as a function in x but discontinuous at $(0, 0)$ as a function in (x, t) . In other words H is a 0-homotopy but not a 1-homotopy.

Moreover, define $G : \mathbb{R} \times I \rightarrow \mathbb{R}$ by $G(x, t) = \int_0^x H(y, t) dy$. Then G a 1-homotopy but not a 2-homotopy.

3. SINGULAR PARTITIONS OF MANIFOLDS

Let M be a smooth manifold equipped with a partition $\mathcal{P} = \{\omega_i\}_{i \in \Lambda}$, i.e., a family of subsets ω_i such that

$$M = \bigcup_{i \in \Lambda} \omega_i, \quad \omega_i \cap \omega_j = \emptyset \quad (i \neq j).$$

In general Λ may be even uncountable and ω_i are not necessarily closed in M . Let also Λ' be a (possibly empty) subset of Λ and $\Sigma = \{\omega_i\}_{i \in \Lambda'}$ be a subfamily of \mathcal{P} thought of as a set of “singular” elements. Then the pair $\Theta = (\mathcal{P}, \Sigma)$ will be called a *singular partition* of M .

Example 3.1. Let F be a vector field on M , \mathcal{P}_F the set of orbits of F , and Σ_F the set of singular points of F . Then the pair $\Theta_F = (\mathcal{P}_F, \Sigma_F)$ will be called the *singular partition of F* .

Example 3.2. Let $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a smooth map, $x \in \mathbb{R}^m$ a point, and $J(f, x)$ the Jacobi matrix of f at x . Then $x \in \mathbb{R}^m$ is called *critical* for f if $\text{rank } J(f, x) < \min\{m, n\}$. Otherwise x is *regular*. This definition naturally extends to maps between manifolds.

Let M, N be smooth manifolds and $f : M \rightarrow N$ a smooth map. Denote by Σ_f the set of critical points of f . Consider the following partition \mathcal{P}_f of M : a subset $\omega \subset M$ belongs to \mathcal{P}_f iff ω is either a critical point of f , or a connected component of the set of the form $f^{-1}(y) \setminus \Sigma_f$ for some $y \in N$. Then the pair $\Theta_f = (\mathcal{P}_f, \Sigma_f)$ will be called the *singular partition of f* . Evidently, every $\omega \in \mathcal{P}_f \setminus \Sigma_f$ is a submanifold of M .

Example 3.3. Assume that, in Example 3.2, $\dim M = \dim N + 1$ and both M and N are orientable. Then every element of $\mathcal{P}_f \setminus \Sigma_f$ is one-dimensional and orientations of M and N allow to *coherently orient all the elements of $\mathcal{P}_f \setminus \Sigma_f$* . Moreover, it is even possible to construct a vector field F on M such that the singular partitions Θ_f and Θ_F coincide.

In particular, let M be an orientable surface and $f : M \rightarrow \mathbb{R}$ be a smooth function. Then M admits a symplectic structure, and in this case we can assume that F is the corresponding Hamiltonian vector field of f .

Example 3.4. Let \mathcal{F} be a foliation on M with singular leaves, \mathcal{P} be the set of leaves of \mathcal{F} , and Σ be the set of its singular leaves (having non-maximal dimension). Then the pair $\Theta_{\mathcal{F}} = (\mathcal{P}, \Sigma)$ will be called the *singular partition of \mathcal{F}* . This example generalizes all previous ones.

Let $\Theta = (\mathcal{P}, \Sigma)$ be a singular partition on M . For every open subset $V \subset M$ denote by $\mathcal{E}(\Theta, V)$ the subset of $C^\infty(V, M)$ consisting of maps $f : V \rightarrow M$ such that

- (1) $f(\omega_i \cap V) \subset \omega_i$ for all $\omega_i \in \mathcal{P}$ and
- (2) f is a local diffeomorphism at every point z belonging to some singular element $\omega \in \Sigma$.

Let also $\mathcal{D}(\Theta, V)$ be the subset of $\mathcal{E}(\Theta, V)$ consisting of *immersions*, i.e., *local diffeomorphisms*. For $V = M$ we abbreviate

$$\mathcal{E}(\Theta) = \mathcal{E}(\Theta, M), \quad \mathcal{D}(\Theta) = \mathcal{D}(\Theta, M).$$

For every $r \in \overline{\mathbb{N}}_0$ denote by $\mathcal{E}_{\text{id}}(\Theta, V)^r$, resp. $\mathcal{D}_{\text{id}}(\Theta, V)^r$, the path-component of the identity inclusion $i_V : V \subset M$ in $\mathcal{E}(\Theta, V)$, resp. in $\mathcal{D}(\Theta, V)$, with respect to the induced C_W^r -topology, see Section 2.

Thus $\mathcal{E}_{\text{id}}(\Theta, V)^r$ (resp. $\mathcal{D}_{\text{id}}(\Theta, V)^r$) consists of maps (resp. immersions) $V \subset M$ which are r -homotopic (r -isotopic) to $i_V : V \subset M$ in $\mathcal{E}(\Theta, V)$ (resp. in $\mathcal{D}(\Theta, V)$).

Evidently,

$$(3.1) \quad \mathcal{E}_{\text{id}}(\Theta, V)^\infty \subset \dots \subset \mathcal{E}_{\text{id}}(\Theta, V)^1 \subset \mathcal{E}_{\text{id}}(\Theta, V)^0,$$

and similar relations hold for $\mathcal{D}_{\text{id}}(\Theta, V)^r$.

The following notion turns out to be useful for studying singular partitions of vector fields.

3.5. Shift-map of a vector field. Let F be a vector field on M and

$$\Phi : M \times \mathbb{R} \supset \text{dom}(\Phi) \rightarrow M$$

be the local flow of F defined on some open neighbourhood $\text{dom}(\Phi)$ of $M \times 0$ in $M \times \mathbb{R}$. For every open subset $V \subset M$ let also

$$\text{func}(\Phi, V) = \{\alpha \in C^\infty(V, \mathbb{R}) : \Gamma_\alpha \subset \text{dom}(\Phi)\},$$

where $\Gamma_\alpha = \{(x, \alpha(x)) : x \in V\} \subset M \times \mathbb{R}$ is the graph of α . Then $\text{func}(\Phi, V)$ is the largest subset of $C^\infty(V, \mathbb{R})$ on which the following *shift-map* is defined:

$$\varphi_V : \text{func}(\Phi, V) \rightarrow C^\infty(V, M), \quad \varphi_V(\alpha)(x) = \Phi(x, \alpha(x)),$$

for $\alpha \in \text{func}(\Phi, V)$, $x \in V$.

Lemma 3.6. *Let Θ_F be the singular partition of M by orbits of F . Then*

$$(3.2) \quad \text{im}(\varphi_V) \subset \mathcal{E}_{\text{id}}(\Theta_F, V)^\infty.$$

Moreover, if $\mathcal{D}_{\text{id}}(\Theta_F, V)^r \subset \text{im}(\varphi)$ for some $r \in \overline{\mathbb{N}}_0$, then

$$\mathcal{D}_{\text{id}}(\Theta_F, V)^\infty = \dots = \mathcal{D}_{\text{id}}(\Theta_F, V)^{r+1} = \mathcal{D}_{\text{id}}(\Theta_F, V)^r.$$

Proof. Let $\alpha \in \text{func}(\Phi, V)$ and $f = \varphi(\alpha)$, i.e., $f(x) = \Phi(x, \alpha(x))$. Then $f(\omega \cap V) \subset \omega$ for every orbit of F . Moreover by [9, Lemma 20] f is a local diffeomorphism at a point $x \in V$ iff $d\alpha(F)(x) \neq -1$, where $d\alpha(F)(x)$ is the Lie derivative of α along F at x . Hence f is so at every singular point z of F , since $d\alpha(F)(z) = 0 \neq -1$, [9, Corollary 21]. Therefore $f \in \mathcal{E}(\Theta_F, V)$. Moreover an ∞ -homotopy of f to $i_V : V \subset M$ in $\mathcal{E}(\Theta_F, V)$ can be given by $f_t(x) = \Phi(x, t\alpha(x))$. Thus $f \in \mathcal{E}_{\text{id}}(\Theta_F, V)^\infty$.

Finally, suppose that $f \in \mathcal{D}_{\text{id}}(\Theta_F, V)^r$. Then the restriction of f to any non-constant orbit ω of F is an orientation preserving local diffeomorphism. Therefore $d\alpha(F)(z) > -1$ on all of V . Hence $d(t\alpha)(F)(z) > -1$ for all $t \in I$ as well, i.e., $f_t \in \mathcal{D}(\Theta_F, V)$. This implies that $f \in \mathcal{D}_{\text{id}}(\Theta_F, V)^\infty$. \square

Example 3.7. Let A be a real non-zero $(m \times m)$ -matrix, $F(x) = Ax$ be the corresponding linear vector field on \mathbb{R}^m , and V be a neighbourhood of the origin 0 . Then the shift-map φ_V is given by

$$\varphi(\alpha)(x) = \Phi(x, \alpha(x)) = e^{A\alpha(x)}x.$$

It is shown in [9] that in this case $\text{im}(\varphi_V) = \mathcal{E}_{\text{id}}(\Theta_F, V)^0$. Hence for all $r \in \overline{\mathbb{N}}_0$ we have

$$\text{im}(\varphi_V) = \mathcal{E}_{\text{id}}(\Theta_F, V)^\infty = \dots = \mathcal{E}_{\text{id}}(\Theta_F, V)^r,$$

$$\mathcal{D}_{\text{id}}(\Theta_F, V)^\infty = \dots = \mathcal{D}_{\text{id}}(\Theta_F, V)^r.$$

4. INVARIANT CONTRACTIONS

Let $\Theta = (\mathcal{P}, \Sigma)$ be a singular partition on a manifold M . We will say that a subset $V \subset M$ is Θ -invariant, if it consists of full elements of Θ , i.e., if $\omega \in \mathcal{P}$ and $\omega \cap V \neq \emptyset$, then $\omega \subset V$.

Definition 4.1. Let $Z \subset M$ be a closed subset such that every point $z \in Z$ is a singular element of Θ , i.e., $\{z\} \in \Sigma$. Say that Θ has an **invariant r -contraction to Z** if there exists a closed Θ -invariant neighbourhood V of Z being a smooth submanifold of M and a homotopy $r : V \times I \rightarrow V$ such that:

- (i) $r_1 = \text{id}_V$;

- (ii) r_0 is a proper retraction of V to Z , i.e., $r_0(V) = Z$, $r_0(z) = z$ for $z \in Z$, and $r_0^{-1}(K)$ is compact for every compact $K \subset Z$;
- (iii) for every $t \in (0, 1]$ the map r_t is a closed C^r -embedding of V into V such that for each $\omega \in \mathcal{P}$ (resp. $\omega \in \Sigma$) its image $r_t(\omega)$ is also an element of \mathcal{P} (resp. Σ);
- (iv) for each $z \in Z$ the set $V_z = r_0^{-1}(z)$ is Θ -invariant, and $r_t(V_z) \subset V_z$ for all $t \in I$.

Since V is Θ -invariant, it follows from (iii) that so is its image $r_t(V)$.

Example 4.2. Define $f : \mathbb{R}^m \rightarrow \mathbb{R}$ by $f(x_1, \dots, x_m) = \sum_{i=1}^m x_i^2$. Evidently, the singular partition Θ_f consists of the origin 0 and concentric spheres centered at 0 . For every $s > 0$ let $V_s = f^{-1}[0, s^2]$ be a closed m -disk of radius s . Then V_s is Θ_f -invariant and its invariant contraction of V_s to $Z = \{0\}$ can be given by $r(x, t) = tx$.

Example 4.3. The previous example can be parametrized as follows. Let $p : M \rightarrow Z$ be an m -dimensional vector bundle over a connected, smooth manifold Z . We will identify Z with the image $Z \subset M$ of the corresponding zero-section of p . Suppose that we are given a norm $\|\cdot\|$ on fibers such that the following function $f : M \rightarrow \mathbb{R}$ is smooth:

$$f(\xi, z) = \|\xi\|^2, \quad z \in Z, \quad \xi \in p^{-1}(z).$$

Define the following singular partition $\Theta = (\mathcal{P}, \Sigma)$ on M , where \mathcal{P} consists of the subsets $\omega_{s,z} = f^{-1}(s) \cap p^{-1}(z)$ for $s \geq 0$ and $z \in Z$, and $\Sigma = \{\omega_{0,z} = \{z\} : z \in Z\}$ consists of points of Z . Thus every fiber $p^{-1}(z)$ is Θ -invariant, and the restriction of Θ to $p^{-1}(z)$ is the same as the one in Example 4.2.

Fix $s > 0$ and put $V = f^{-1}[0, s]$. Then V is Θ -invariant and a Θ -invariant contraction of V to Z can be given by $r(\xi, z, t) = (t\xi, z)$.

We will now generalize these examples. Let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a smooth function and suppose that there exists a neighbourhood V of 0 and smooth functions $\alpha_1, \dots, \alpha_m : V \rightarrow \mathbb{R}$ such that

$$(4.1) \quad f = \alpha_1 \cdot f'_{x_1} + \dots + \alpha_m \cdot f'_{x_m}.$$

Equivalently, let $\Delta(f, 0)$ be the *Jacobi ideal* of f in $C^\infty(V, \mathbb{R})$ generated by partial derivatives of f . Then (4.1) means that $f \in \Delta(f, 0)$.

For instance, let f be *quasi-homogeneous of degree d with weights s_1, \dots, s_m* , i.e., $f(t^{s_1}x_1, \dots, t^{s_m}x_m) = t^d f(x_1, \dots, x_m)$ for $t > 0$, see e.g [1, §12]. Equivalently, we may require that the function

$$g(x_1, \dots, x_m) = f(x_1^{s_1}, \dots, x_m^{s_m})$$

be homogeneous of degree d . Then the following *Euler identity* holds true:

$$f = \frac{x_1}{s_1} f'_{x_1} + \dots + \frac{x_m}{s_m} f'_{x_m}.$$

In particular, f satisfies (4.1). Moreover in the complex analytical case¹ the identity (4.1) characterizes quasi-homogeneous functions, see [21].

Lemma 4.4. *Let $f : M \rightarrow \mathbb{R}$ be a smooth function and $z \in M$ be an isolated local minimum of f . Suppose that f satisfies condition (4.1) at z , i.e., $f \in \Delta(f, z)$. Then the singular partition Θ_f admits an invariant contraction to z .*

Proof. Since the situation is local, we may assume that $M = \mathbb{R}^m$, $z = 0$ is a unique critical point of f being its global minimum, $f(0) = 0$, and there exists an $\varepsilon > 0$ such that $V = f^{-1}[0, \varepsilon]$ is a smooth compact m -dimensional manifold with boundary $L = f^{-1}(\varepsilon)$.

First we give a precise description of the partition \mathcal{P}_f on V . Let F be any gradient like vector field on V for f , i.e., $df(F)(x) > 0$ for $x \neq 0$. Then following [15, Th. 3.1] we can construct a diffeomorphism

$$\eta : V \setminus \{0\} \rightarrow L \times (0, \varepsilon]$$

¹I would like to thank V. A. Vasilyev for referring me to the paper [21] by K. Saito.

such that $f \circ \eta^{-1}(y, t) = t$ for all $(y, t) \in L \times (0, \varepsilon]$, see Figure 4.1.

Let $CL = L \times [0, \varepsilon] / \{L \times 0\}$ be the cone over L and $L_t = f^{-1}(t)$ for $t \in (0, \varepsilon]$. Since $\text{diam}(L_t) \rightarrow 0$ when $t \rightarrow 0$, we obtain that η extends to a homeomorphism $\eta : V \rightarrow CL$ by $\eta(0) = \{L \times 0\}$.

For every $t \in (0, \varepsilon]$ put $L_t = f^{-1}(t)$. Then η diffeomorphically maps L_t onto $L \times \{t\}$.




FIGURE 4.1

Lemma 4.5. *L is homotopy equivalent to S^{m-1} .*

We will prove this lemma below. Then it will follow from the generalized Poincaré conjecture that L is homeomorphic with the sphere S^{m-1} , and even diffeomorphic to S^{m-1} if $k = m - 1 \neq 4$. For $k = 1, 2$ this statement is rather elementary, for $k = 3$ this follows from a recent work of G. Perelman [18, 19], for $k = 4$ from M. Freedman [3], and for $k \geq 5$ from S. Smale [23], see also [14].

In particular, every L_t is connected, whence the partition \mathcal{P}_f on V consists of a unique singular element $\{0\} \in \Sigma_f$ and sets $L_t, t \in (0, \varepsilon]$.

Let us recall the definition of η . Notice that every orbit of F starts at 0 and transversely intersects every L_t . For each $x \in V \setminus \{0\}$ denote by $q(x)$ a unique point of the intersection of the orbit of x with $L = L_\varepsilon = \partial V$. Then $\eta : V \setminus \{0\} \rightarrow L \times (0, \varepsilon]$ can be given by the following formula:

$$\eta(x) = (q(x), f(x)).$$

Also notice that if $\phi : [0, \varepsilon] \rightarrow [0, \varepsilon]$ is a (not necessarily surjective) C^∞ embedding such that $\phi(0) = 0$, then we can define the embedding

$$w_\phi : CL \rightarrow CL, \quad w_\phi(y, t) = (y, \phi(t))$$

and therefore the embedding $r_\phi = \eta \circ w_\phi \circ \eta^{-1} : V \rightarrow V$. Then r_ϕ is C^∞ on $V \setminus 0$ and diffeomorphically maps L_t onto $L_{\phi(t)}$ for all $t \in (0, \varepsilon]$. Moreover $r_\phi(0) = 0$, but in general r_ϕ is not even smooth at 0.

Suppose now that $f \in \Delta(f, 0)$, i.e., we have a presentation (4.1). Consider the following vector field

$$F = \alpha_1 \frac{\partial}{\partial x_1} + \dots + \alpha_m \frac{\partial}{\partial x_m}.$$

Then (4.1) means that $f = df(F)$. Since $f(x) > 0$ for $x \neq 0$, it follows that F is a gradient like vector field for f . Therefore we can construct a homeomorphism $\eta : V \rightarrow CL$ using F as above. It follows from [10] that in our case this η has the following feature:

- if $\phi : [0, \varepsilon] \rightarrow [0, \varepsilon]$ is a C^∞ embedding such that $\phi(0) = 0$, then the corresponding embedding $r_\phi : V \rightarrow V$ is a diffeomorphism onto its image. Moreover, if $\phi_s, (s \in I)$, is a C^∞ isotopy, then so is $r_{\phi_s} : V \rightarrow V$.

In particular, consider the following homotopy

$$\phi : [0, \varepsilon] \times I \rightarrow [0, \varepsilon], \quad \phi(t, s) = t(1 - s)$$

which contracts $[0, \varepsilon]$ to a point and being an isotopy for $t > 0$. Then the induced homotopy $r : V \times I \rightarrow V$ is an invariant ∞ -contraction of Θ_f to 0. \square

Proof of Lemma 4.5. It suffices to establish that

$$(4.2) \quad \pi_k L \approx \pi_k S^{m-1} = \begin{cases} 0, & k = 0, \dots, m-2, \\ \mathbb{Z}, & k = m-1. \end{cases}$$

Then the generator $\mu : S^{m-1} \rightarrow L$ of $\pi_{m-1} L \approx \mathbb{Z}$ will yield isomorphisms of the homotopy groups $\pi_k S^{m-1} \approx \pi_k L$ for all $k \leq m-1 = \dim S^{m-1} = \dim L$. Now by the well-known Whitehead's theorem μ will be a homotopy equivalence between S^{m-1} and L .

For the calculation of homotopy groups of L consider the exact sequence of homotopy groups of the pair (V, L) :

$$(4.3) \quad \cdots \rightarrow \pi_{k+1}(V, L) \rightarrow \pi_k L \rightarrow \pi_k V \rightarrow \cdots$$

Since V is homeomorphic with the cone CL , V is contractible, whence $\pi_k V = 0$ for all $k \geq 0$.

Moreover, $\pi_{k+1}(V, L) = 0$ for all $k = 0, \dots, m-2$. Indeed, let

$$\xi : (D^{k+1}, S^k) \rightarrow (V, L)$$

be a continuous map. We have to show that ξ is homotopic (as a map of pairs) to a map into L . Since $k+1 \leq m-1 < \dim V$, ξ is homotopic to a map into $V \setminus \{0\}$. But L is a deformation retract of $V \setminus \{0\}$, therefore ξ is homotopic to a map into L .

Now it follows from (4.3) that $\pi_k L = 0$ for $k = 0, \dots, m-2$. Hence from Hurewicz's theorem we obtain that $\pi_{m-1} L \approx H_{m-1}(L, \mathbb{Z})$. It remains to note that L is a connected closed orientable $(m-1)$ -manifold, whence (by Poincaré duality) $H_{m-1}(L, \mathbb{Z}) \approx \mathbb{Z}$. This proves (4.2). \square

Remark 4.6. If f is a quasi-homogeneous function of degree d with weights s_1, \dots, s_m , then we can define an invariant contraction by

$$r(x_1, \dots, x_m, t) = (t^{s_1} x_1, \dots, t^{s_m} x_m).$$

If f is homogeneous, then we can even put $r(x, t) = tx$.

Theorem 4.7. Let $\Theta = (\mathcal{P}, \Sigma)$ be a singular partition on a manifold M , and $Z \subset M$ be a closed subset such that every $z \in Z$ is a singular element of \mathcal{P} , i.e., $\{z\} \in \Sigma$. Suppose that Θ has an invariant ∞ -contraction to Z defined on a Θ -invariant neighbourhood V of Z . Let also $h \in \mathcal{E}(\Theta, V)$ be a map fixed outside some neighbourhood U of z such that $\bar{U} \subset \text{Int} V$. Then $h \in \mathcal{E}_{\text{id}}(\Theta, V)^0$, though h not necessarily belongs to $\mathcal{E}_{\text{id}}(\Theta, V)^r$ for some $r \geq 1$.

Remark 4.8. Let Θ_f be the singular partition of $f(x) = \|x\|^2$ as in Example 4.2, V be the unit n -disk centered at 0 and $r : V \times I \rightarrow V$ be the invariant contraction of Θ_f to a point $Z = \{0\}$ defined by $r(x, t) = tx$. Then a 0-homotopy between h and id_V can be defined by

$$H_t(x) \begin{cases} th(\frac{x}{t}), & t > 0, \\ 0, & t = 0. \end{cases}$$

c.f. [5, Ch. 4, Theorems 5.3 & 6.7]. Theorem 4.7 generalizes this example.

Proof of Theorem 4.7. Let $r : V \times I \rightarrow V$ be an invariant ∞ -contraction of Θ_F to Z . Define the following map $H : V \times I \rightarrow M$ by

$$H(x, t) = \begin{cases} r_t \circ h \circ r_t^{-1}(x), & \text{if } t > 0 \text{ and } x \in r_t(V), \\ x, & \text{otherwise.} \end{cases}$$

We claim that H is a 0-homotopy (i.e. just a homotopy) between h and the identity inclusion $i_V : V \subset M$ in $\mathcal{E}(\Theta, V)$. To make this more obvious we rewrite the formulas for H in another way.

The homotopy r can be regarded as the composition

$$r = p_1 \circ \tilde{r} : V \times I \xrightarrow{\tilde{r}} V \times I \xrightarrow{p_1} V,$$

where \tilde{r} is the following level-preserving map

$$\tilde{r} : V \times I \rightarrow V \times I, \quad \tilde{r}(x, t) = (r(x, t), t),$$

and $p_1 : V \times I \rightarrow V$ is the projection to the first coordinate. It follows from the definition




FIGURE 4.2

that r yields a level-preserving embedding $V \times (0, 1]$ to $V \times I$, see Figure 4.2. Denote

$$R' = \tilde{r}(V \times (0, 1]), \quad R = \tilde{r}(V \times I).$$

Then $R \setminus R' = Z \times 0$. Define also the following map

$$\tilde{h} : V \times I \rightarrow V \times I, \quad \tilde{h}(x, t) = (h(x), t).$$

In these terms, the homotopy H is defined by

$$H = p_1 \circ \tilde{H} : V \times I \xrightarrow{\tilde{H}} V \times I \xrightarrow{p_1} V,$$

where $\tilde{H} : V \times I \rightarrow V \times I$ is a level-preserving map given by

$$\tilde{H}(x, t) = \begin{cases} \tilde{r} \circ \tilde{h} \circ \tilde{r}^{-1}(x, t), & (x, t) \in R', \\ (x, t), & (x, t) \in (V \times I) \setminus R'. \end{cases}$$

Now we can prove that H has the desired properties.

Since $r_1 = \text{id}_V$, we have $H_1 = h$. Moreover $H_0 = \text{id}_V$.

1. Continuity of \tilde{H} on $V \times (0, 1]$. Notice that $\tilde{r} \circ \tilde{h} \circ \tilde{r}^{-1}$ is well-defined and continuous on R' . Moreover, since h is fixed on $V \setminus U$, it follows that \tilde{h} is fixed on $(V \setminus U) \times I$, whence $\tilde{r} \circ \tilde{h} \circ \tilde{r}^{-1}$ is fixed on the subset $\tilde{r}((V \setminus U) \times (0, 1]) \subset R'$. This implies that \tilde{H} is continuous on $V \times (0, 1]$.

2. Continuity of \tilde{H} when $t \rightarrow 0$. Let $z \in V$. Then $\tilde{H}(z, 0) = (z, 0)$.

Suppose that $z \in V \setminus Z$. Since Z is closed in V , \tilde{H} is also fixed and therefore continuous on some neighbourhood of $(z, 0)$ in $(V \times I) \setminus R$.

Let $z \in Z$ and let W be a neighbourhood of $(z, 0)$ in $V \times I$. We have to find another neighbourhood W' of $(z, 0)$ such that $\tilde{H}(W') \subset W$.

Recall that for every $y \in Z$ we denoted $V_y = r_0^{-1}(y)$. Then V_y is compact and Θ -invariant.

Claim. *There exist $\varepsilon > 0$ and an open neighbourhood N of z in V such that $\overline{N} \times [0, \varepsilon] \subset W$ and*

$$(4.4) \quad \tilde{r}(V_y \times [0, \varepsilon]) \subset W, \quad (y \in \overline{N} \cap Z).$$

Proof. Let N be an open neighbourhood of z such that \overline{N} is compact and $\overline{N} \times 0 \subset W$. Denote

$$Q = r_0^{-1}(\overline{N} \cap Z) = \bigcup_{y \in \overline{N} \cap Z} r_0^{-1}(y) = \bigcup_{y \in \overline{N} \cap Z} V_y.$$

Then Q is a compact subset of V , and $\tilde{r}^{-1}(W)$ is an open neighbourhood of $Q \times 0$ in $V \times I$. Hence there exists $\varepsilon > 0$ such that $Q \times [0, \varepsilon] \subset \tilde{r}^{-1}(W)$. This implies (4.4). Decreasing ε if necessary we can also assume that $\overline{N} \times [0, \varepsilon] \subset W$ as well. \square

Denote $W' = N \times [0, \varepsilon]$. We claim that $\tilde{H}(W') \subset W$.

Let $(x, t) \in W'$. If either $(x, t) \in W' \setminus R'$ or $t = 0$, then $\tilde{H}(x, t) = (x, t) \in W' \subset W$.

Suppose that $(x, t) \in W' \cap R'$. Then $t > 0$. Let also $y = r_0(x) \in Z$. Then $\tilde{r}^{-1}(x, t) \in V_y \times t$ for all $t \in I$. Hence

$$\tilde{r} \circ \tilde{h} \circ \tilde{r}^{-1}(x, t) \in \tilde{r} \circ \tilde{h}(V_y \times t) \subset \tilde{r}(V_y \times t) \stackrel{(4.4)}{\subset} W.$$

In the second inclusion we have used a Θ -invariance of V_y and the assumption that $h \in \mathcal{E}(\Theta, V)$.

3. Proof that $H_t \in \mathcal{E}(\Theta, V)$ for $t \in I$. We have to show that (i) for every $t \in I$ the mapping H_t is C^∞ , (ii) $H_t(\omega) \subset \omega$ for every element $\omega \in \mathcal{P}$ included in V , and (iii) H_t is a local diffeomorphism at every point z belonging to some $\omega \in \Sigma$.

(i) Since $r_t, t > 0$, is C^∞ and h is fixed on $V \setminus U$, it follows that H_t is C^∞ as well.

(ii) Let $\omega \subset V$ be an element of \mathcal{P} (resp. Σ).

If $\omega \subset V \setminus r_t(V)$, then H_t is fixed on ω , whence $H_t(\omega) = \omega \in \mathcal{P}$ (resp. Σ).

Suppose that $\omega \subset r_t(V)$. Since $r_t(V)$ is Θ -invariant, $\omega = r_t(\omega')$ for some another element $\omega' \in \mathcal{P}$ (resp. Σ). Then $h(\omega') \subset \omega'$, whence

$$H_t(\omega) = r_t \circ h \circ r_t^{-1}(\omega) = r_t \circ h(\omega') \subset r_t(\omega') = \omega.$$

(iii) Suppose that $\omega \in \Sigma$ and let $x \in \omega$.

If $x \in V \setminus r_t(U)$, then H_t is fixed in a neighbourhood of x , and therefore it is a local diffeomorphism at x .

Suppose that $x = r_t(x') \in r_t(U)$ for some $x' \in U$ and let $\omega' \in \Sigma$ be the element containing x' . Then h is a local diffeomorphism at x' , whence $H_t = r_t \circ h \circ r_t^{-1}$ is a local diffeomorphism at x . \square

5. STABILIZERS OF SMOOTH FUNCTIONS

Let $B^m \subset \mathbb{R}^m$ be the unit disk centered at the origin 0, $S^{m-1} = \partial B^m$ be its boundary sphere, $f : B^m \rightarrow \mathbb{R}$ be a C^∞ function, and Θ_f be the singular partition of f .

Let $\mathcal{S}(f) = \{h \in \mathcal{D}(B^m) : f \circ h = f\}$ be the stabilizer of f with respect to the right action of the group $\mathcal{D}(B^m)$ of diffeomorphisms of B^m on the space $C^\infty(B^m, \mathbb{R})$. Denote by $\mathcal{S}^+(f)$ the subgroup of $\mathcal{S}(f)$ consisting of orientation preserving diffeomorphisms. For $r \in \overline{\mathbb{N}}_0$ let also $\mathcal{S}_{\text{id}}(f)^r$ be the identity component of $\mathcal{S}(f)$ with respect to the C^r -topology. Then

$$\mathcal{S}_{\text{id}}(f)^\infty \subset \cdots \subset \mathcal{S}_{\text{id}}(f)^{r+1} \subset \mathcal{S}_{\text{id}}(f)^r \subset \cdots \subset \mathcal{S}_{\text{id}}(f)^0 \subset \mathcal{S}^+(f).$$

Theorem 5.1. *Let $m \geq 2$, $f : B^m \rightarrow [0, 1]$ be a C^∞ function such that 0 is a unique critical point of f being its global minimum, $f(0) = 0$, and $f(S^{m-1}) = 1$. Denote by \mathcal{S} the subgroup of $\mathcal{S}(f)$ consisting of diffeomorphisms h such that $h|_{S^{m-1}} : S^{m-1} \rightarrow S^{m-1}$ is C^∞ -isotopic to $\text{id}_{S^{m-1}}$. Suppose also that the singular partition Θ_f of f has an invariant ∞ -contraction to 0. Then $\mathcal{S} \subset \mathcal{S}_{\text{id}}(f)^0$.*

If $m = 2, 3, 4$, then $\mathcal{S} = \mathcal{S}_{\text{id}}(f)^0 = \mathcal{S}^+(f)$.

For the proof we need the following two simple standard statements concerning smoothing homotopies at the beginning and at the end, see e.g. [20, pp. 74 & 118] and [16, p. 205]. Let M be a closed smooth manifold.

Claim 5.2. *Let $a, b, c \in \mathbb{R}$ be numbers such that $0 < a < b < c$, and $N = M \times (0, c]$. Then we have a foliation on N by submanifolds $M \times t$, $t \in (0, c]$. Let also $h : N \rightarrow N$ be a C^∞ leaf preserving diffeomorphism, i.e., $h(x, t) = (\phi(x, t), t)$ for some C^∞ map $\phi : M \times (0, c] \rightarrow M$ such that for every $t \in (0, c]$ the map $\phi_t : M \rightarrow M$ is a diffeomorphism.*

Then there exists a leaf preserving isotopy relatively to $M \times (0, a]$ of h to a diffeomorphism $\hat{h}(x, t) = (\hat{\phi}(x, t), t)$ such that $\hat{\phi}_t = \phi_c$ for all $t \in [b, c]$.

Proof. Let $\mu : (0, c] \rightarrow (0, c]$ be a C^∞ function such that $\mu(t) = t$ for $t \in (0, a]$ and $\mu(t) = c$ for $t \in [b, c]$, see Figure 5.1a). Define the following leaf preserving isotopy $H : N \times I \rightarrow N$ by

$$H_s(x, t) = (\phi(x, (1 - s)t + s\mu(t)), t).$$

Then it easy to see that $H_0 = \text{id}_N$, $H_t = h$ on $M \times (0, a]$ and $\hat{h} = H_1$ satisfies conditions of our claim. \square

Claim 5.3. Let $d < e \in (0, 1)$ and $G : M \times I \rightarrow M$ be a C^∞ homotopy (isotopy). Then there exists another C^∞ homotopy (isotopy) $G' : M \times I \rightarrow M$ such that $\hat{G}_t = G_0$ for $t \in [0, d]$ and $\hat{G}_t = G_1$ for $t \in [e, 1]$.

Proof. Take any C^∞ function $\nu : I \rightarrow I$ such that $\nu[0, d] = 0$ and $\nu[e, 1] = 1$, and put $\hat{G}_t = G_{\nu(t)}$, see Figure 5.1b). \square




FIGURE 5.1

Proof of Theorem 5.1. Since 0 is a unique critical point of f and f is constant on S^{m-1} , it follows from the arguments of the proof of Lemma 4.4 that there exists a diffeomorphism $\eta : D^m \setminus 0 \rightarrow S^{m-1} \times (0, 1]$ such that $f \circ \eta^{-1}(y, t) = t$. In particular, for every $t \in [0, 1]$ the set $f^{-1}(t)$ is diffeomorphic with S^{m-1} . Since S^{m-1} is connected, we obtain that $\mathcal{D}(\Theta_f) = \mathcal{S}(f)$.

Therefore $\mathcal{D}_{\text{id}}(\Theta_f)^r = \mathcal{S}_{\text{id}}(f)^r$ for all $r \in \overline{\mathbb{N}}_0$.

By assumption there exists an invariant ∞ -contraction of Θ_f to 0 defined on some Θ_f -invariant neighbourhood V of 0. Therefore we can assume that $V = f^{-1}[0, 2c]$ for some $c \in (0, \frac{1}{2})$.

Lemma 5.4. *There exists a C^∞ -isotopy of h in \mathcal{S} to a diffeomorphism \hat{h} fixed on $f^{-1}[c, 1]$. Then it follows from Theorem 4.7 that \hat{h} and therefore h belong to $\mathcal{D}_{\text{id}}(\Theta_f)^0 = \mathcal{S}_{\text{id}}(f)^0$.*

Proof. Since h preserves f , it follows that the following diffeomorphism

$$g = \eta \circ h \circ \eta^{-1} : S^{m-1} \times (0, 1] \rightarrow S^{m-1} \times (0, 1]$$

is leaf preserving, i.e., $h(S^{m-1} \times t) = S^{m-1} \times t$ for all $t \in (0, 1]$. Then by Claim 5.2 we can assume that $g|_{S^{m-1} \times t} = h|_{S^{m-1}}$ for all $t \in [0.5, 1]$.

Take any $a \in (0, c)$. It suffices to find a leaf preserving isotopy relatively to $S^{m-1} \times (0, a]$ of g to a diffeomorphism \hat{g} fixed on $S^{m-1} \times [c, 1]$. This isotopy will yield an isotopy relatively to $f^{-1}[0, a]$ of h in \mathcal{S} to a diffeomorphism \hat{h} which is fixed on $f^{-1}[c, 1]$.

By assumptions of our theorem there exists a C^∞ isotopy $G : S^{m-1} \times [1, 2] \rightarrow S^{m-1}$ such that $G_1 = h|_{S^{m-1}}$ and $G_2 = \text{id}_{S^{m-1}}$. By Claim 5.3 we can assume that $G_t = h|_{S^{m-1}}$

for all $t \in [1, 1.5]$. Hence g and G yield the following C^∞ leaf preserving diffeomorphism:

$$T : S^{m-1} \times (0, 2] \rightarrow S^{m-1} \times (0, 2], \quad T(y, s) = \begin{cases} g(y, s), & s \in (0, 1], \\ G(y, s), & s \in [1, 2]. \end{cases}$$

Notice that $T(y, 2) = G(y, 2) = y$. Then, by Claim 5.3, T is isotopic via a leaf preserving isotopy relatively $S^{m-1} \times (0, a]$ to a diffeomorphism \hat{T} which is fixed on $S^{m-1} \times (c, 2]$. Denote $\hat{g} = \hat{T}|_{S^{m-1} \times (0, 1]}$. The restriction of this isotopy to $S^{m-1} \times (0, 1]$ gives a leaf preserving isotopy relatively $S^{m-1} \times (0, a]$ of g to a diffeomorphism \hat{g} with desired properties. The construction of homotopy is schematically presented in Figure 5.2. \square




FIGURE 5.2

Suppose now that $m = 2, 3, 4$. Then every orientation preserving diffeomorphism of S^{m-1} is C^∞ -isotopic to $\text{id}_{S^{m-1}}$, whence $\mathcal{S} = \mathcal{S}^+(f)$, and therefore $\mathcal{S} = \mathcal{S}_{\text{id}}(f)^0 = \mathcal{S}^+(f)$. For $m = 2$ this is rather trivial, for $m = 3$ is proved by S. Smale [22], and for $m = 4$ by A. Hatcher [4].

If $m \geq 5$, then $\mathcal{D}^+(S^{m-1})$ is not connected in general, see e.g. [17], and therefore Theorem 4.7 is not applicable. \square

6. LINEAR SYMMETRIES OF HOMOGENEOUS POLYNOMIALS

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a homogeneous polynomial of degree $p \geq 2$ given by (1.1),

$$f(x, y) = \pm \prod_{i=1}^l L_i^{\alpha_i}(x, y) \cdot \prod_{j=1}^k Q_j^{\beta_j}(x, y).$$

Denote

$$\mathcal{LS}(f) = \mathcal{S}^+(f) \cap \text{GL}^+(2, \mathbb{R}).$$

Thus $\mathcal{LS}(f)$ consists of preserving orientation linear automorphisms $h : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $f \circ h = f$. Also notice that $\mathcal{LS}(f)$ is a closed subgroup of $\text{GL}^+(2, \mathbb{R})$, and therefore it is a Lie group. Denote by $\mathcal{LS}(f)_0$ the connected component of the unit matrix $\text{id}_{\mathbb{R}^2}$ in $\mathcal{LS}(f)$.

In this section we recall the structure of $\mathcal{LS}(f)$. Notice that we may make linear changes of coordinates to reduce f to a convenient form. Then $\mathcal{LS}(f)$ will change to a conjugate subgroup in $\text{GL}^+(2, \mathbb{R})$.

The following statement is evident.

Lemma 6.1. *If $\deg f$ is even, then $f(-z) \equiv f(z)$, i.e., $-\text{id}_{\mathbb{R}^2} \in \mathcal{LS}(f)$. Therefore in this case $\mathcal{LS}(f)$ is a non-trivial group.*

We will distinguish the following five cases of f .

(A) $l = 1, k = 0, f = L_1^{\alpha_1}$. By a linear change of coordinates we can assume that $L_1(x, y) = y$ and thus $f(x, y) = y^{\alpha_1}$. Then

$$\mathcal{LS}_0(y^{\alpha_1}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} : a > 0 \right\}.$$

If α_1 is odd then $\mathcal{LS}_0 = \mathcal{LS}(f)$, otherwise, $\mathcal{LS}(f)$ consists of two connected components \mathcal{LS}_0 and $-\mathcal{LS}_0$.

(B) $l = 2, k = 0, f = L_1^{\alpha_1} L_2^{\alpha_2}$. By a linear change of coordinates we can assume that $L_1(x, y) = x, L_2(x, y) = y$ and thus $f(x, y) = x^{\alpha_1} y^{\alpha_2}$. Then

$$\mathcal{LS}_0(x^{\alpha_1} y^{\alpha_2}) = \left\{ \begin{pmatrix} e^{\alpha_2 t} & 0 \\ 0 & e^{-\alpha_1 t} \end{pmatrix} : t \in \mathbb{R} \right\}.$$

Moreover, $\mathcal{LS}(f)/\mathcal{LS}_0$ is isomorphic with some subgroup of \mathbb{Z}_4 generated by the rotation of \mathbb{R}^2 by $\pi/2$.

(C) $l = 0, k = 1, f = Q_1^{\beta_1}$. By a linear change of coordinates we can assume that $Q_1(x, y) = x^2 + y^2$, whence $f(x, y) = (x^2 + y^2)^{\beta_1}$. Then

$$\mathcal{LS}(x^2 + y^2) = SO(2, \mathbb{R}) = \left\{ \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} : t \in [0, 2\pi) \right\}.$$

The above statements are elementary and we left them to the reader. Notice also that, in the cases (A)–(C), $l + 2k \leq 2$. The remaining two cases are the following:

(D) $l = 0, k \geq 2, f = Q_1^{\beta_1} \cdots Q_k^{\beta_k}$. In this case $\deg f$ is even, whence $\mathcal{LS}(f)$ is non-trivial.

(E) $l \geq 1, l + 2k \geq 3$.

Lemma 6.2. *In the cases (D) and (E), $\mathcal{LS}(f)$ is a finite cyclic subgroup of $GL^+(2, \mathbb{R})$. Moreover, in the case (E), $\mathcal{LS}(f)$ is a subgroup of \mathbb{Z}_{2l} .*

Proof. In fact cyclicity of $\mathcal{LS}(f)$ for the case $l + 2k - 1 \geq 2$ can be extracted from the paper of W. C. Huffman, [6], where the symmetries of complex binary forms are classified. Regard $f : \mathbb{C}^2 \rightarrow \mathbb{C}$ as a complex polynomial with real coefficients. Then by [6] the subgroup $\mathcal{LS}_{\mathbb{C}}(f)$ of $GL(2, \mathbb{C})$ consisting of *complex* symmetries of f turned out to be of one of the following types: cyclic, dihedral, tetrahedral, octahedral, and icosahedral. Notice $\mathcal{LS}(f)$ is the subgroup of $\mathcal{LS}_{\mathbb{C}}(f)$ consisting of *preserving orientation real* symmetries of f , i.e., automorphisms which also leave invariant the 2-plane $\mathbb{R}^2 \subset \mathbb{C}^2$ of real coordinates and preserve its orientation. Then it follows from the structure of symmetries of regular polyhedrons that $\mathcal{LS}(f)$ must be cyclic.

Nevertheless, since we need a very particular case of [6] and for the sake of completeness, we will present a short elementary proof. It suffices to show that $\mathcal{LS}(f)$ is finite, see 6.6. This will imply that $\mathcal{LS}(f)$ is isomorphic to a finite subgroup of $SO(2)$, and therefore is cyclic. Also notice that the fact that $\mathcal{LS}(f)$ is discrete also follows from [12]. First we establish the following three statements.

Claim 6.3. *Let $h \in GL^+(2, \mathbb{R})$ and Q be a positive definite quadratic form such that $Q \circ h = tQ$ for some $t > 0$. Then $t = \det(h)$.*

Proof. By a linear change of coordinates we can assume that $Q(z) = |z|^2$. Then $h(z) = \sqrt{t}e^{i\psi}z$ for some $\psi \in \mathbb{R}$, hence $\det(h) = t$. □

Claim 6.4. *Let Q_1, Q_2 be a positive definite quadratic form such that $\frac{Q_1}{Q_2} \neq \text{const}$. Let also $h \in GL^+(2, \mathbb{R})$ be such that $Q_i \circ h = tQ_i$ for $i = 1, 2$, where $t = \det(h)$. Then $h(z) = \pm\sqrt{t}z$.*

Proof. We can assume that $Q_1(x, y) = x^2 + y^2$ and $Q_2(x, y) = ax^2 + by^2$, where $a, b > 0$ and either $a \neq 1$ or $b \neq 1$. Denote $g(z) = h(z)/\sqrt{t}$. Then $Q_i \circ g = Q_i$, i.e., g preserves every circle $x^2 + y^2 = \text{const}$ and every ellipse $ax^2 + by^2 = \text{const}$. Therefore $g = \pm \text{id}_{\mathbb{R}^2}$, and $h(z) = \pm\sqrt{t}z$. □

Claim 6.5. *If $t \cdot \text{id}_{\mathbb{R}^2} \in \mathcal{LS}(f)$ for some $t \in \mathbb{R}$, then $t = \pm 1$.*

Proof. Let $z \in \mathbb{R}^2$ be such that $f(z) \neq 0$. Since f is homogeneous, we have $f(z) = f(tz) = t^{\deg f} f(z)$, whence $t = \pm 1$. □

Let $h \in \mathcal{LS}(f)$. Since L_i and Q_j are irreducible over \mathbb{R} , so are $L_i \circ h$ and $Q_j \circ h$. Therefore the identity $f \circ h = f$ implies that “ h permutes L_i and Q_j up to non-zero multiples”. This means that for every i there exist i' and $s_i \in \mathbb{R} \setminus \{0\}$, and for every j there exist j' and $t_j > 0$ such that

$$L_i(h(z)) = s_i L_{i'}(z), \quad Q_j(h(z)) = t_j Q_{j'}(z).$$

Denote by $\text{Sym}(r)$ the group of permutations of r symbols. Then we have a well-defined homomorphism

$$\mu : \mathcal{LS}(f) \rightarrow \text{Sym}(l) \times \text{Sym}(k)$$

associating to every $h \in \mathcal{LS}$ its permutations of L_i and Q_j .

Claim 6.6. *If $l + 2k \geq 3$, then $\ker \mu \subset \{\pm \text{id}_{\mathbb{R}^2}\}$, whence $\mathcal{LS}(f)$ is a finite group.*

Proof. Let $h \in \ker \mu$. Thus $L_i \circ h = s_i L_i$ and $Q_j \circ h = t_j Q_j$ for all i, j . We will show that $h = t \cdot \text{id}_{\mathbb{R}^2}$ for some $t \neq 0$. Then it will follow from Claim 6.5 that $h = \pm \text{id}$.

Notice that h preserves every line $\{L_i = 0\}$ and thus has l distinct eigen directions.

a) Therefore if $l \geq 3$, then $h = t \cdot \text{id}_{\mathbb{R}^2}$ for some $t \in \mathbb{R}$.

b) Moreover, if $k \geq 2$, then by Claim 6.4, we also have $h = \pm \text{id}_{\mathbb{R}^2}$.

c) Suppose that $1 \leq l \leq 2$ and $k = 1$. We can assume that $Q(z) = |z|^2$ and that $h(z) = t e^{i\psi} z$ for some $t > 0$ and $\psi \in \mathbb{R}$. Since h has $l \geq 1$ eigen directions, we obtain that $h(z) = \pm \sqrt{t} z$. □

Thus $\mathcal{LS}(f) \approx \mathbb{Z}_n$ for some $n \in \mathbb{N}$. Let h be a generator of $\mathcal{LS}(f)$. Then we can assume that $h(z) = e^{2\pi i/n} z$. It remains to prove the latter statement.

Claim 6.7. *Suppose that $l \geq 1$. Then n divides $2l$, whence $\mathcal{LS}(f)$ is isomorphic to a subgroup of \mathbb{Z}_{2l} .*

Proof. Since $f \circ h = f$, it follows that $h(f^{-1}(0)) = f^{-1}(0)$. By assumption $l \geq 1$, whence $f^{-1}(0) = \bigcup_{i=1}^l \{L_i = 0\}$ is the union of l lines passing through the origin. This set can be viewed as the union of $2l$ rays starting at the origin, and these rays are cyclically shifted by h . Moreover, if h^t preserves at least one of these rays, then $h^t = \text{id}_{\mathbb{R}^2}$. Therefore n divides $2l$. □

Lemma 6.2 is completed. □

7. PROOF OF THEOREM 1.1.

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a homogeneous polynomial of degree $p \geq 1$ given by (1.1)

$$f(x, y) = \pm \prod_{i=1}^l L_i^{\alpha_i}(x, y) \cdot \prod_{j=1}^k Q_j^{\beta_j}(x, y).$$

We will refer to the cases (A)–(E) of f considered in the previous section. We have to show that $\mathcal{S}_{\text{id}}(f)^\infty = \dots = \mathcal{S}_{\text{id}}(f)^1$ and that $\mathcal{S}_{\text{id}}(f)^1 \neq \mathcal{S}_{\text{id}}(f)^0$ iff f is of the case (D).

Our first aim is to identify $\mathcal{S}_{\text{id}}(f)^r$ with the group $\mathcal{D}_{\text{id}}(\Theta_G)^r$ for some vector field G on \mathbb{R}^2 , see Lemma 7.1. Then we will use the shift map of G . Denote

$$D = \pm \prod_{i=1}^l L_i^{\alpha_i - 1} \cdot \prod_{j=1}^k Q_j^{\beta_j - 1}.$$

Then

$$f = L_1 \cdots L_l \cdot Q_1 \cdots Q_k \cdot D$$

and it is easy to see that D is the greatest common divisor of f'_x and f'_y in the ring $\mathbb{R}[x, y]$.

Let $F = -f'_y \frac{\partial}{\partial x} + f'_x \frac{\partial}{\partial y}$ be the Hamiltonian vector field of f on \mathbb{R}^2 and

$$G = F/D = -(f'_y/D) \frac{\partial}{\partial x} + (f'_x/D) \frac{\partial}{\partial y}.$$

We will call G the *reduced* Hamiltonian vector field of f . Notice that

$$\deg G = l + 2k - 1$$

and the coordinate functions of G are relatively prime in $\mathbb{R}[x, y]$.

As noted in Example 3.3 the singular partitions Θ_f and Θ_F coincide. Let us describe the singular partition $\Theta_G = (\mathcal{P}_G, \Sigma_G)$. Recall that elements of \mathcal{P}_G are the orbits of G and Σ_G consists of zeros of G . Consider the following cases, see Figure 7.1.

(A) $f = y^{\alpha_1}$. Then $D = y^{\alpha_1 - 1}$ and $F(x, y) = \alpha_1 y^{\alpha_1 - 1} \frac{\partial}{\partial y}$. Hence $G(x, y) = \alpha_1 \frac{\partial}{\partial y}$ is a constant vector field and the partition Θ_G consists of horizontal lines $\{y = \text{const}\}$ being non-singular elements of Θ_G .

(C) and (D) $f = Q_1^{\beta_1} \cdots Q_k^{\beta_k}$. In this case $\Theta_F = \Theta_G$. The origin is a unique singular element of Θ_G . All other elements of Θ_G are level-sets $f^{-1}(c)$ of f for $c > 0$.

(B) and (E) either $l = 2$ and $k = 0$ or $l \geq 1$ and $k \geq 1$. In both cases the set of singular points of Θ_F consist of the origin and the set

$$D^{-1}(0) = \bigcup_{i : \alpha_i \geq 2} \{L_i = 0\}$$

of zeros of D being the union of those lines $\{L_i = 0\}$ for which L_i is a multiple factor of f . Since after division of F by D the coordinate functions of $G = F/D$ are relatively prime, it follows that 0 is a unique singular element of Θ_G . Hence non-singular elements of Θ_G are the connected components $f^{-1}(c)$ for $c \neq 0$ and the half-lines in $f^{-1}(0) \setminus 0$, see Figure 7.2.




FIGURE 7.1




FIGURE 7.2. Case (B). Hamiltonian and reduced Hamiltonian vector fields for $f(x, y) = x y^2$.

Since f is constant along orbits of F and G , it follows that

$$(7.1) \quad \mathcal{D}(\Theta_F) \subset \mathcal{D}(\Theta_G) \subset \mathcal{S}(f).$$

Also notice that $\mathcal{D}(\Theta_F)$ consists of those $h \in \mathcal{D}(\Theta_G)$ which fixes every critical point of f .

Lemma 7.1. $\mathcal{D}_{\text{id}}(\Theta_G)^r = \mathcal{S}_{\text{id}}(f)^r$ for all $r \in \overline{\mathbb{N}}_0$.

Proof. It follows from (7.1) that $\mathcal{D}_{\text{id}}(\Theta_G)^r \subset \mathcal{S}_{\text{id}}(f)^r$.

Conversely, let $h \in \mathcal{S}_{\text{id}}(f)^r$, so there exists an r -isotopy $h_t : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ between $h_0 = \text{id}_{\mathbb{R}^2}$ and $h_1 = h$ in $\mathcal{S}(f)$, i.e.,

$$(7.2) \quad f \circ h_t = f, \quad t \in I.$$

We claim that every $h_t \in \mathcal{D}(\Theta_G)$, i.e., $h_t(\omega) = \omega$ for every element ω of Θ_G . This will mean that $\{h_t\}$ is an r -isotopy in $\mathcal{D}(\Theta_G)$, whence $h \in \mathcal{D}_{\text{id}}(\Theta_G)^r$.

It follows from (7.2) that $h_t(f^{-1}(c)) = f^{-1}(c)$ for every $c \in \mathbb{R}$ and $h_t(\Sigma_f) = \Sigma_f$. Since $h_0 = \text{id}_{\mathbb{R}^2}$ preserves every connected component ω of $f^{-1}(c) \setminus \Sigma_f$, so does h_t , $t \in I$. If either $c \neq 0$, or $c = 0$ but f is of either the cases (A), (C), or (D), then by definition every such ω is an element of Θ_G .

Let $c = 0$. We claim that in the cases (B) and (E) $h_t(0) = 0$ for all $t \in I$. Indeed, in these cases the origin is “the most degenerate point among all other points of $f^{-1}(0)$ ”. This means the following.

For every $z \in f^{-1}(0)$ denote by p_z the least number such that p_z -jet of f at z does not vanish, i.e., $j^{p_z-1}(f, z) = 0$ while $j^{p_z}(f, z) \neq 0$. In other words, the Taylor series of f at z starts with terms of order p_z . It is easy to see that for the origin $p_0 = \deg f$, while for all other points $z \in f^{-1}(0)$ we have that $p_z < \deg f$. Also notice that this number p_z is preserved by any diffeomorphism $h \in \mathcal{S}(f)$, i.e., $p_{h(z)} = p_z$. It follows that $h_t(0) = 0$.

It remains to note that by continuity every h_t preserves connected components of $D^{-1}(0) \setminus \{0\}$. Hence $h_t \in \mathcal{D}(\Theta_G)$ for all $t \in I$. \square

Now we can complete Theorem 1.1. Let Φ be the local flow on \mathbb{R}^2 generated by G , and φ be the shift map of G , see Section 3.5. The following statement was established in [9].

Lemma 7.2. *In the cases (A)-(C) (i.e., when $\deg G \leq 1$) for every $h \in \mathcal{E}_{\text{id}}(\Theta_G)^0$ there exists a smooth function $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $h(z) = \Phi(x, \sigma(x))$, i.e., $\text{im}(\varphi) = \mathcal{E}_{\text{id}}(\Theta_G)^0$.*

It follows from Lemmas 3.6 and 7.1 that, in the cases (A)-(C),

$$(7.3) \quad \mathcal{S}_{\text{id}}(f)^\infty = \mathcal{D}_{\text{id}}(\Theta_G)^\infty = \dots = \mathcal{D}_{\text{id}}(\Theta_G)^0 = \mathcal{S}_{\text{id}}(f)^0.$$

The following lemma is a consequence of results of [12, 13].

Lemma 7.3. *In the cases (D) and (E) (i.e., when $\deg G \geq 2$) $\text{im}(\varphi)$ consists of all $h \in \mathcal{E}(\Theta_G)$ whose tangent map $T_0h : T_0\mathbb{R}^2 \rightarrow T_0\mathbb{R}^2$ at 0 is the identity.*

Corollary 7.4. $\mathcal{E}_{\text{id}}(\Theta_G)^1 \subset \text{im}(\varphi)$, whence similarly to (7.3) we get $\mathcal{S}_{\text{id}}(f)^\infty = \dots = \mathcal{S}_{\text{id}}(f)^1$.

Proof of Corollary. Let $h \in \mathcal{E}_{\text{id}}(\Theta_G)^1$. Then there exists a 1-homotopy h_t between $h_0 = \text{id}_{\mathbb{R}^2}$ and $h_1 = h$ in $\mathcal{E}(\Theta_G)$. In particular, T_0h_t is continuous in t .

Since f is homogeneous, it follows from [9, Lemma 36], that T_0h_t regarded as a linear automorphism of \mathbb{R}^2 also must preserve f , i.e., $T_0h_t \in \mathcal{LS}(f)$. Therefore the family of maps T_0h_t can be regarded as a homotopy in $\mathcal{LS}(f)$. But by Lemma 6.2 in the cases (D) and (E) the group $\mathcal{LS}(f)$ is discrete, whence all T_0h_t coincide with the identity map $T_0h_0 = \text{id}_{\mathbb{R}^2}$. In particular, $T_0h = \text{id}_{\mathbb{R}^2}$, whence by Lemma 7.3 $h \in \text{im}(\varphi)$. \square

It remains to show that $\mathcal{S}_{\text{id}}(f)^1$ and $\mathcal{S}_{\text{id}}(f)^0$ coincide in the case (E) and are distinct in the case (D).

(D) Let f be a product of at least two distinct definite quadratic forms. Then by Theorem 5.1 $\mathcal{S}_{\text{id}}(f)^0 = \mathcal{S}^+(f)$. Moreover, since $\deg f$ is even, we have that $-\text{id}_{\mathbb{R}^2} \in \mathcal{S}^+(f) = \mathcal{S}_{\text{id}}(f)^0$, see Lemma 6.1. On the other hand by Lemma 7.3 for every $h \in \mathcal{S}_{\text{id}}(f)^1$ its tangent map $T_0h = \text{id}_{\mathbb{R}^2} \neq -\text{id}_{\mathbb{R}^2}$. Hence $\mathcal{S}_{\text{id}}(f)^1 \neq \mathcal{S}_{\text{id}}(f)^0$.

(E) In this case $f^{-1}(0)$ is a union of $l \geq 1$ straight lines $\{L_i = 0\}$ passing through the origin. Let $h \in \mathcal{D}_{\text{id}}(\Theta_G)^0$. Since there exists a homotopy between h and $\text{id}_{\mathbb{R}^2}$ in

$\mathcal{D}_{\text{id}}(\Theta_G)^0$, it follows that h preserves every half-line of $f^{-1}(0) \setminus \{0\}$. Therefore so does $T_0h \in \mathcal{LS}(f)$. Then it follows from Claim 6.7 that $T_0h = \text{id}_{\mathbb{R}^2}$, whence by Lemma 7.3 $h \in \text{im}(\varphi)$. Thus $\mathcal{D}_{\text{id}}(\Theta_G)^0 \subset \text{im}(\varphi)$.

Now we get from Lemma 3.6 that

$$\mathcal{S}_{\text{id}}(f)^1 = \mathcal{D}_{\text{id}}(\Theta_G)^1 = \mathcal{D}_{\text{id}}(\Theta_G)^0 = \mathcal{S}_{\text{id}}(f)^0.$$

Theorem 1.1 is completed.

Acknowledgments. I would like to thank V. V. Sharko and E. Polulyakh for useful discussions. I also thank anonymous referee for careful reading of this manuscript and critical remarks which allowed for clarification of the present exposition.

REFERENCES

1. V. I. Arnold, A. N. Varchenko, S. M. Gusein-Zade, *Singularities of Differentiable Mappings. I*, Nauka, Moscow, 1982. (Russian)
2. E. N. Dancer, *Degenerate critical points, homotopy indices and Morse inequalities. II*, J. Reine Angew. Math. **382** (1987), 145–164.
3. M. H. Freedman, *The topology of four-dimensional manifolds*, J. Differential Geom. **17** (1982), no. 3, 357–453.
4. A. Hatcher, *A proof of Smale Conjecture* $\text{Diff}(S^3) \cong O(4)$, Ann. of Math. **117** (1983), 533–607.
5. M. W. Hirsch, *Differential Topology*, Series: Graduate Texts in Mathematics, **33**, Springer-Verlag, 1976.
6. W. C. Huffman, *Polynomial invariants of finite linear groups of degree two*, Canad. J. Math. **32** (1980), 317–330.
7. K. Kuratowski, *Topology. II*, Academic Press, New York—London, 1968.
8. J. A. Leslie, *On a differential structure for the group of diffeomorphisms*, Topology **6** (1967), no. 4, 263–271.
9. S. Maksymenko, *Smooth shifts along trajectories of flows*, Topology Appl. **130** (2003), no. 2, 183–204; <http://lanl.arxiv.org/abs/math/0106199>.
10. S. Maksymenko, *Stabilizers and orbits of smooth functions*, Bulletin des Sciences Mathématiques **130** (2006), 279–311; <http://xxx.lanl.gov/abs/math/0411612>.
11. S. Maksymenko, *Homotopy types of stabilizers and orbits of Morse functions on surfaces*, Annals of Global Analysis and Geometry **29** (2006), no. 3, 241–285; <http://xxx.lanl.gov/abs/math/0310067>.
12. S. Maksymenko, *∞ -jets of diffeomorphisms preserving orbits of vector fields*, Central European Journal Math. **7** (2009), no. 2, 272–298; <http://xxx.lanl.gov/abs/math/0708.0737>.
13. S. Maksymenko, *Hamiltonian vector fields of homogeneous polynomials on the plane*, Problems of topology and related questions, Proceedings of Institute of Mathematics of NAS of Ukraine, vol. 3, no. 3, 2006, pp. 269–308. (Ukrainian); <http://xxx.lanl.gov/abs/math/0709.2511>.
14. J. Milnor, *Lectures on the h-Cobordism Theorem*, Princeton, New Jersey, 1965.
15. J. Milnor, *Morse Theory*, Princeton, New Jersey, 1963.
16. J. Milnor, A. Wallace, *Differential Topology*, Mir, Moscow, 1972. (Russian)
17. S. P. Novikov, *Differentiable sphere bundles*, Izv. Acad. Nauk SSSR **29** (1965), 71–96.
18. G. Perelman, *The entropy formula for the Ricci flow and its geometric application*, 11 Nov 2002, <http://arxiv.org/abs/math.DG/0211159>.
19. G. Perelman, *Ricci flow with surgery on three-manifolds*, 10 Mar 2003, <http://arxiv.org/abs/math.DG/0303109>.
20. L. S. Pontryagin, *Smooth Manifolds and Their Applications in Homotopy Theory*, Nauka, Moscow, 1985. (Russian)
21. K. Saito, *Quasihomogene isolierte Singularitäten von Hyperflächen*, Inv. Math. **14** (1971), 123–142.
22. S. Smale, *Diffeomorphisms of the 2-sphere*, Proc. Amer. Math. Soc. **10** (1959), 621–626.
23. S. Smale, *Generalized Poincaré’s conjecture in dimensions greater than four*, Ann. of Math. **74** (1961), 391–406.

TOPOLOGY DEPARTMENT, INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE,
3 TERESHCHENKIVS’KA, KYIV, 01601, UKRAINE
E-mail address: maks@imath.kiev.ua

Received 10/06/2008; Revised 01/12/2008