

## ON \*-REPRESENTATIONS OF THE PERTURBATION OF TWISTED CCR

KONSTANTIN M. SUKRETNÝĚ

ABSTRACT. A classification of irreducible \*-representations of a certain deformation of twisted canonical commutation relations is given.

### 1. INTRODUCTION

In this note we study representations of a \*-algebra  $A_\alpha$  defined by generators  $a_i, a_i^*$ ,  $i = 1, \dots, d$ , satisfying the commutation relations of the following form:

$$(1) \quad \begin{aligned} a_i^* a_i &= 1 + \alpha_i^2 a_i a_i^* - \sum_{j < i} (1 - \alpha_j^2) a_j a_j^*, \\ a_i^* a_j &= \alpha_i a_j a_i^*, \quad i < j, \\ a_j a_i &= \alpha_i a_i a_j, \quad i, j = 1, \dots, d, \quad i < j, \end{aligned}$$

where we additionally suppose that  $\alpha_i^2 = \mu^{n_i}$ ,  $0 < \mu < 1$ ,  $n_i \in \mathbb{N}$ ,  $i = 1, \dots, d$ . When  $n_i = 2$ ,  $i = 1, \dots, d$ , we get the twisted canonical commutation relations (TCCR) constructed and studied by W. Pusz and S. L. Woronowicz, see [5]. These relations also belong to the class of generalized canonical commutation relations (GCCR), defined in [3].

The aim of this paper is to study irreducible representations of  $A_\alpha$  by, possibly unbounded, Hilbert space operators. Note that representations of TCCR were classified in [5]. The description of bounded representations of GCCR was obtained in [3]. In [2] the authors proved that the Fock representation of the universal enveloping  $C^*$ -algebra generated by GCCR is faithful.

To deal with the unbounded representations one has firstly to give a precise definition of a family of unbounded operators satisfying relations (1). To do so, let us perform some formal manipulations with generators and relations.

Construct the polar decompositions of  $a_i^*$ ,  $a_i^* = U_i C_i$ , where  $C_i^2 = a_i a_i^*$ ,  $U_i$  is a partial isometry and  $\ker U_i = \ker C_i = \ker a_i^*$ . Then the commutation relations (1) take the following form:

$$(2) \quad \begin{aligned} C_i^2 U_i^* &= U_i^* \left( 1 + \alpha_i^2 C_i^2 - \sum_{j < i} (1 - \alpha_j^2) C_j^2 \right), \\ C_i^2 U_j^* &= \alpha_j^2 U_j^* C_i^2, \quad j < i, \\ C_i^2 U_j^* &= U_j^* C_i^2, \quad j > i, \end{aligned}$$

$$(3) \quad C_i C_j = C_j C_i, \quad U_j U_i = U_i U_j, \quad U_j^* U_i = U_i U_j^*, \quad i \neq j.$$

---

2000 *Mathematics Subject Classification*. Primary 47L60, 47L30, 47A67; Secondary 81R10.  
*Key words and phrases*. Deformed commutation relations, irreducible \*-representations.

Consider the functions

$$F_j(x_1, \dots, x_d) = \left( x_1, \dots, x_{j-1}, 1 + \alpha_j^2 x_j - \sum_{k < j} (1 - \alpha_k^2) x_k, \right. \\ \left. \alpha_j^2 x_{j+1}, \dots, \alpha_j^2 x_d \right), \quad j = 1, \dots, d.$$

Then, in a compact form, relations (2) can be written as follows:

$$(C_1^2, \dots, C_d^2) U_j^* = U_j^* F_j(C_1^2, \dots, C_d^2), \quad j = 1, \dots, d.$$

Note that, in the bounded case, the relations (1) and (2, 3) are equivalent.

**Definition 1.** (see [4]). *Let a family of self-adjoint operators  $\mathcal{C} = \{C_i^2, i = 1, \dots, d\}$  commute on a dense invariant domain of analytic vectors. We say that the family  $\mathcal{C}$  and partial isometries  $\{U_i, i = 1, \dots, d\}$  satisfy relations (2) if for any Borel set  $\Delta \subset \mathbb{R}^d$  and any  $j = 1, \dots, d$  one has*

$$E_{\mathcal{C}}(\Delta) U_j^* = U_j^* E_{\mathcal{C}}(F_j^{-1}(\Delta)),$$

where  $E_{\mathcal{C}}(\cdot)$  is the joint resolution of identity of the family  $\mathcal{C}$ .

**Definition 2.** *Let families  $\mathcal{C}$  and  $\{U_i, i = 1, \dots, d\}$  satisfy the conditions of the definition above and  $\ker U_i = \ker C_i, i = 1, \dots, d$ , then we say that the family of operators  $a_i^* = U_i C_i, i = 1, \dots, d$  is an unbounded representation of relations (1).*

## 2. REPRESENTATIONS OF $A_\alpha$

In this section we will use a dynamical system method developed in a series of papers by Yu. Samoilenko, V. Ostrovkyi, L. Turowska, E. Vaisleb et al., see [4] and the references therein.

Our considerations will be based on an analysis of the spectrum of  $C_1^2$  in the irreducible representation. Since

$$C_1^2 U_1^* = U_1^* (1 + \alpha_1^2 C_1^2), \quad C_1^2 U_j^* = U_j^* C_1^2, C_1 C_j = C_j C_1, \quad j \geq 2$$

in an irreducible representation of (2,3), the spectrum of  $C_1^2$  is coincides with the positive orbit of the dynamical system  $(f_1, \mathbb{R})$ , where  $f_1(t) = 1 + \alpha_1^2 t$ , see [4]. Such orbits can be subdivided onto the following three types:

- (1) Fock orbit,  $O_F = \left\{ \frac{1 - \alpha_1^{2n}}{1 - \alpha_1^2}, n \in \mathbb{Z}_+ \right\}$ ;
- (2) fixed point  $O_{fix} = \left\{ \frac{1}{1 - \alpha_1^2} \right\}$ ;
- (3) unbounded orbits, labeled by  $x_1 \in \tau_{y_1} = (1 + \alpha_1^2 y_1, y_1], y_1 > \frac{1}{1 - \alpha_1^2}$  is fixed,

$$O_{x_1} = \left\{ \frac{1 - \alpha_1^{2n}}{1 - \alpha_1^2} + \alpha_1^{2n} x_1, n \in \mathbb{Z} \right\}.$$

In the following propositions we give a description of irreducible representations of  $A_\alpha$  when the spectrum of  $C_1^2$  is assumed to coincide with one of the orbits above.

We start with the most simple case.

**Proposition 1.** *Let in irreducible representation of  $A_\alpha$  one has  $\sigma(C_1^2) = \overline{O_F}$ , then, up to a unitary equivalence,  $\mathcal{H} = l_2(\mathbb{Z}_+) \otimes \mathcal{K}$  and*

$$C_1^2 = d(f_1) \otimes \mathbf{1}, \quad U_1^* = S \otimes \mathbf{1}, \\ C_i^2 = d(\alpha_1^2) \otimes \widehat{C}_i^2, \quad U_i^* = \mathbf{1} \otimes \widehat{U}_i^*, \quad i = 2, \dots, d,$$

where, for the standard basis of  $l_2(\mathbb{Z}_+)$  denoted by  $\{e_n, n \in \mathbb{Z}_+\}$ , one has

$$d(f_1)e_n = f_1^n(0)e_n = \frac{1 - \alpha_1^{2n}}{1 - \alpha_1^2} e_n, \quad d(\alpha_1^2)e_n = \alpha_1^{2n} e_n, \quad n \in \mathbb{Z}_+$$

and the family of the operators  $\{\widehat{C}_i, \widehat{U}_i, i = 2, \dots, d\}$  is irreducible on  $\mathcal{K}$  and satisfies the relations (2,3) with  $d - 1$  generators.

*Proof.* The proof is analogous to the proof of the propositions below and the most trivial among them, so we omit it here.  $\square$

Let us now suppose that  $\sigma(C_1^2) = \{\frac{1}{1-\alpha_1^2}\}$  and  $d > 3$ . Fix  $y_2 > 0$ , put  $\sigma_{y_2} = (\mu^l y_2, y_2]$ , where  $l = \mathbf{GCD}(n_1, n_2)$ ,  $\alpha_i^2 = \mu^{n_i}$ ,  $i = 1, 2$ . Let also  $n_i = lk_i$ ,  $i = 1, 2$  and  $l = n_1 m_1 + n_2 m_2$ .

**Proposition 2.** *If  $d > 2$  and  $\sigma(C_1^2) = \{\frac{1}{1-\alpha_1^2}\}$  and  $C_2^2 \neq 0$  in the irreducible representation, then, up to a unitary equivalence,  $\mathcal{H} = l_2(\mathbb{Z}) \otimes \bigotimes_{i=3}^d l_2(\mathbb{Z}_+)$  and*

$$\begin{aligned}
 C_1^2 &= \frac{1}{1-\alpha_1^2} \mathbf{1} \otimes \bigotimes_{2 < k \leq d} \mathbf{1}, \\
 U_j^* &= e^{i\phi_j} E^{k_j} \otimes \bigotimes_{2 < k \leq d} \mathbf{1}, \quad j = 1, 2, \quad m_1 \phi_1 + m_2 \phi_2 = 0, \quad \text{mod } 2\pi, \\
 C_2^2 &= x_2 D(\mu^l) \otimes \bigotimes_{2 < k \leq d} \mathbf{1}, \quad x_2 \in \sigma_{y_2}, \\
 C_i^2 &= D(\mu^l) \otimes \bigotimes_{2 < k < i} \widehat{d}(\alpha_k^2) \otimes \widehat{d}(h_i(0, x_2)) \otimes \bigotimes_{i < k \leq d} \mathbf{1}, \quad i = 3, \dots, d, \\
 U_i^* &= \mathbf{1} \otimes \bigotimes_{2 < k < i} \mathbf{1} \otimes \widehat{S} \otimes \bigotimes_{i < k \leq d} \mathbf{1}, \quad i = 3, \dots, d,
 \end{aligned}$$

and

$$D(\mu^l), \quad E: l_2(\mathbb{Z}) \rightarrow l_2(\mathbb{Z}), \quad D(\mu^l)e_n = \mu^{nl}e_n, \quad Ee_n = e_{n+1}, \quad n \in \mathbb{Z},$$

where  $\{e_n, n \in \mathbb{Z}\}$  is the standard basis of  $l_2(\mathbb{Z})$ ;

$$h_i(t, x_2) = -(1 - \alpha_2^2)x_2 + \alpha_2^2 t,$$

$$\widehat{d}(h_i(0, x_2)), \quad \widehat{S}, \quad \widehat{d}(\lambda): l_2(\mathbb{Z}_-) \rightarrow l_2(\mathbb{Z}_-), \quad \widehat{S}e_0 = 0, \quad \widehat{S}e_{-n} = e_{-n+1}, \quad n \geq 1,$$

$$\widehat{d}(h_i(0, x_2))e_{-n} = h_i^{-n}(0, x_2)e_{-n}, \quad \widehat{d}(\lambda)e_{-n} = \lambda^{-n}e_{-n}, \quad n \in \mathbb{Z}_+,$$

where  $\{e_{-n}, n \in \mathbb{Z}_+\}$  is the standard basis of  $l_2(\mathbb{Z}_-)$ .

*Proof.* Since  $C_1^2 = \frac{1}{1-\alpha_1^2} \mathbf{1}$  and  $\ker U_1 = \ker U_1^* = \{0\}$ ,  $U_1$  is a unitary operator. Furthermore, one has

$$C_i^2 U_i^* = U_i^* \left( \alpha_i^2 C_i^2 - \sum_{1 < j < i} (1 - \alpha_j^2) C_j^2 \right), \quad i \geq 2.$$

In particular,  $C_2^2 U_i^* = \alpha_i^2 U_i^* C_2^2$ ,  $i = 1, 2$ . Since  $C_2^2 U_j = U_j C_2^2$ ,  $C_2^2 C_j^2 = C_j^2 C_2^2$ ,  $j > 2$ , the spectrum of  $C_2^2$  is concentrated on the positive orbit of the mapping  $t \mapsto \mu^l t$ ,  $l = \mathbf{GCD}(n_1, n_2)$ . If  $C_2^2 \neq 0$ , then  $\sigma(C_2^2) = \{\mu^{nl} x_2, n \in \mathbb{Z}\}$  for some  $x_2 \in \sigma_{y_2}$  and all eigenvalues have the same multiplicities, see [4]. Then one can choose a basis in the representation space  $\mathcal{H}$  so that  $\mathcal{H} \simeq l_2(\mathbb{Z}) \otimes \mathcal{K}_1$  and

$$C_2^2 = x_2 D(\mu^l) \otimes \mathbf{1}.$$

Let  $l = n_1 m_1 + n_2 m_2$ ,  $m_1, m_2 \in \mathbb{Z}$ , put  $U := U_1^{m_1} U_2^{m_2}$ , then

$$C_2^2 U^* = \mu^l U^* C_2^2,$$

and using unitary equivalence one can get  $U^* = E \otimes \mathbf{1}$ . Then the relations

$$C_2^2 U_i^* = \mu^{n_2} U_i^* C_2^2, \quad U U_i = U_i U, \quad i = 1, 2,$$

imply that  $U_i^* = E^{k_i} \otimes \widetilde{U}_i^*$ , where  $n_i = lk_i$ ,  $i = 1, 2$ , and  $\widetilde{U}_1, \widetilde{U}_2$  are unitaries. Analogously, from

$$C_2^2 C_j^2 = C_j^2 C_2^2, \quad C_j^2 U^* = \mu^l U^* C_j^2, \quad U U_j = U_j U, \quad C_2^2 U_j = U_j C_2^2, \quad j > 2,$$

we have  $C_j^2 = D(\mu^l) \otimes \tilde{C}_j^2$  and  $U_j = \mathbf{1} \otimes \tilde{U}_j$ ,  $j > 2$ .

One can verify directly that the family  $\{C_i^2, U_i, i = 1, \dots, d\}$  is irreducible iff the family  $\{\tilde{C}_i^2, \tilde{U}_i, i = 1, \dots, d\}$  is irreducible and the second family determines the first one up to a unitary equivalence.

Let us now rewrite the relations (2,3) in terms of the operators  $\tilde{C}_i^2, \tilde{U}_i$ . It is easy to show that (2,3) are equivalent to

$$(4) \quad \tilde{C}_j^2 \tilde{U}_i^* = \tilde{U}_i^* \tilde{C}_j^2, \quad \tilde{U}_i \tilde{U}_j = \tilde{U}_j \tilde{U}_i, \quad i = 1, 2, \quad j > 2$$

and

$$(5) \quad \begin{aligned} \tilde{C}_i^2 \tilde{U}_i^* &= \tilde{U}_i^* \left( -(1 - \alpha_2^2)x_2 + \alpha_i^2 \tilde{C}_i^2 - \sum_{3 \leq j < i} (1 - \alpha_j^2) \tilde{C}_j^2 \right), \quad i = 3, \dots, d, \\ \tilde{C}_i^2 \tilde{U}_j^* &= \alpha_j^2 \tilde{U}_j^* \tilde{C}_i^2, \quad i > j, \quad \tilde{C}_i^2 \tilde{U}_j^* = \tilde{U}_j^* \tilde{C}_i^2, \quad i < j, \\ \tilde{U}_i \tilde{U}_j^* &= \tilde{U}_j^* \tilde{U}_i, \quad \tilde{U}_i \tilde{U}_j = \tilde{U}_j \tilde{U}_i, \quad \tilde{C}_i \tilde{C}_j = \tilde{C}_j \tilde{C}_i, \quad i \neq j. \end{aligned}$$

Since  $\tilde{U}_i, i = 1, 2$  are unitaries, the Schur lemma and relations (4) imply that  $\tilde{U}_i = e^{i\phi_i} \mathbf{1}$ ,  $i = 1, 2$ ,  $\phi_1 m_1 + \phi_2 m_2 = 0 \pmod{2\pi}$ .

Furthermore, since

$$\tilde{C}_3^2 \tilde{U}_3^* = \tilde{U}_3^* \left( -(1 - \alpha_2^2)x_2 + \alpha_3^2 \tilde{C}_3^2 \right), \quad \tilde{C}_3^2 \tilde{U}_j^* = \tilde{U}_j^* \tilde{C}_3^2, \quad j > 3,$$

in the irreducible representation, the spectrum of  $\tilde{C}_3^2$  is concentrated on the positive orbit of the mapping

$$h_3(t, x_2) = -(1 - \alpha_2^2)x_2 + \alpha_3^2 t.$$

For this mapping we have the unique positive orbit, the anti-Fock one

$$\sigma(\tilde{C}_3^2) = \overline{\{h_3^{-n}(0, x_2), n \in \mathbb{Z}_+\}}$$

and, as above, all eigenvalues have the same multiplicities. Then  $\mathcal{K}_1 = l_2(\mathbb{Z}_-) \otimes \mathcal{K}_2$  and, up to a unitary equivalence,

$$\tilde{C}_3^2 = \hat{d}(h_3(0, x_2)) \otimes \mathbf{1}, \quad \tilde{U}_3^* = \hat{S} \otimes \mathbf{1}$$

and the relations (5) imply that

$$\tilde{C}_j^2 = \hat{d}(\alpha_3^2) \otimes \hat{C}_j^2, \quad \tilde{U}_j^* = \mathbf{1} \otimes \hat{U}_j^*, \quad j > 3,$$

where the family  $\{\hat{C}_j, \hat{U}_j, j > 3\}$  should be irreducible and satisfy the relations (5) with  $d - 3$  generators. Finally, note that the family  $\{\hat{C}_j, \hat{U}_j, j > 3\}$  determines the family  $\{\tilde{C}_j, \tilde{U}_j, j > 2\}$  up to a unitary equivalence. Then the evident induction on the number of generators completes the proof.  $\square$

It remains only to consider the third type of orbits.

**Proposition 3.** *Let*

$$\sigma(C_1^2) = \overline{\left\{ \frac{1 - \alpha_1^{2n}}{1 - \alpha_1^2} + \alpha_1^{2n} x_1, n \in \mathbb{Z} \right\}}$$

in an irreducible representation of  $A_\alpha$  for some fixed  $x_1 \in \tau_{y_1}$ . Then, up to a unitary equivalence, the representation space is  $\mathcal{H} = l_2(\mathbb{Z}) \otimes \bigotimes_{k=2}^d l_2(\mathbb{Z}_-)$  and

$$\begin{aligned} C_1^2 &= D(f_1, x_1) \otimes \bigotimes_{2 \leq k \leq d} \mathbf{1}, & U_1^* &= E \otimes \bigotimes_{2 \leq k \leq d} \mathbf{1}, \\ C_i^2 &= D(\alpha_1^2) \otimes \bigotimes_{2 \leq k < i} \widehat{d}(\alpha_k^2) \otimes \widehat{d}(u_i(0, x_1)) \otimes \bigotimes_{i < k \leq d} \mathbf{1}, \\ U_i^* &= \mathbf{1} \otimes \bigotimes_{2 \leq k < i} \mathbf{1} \otimes \widehat{S} \otimes \bigotimes_{i < k \leq d} \mathbf{1}, & i &= 2, \dots, d, \end{aligned}$$

where

$$D(f_1, x_1): l_2(\mathbb{Z}) \rightarrow l_2(\mathbb{Z}), \quad D(f_1, x_1)e_n = \left( \frac{1 - \alpha_1^{2n}}{1 - \alpha_1^2} + \alpha_1^{2n} x_1 \right) e_n, \quad n \in \mathbb{Z}$$

and  $u_i(t, x_1) = 1 - x_1 + \alpha_i^2 t$ ,  $i = 2, \dots, d$ ,

$$\widehat{d}(u_i(0, x_1)): l_2(\mathbb{Z}_-) \rightarrow l_2(\mathbb{Z}_-), \quad \widehat{d}(u_i(0, x_1))e_{-n} = u_i^{-n}(0, x_1)e_{-n}, \quad n \in \mathbb{Z}_+.$$

*Proof.* As in the proof of Proposition 2, we will use induction on the number of generators. If  $\sigma(C_1^2) = \overline{O_{x_1}}$ , then, up to a unitary equivalence,  $\mathcal{H} = l_2(\mathbb{Z}) \otimes \mathcal{K}_1$  and

$$C_1^2 = D(f_1, x_1) \otimes \mathbf{1}, \quad U_1^* = E \otimes \mathbf{1}.$$

The relations (2,3) imply that

$$C_i^2 = D(\alpha_1^2) \otimes \widetilde{C}_i^2, \quad U_i^* = \mathbf{1} \otimes \widetilde{U}_i^*, \quad i \geq 2,$$

where the family  $\{\widetilde{C}_i, \widetilde{U}_i, i \geq 2\}$  is irreducible and determines  $\{C_i, U_i, i \geq 1\}$  up to a unitary equivalence. Moreover, the following relations are satisfied:

$$\begin{aligned} \widetilde{C}_i^2 \widetilde{U}_i^* &= \widetilde{U}_i^* \left( 1 - x_1 + \alpha_i^2 \widetilde{C}_i^2 - \sum_{2 \leq j \leq i-1} (1 - \alpha_j^2) \widetilde{C}_j^2 \right), \quad i = 2, \dots, d, \\ (6) \quad \widetilde{C}_i^2 \widetilde{U}_j^* &= \alpha_j^2 \widetilde{U}_j^* \widetilde{C}_i^2, \quad i > j, \quad \widetilde{C}_i^2 \widetilde{U}_j^* = \widetilde{U}_j^* \widetilde{C}_i^2, \quad i < j, \\ \widetilde{U}_i \widetilde{U}_j^* &= \widetilde{U}_j^* \widetilde{U}_i, \quad \widetilde{U}_i \widetilde{U}_j = \widetilde{U}_j \widetilde{U}_i, \quad \widetilde{C}_i \widetilde{C}_j = \widetilde{C}_j \widetilde{C}_i \quad i \neq j. \end{aligned}$$

In particular, the spectrum of  $\widetilde{C}_2^2$  is concentrated on the positive orbit of the mapping

$$u_2(t, x_1) = 1 - x_1 + \alpha_2^2 t,$$

since  $x_1 > \frac{1}{1 - \alpha_1^2} > 1$  and  $\alpha_2^2 < 1$ , the unique positive orbit of  $u_2(t, x_1)$  is the anti-Fock orbit. Then the proof is analogous to the final part of the proof of Proposition 2.  $\square$

To get a general description of representations of  $A_\alpha$ , we have to combine the results of Propositions 1,2,3. Namely, let us construct three types of representations.

The *first* is the Fock one:  $\mathcal{H} = \bigotimes_{k=1}^d l_2(\mathbb{Z}_+)$ ,

$$C_j^2 = \bigotimes_{k < j} d(\alpha_k^2) \otimes d(f_j) \otimes \bigotimes_{k > j} \mathbf{1}, \quad U_j^* = \bigotimes_{k < j} \mathbf{1} \otimes S \otimes \bigotimes_{k > j} \mathbf{1}, \quad j = 1, \dots, d.$$

The *second* type is the representations with first  $i - 1$  generators as in the Fock representation and with

$$\sigma(C_i^2) = \overline{\left\{ \alpha_1^{2n_1} \dots \alpha_{i-1}^{2n_{i-1}} \frac{1}{1 - \alpha_i^2}, n_1, \dots, n_{i-1} \in \mathbb{Z}_+ \right\}}.$$

Let firstly  $i < d$ , then fix any  $t_i \in \mathbb{Z}_+$  such that  $i + t_i \leq d$ . If  $i + t_i < d$  put  $s_i := i + t_i + 1$  and fix  $y_{s_i} > 0$ ,  $\sigma_{y_{s_i}} = (\mu^{l_{i s_i}} y_{s_i}, y_{s_i}]$ , where  $l_{i s_i} = \mathbf{GCD}(n_i, n_{s_i})$ ,  $\alpha_i^2 = \mu^{n_i}$ ,  $\alpha_{s_i}^2 = \mu^{n_{s_i}}$ .

Let also  $n_i = l_{i s_i} k_i$ ,  $n_{s_i} = l_{i s_i} k_{s_i}$ . Then construct the family of operators acting on the space

$$\mathcal{H} = \bigotimes_{k=1}^{i-1} l_2(\mathbb{Z}_+) \otimes l_2(\mathbb{Z}) \otimes \bigotimes_{k=s_i+1}^d l_2(\mathbb{Z}_-)$$

by the formulas

$$\begin{aligned} C_j^2 &= \bigotimes_{k<j} d(\alpha_k^2) \otimes d(f_j) \otimes \bigotimes_{k>j, k \geq s_i} \mathbf{1}, \quad U_j^* = \bigotimes_{k<j} \mathbf{1} \otimes S \otimes \bigotimes_{k>j, k \geq s_i} \mathbf{1}, \quad j < i, \\ C_i^2 &= \frac{1}{1 - \alpha_1^2} \bigotimes_{k<i} d(\alpha_k^2) \otimes \bigotimes_{k \geq s_i} \mathbf{1}, \quad U_i^* = e^{i\phi_i} \bigotimes_{k<i} \mathbf{1} \otimes E^{k_i} \otimes \bigotimes_{k>s_i} \mathbf{1}, \\ C_j^2 &= 0, \quad U_j = 0, \quad i < j < s_i - 1, \\ C_{s_i}^2 &= \bigotimes_{k<i} d(\alpha_k^2) \otimes x_{s_i} D(\mu^{l_{i s_i}}) \otimes \bigotimes_{k>s_i} \mathbf{1}, \quad U_{s_i}^* = e^{i\phi_{s_i}} \bigotimes_{k<i} \mathbf{1} \otimes E^{k_{s_i}} \otimes \bigotimes_{k>s_i} \mathbf{1}, \\ C_j^2 &= \bigotimes_{k<i} d(\alpha_k^2) \otimes D(\mu^{l_{i s_i}}) \otimes \bigotimes_{s_i < k < j} \widehat{d}(\alpha_k^2) \widehat{d}(h_j(0, x_{s_i})) \otimes \bigotimes_{k>j} \mathbf{1}, \quad j > s_i, \\ U_j^* &= \bigotimes_{k<i} \mathbf{1} \otimes \bigotimes_{s_i \leq k < j} \widehat{S} \otimes \bigotimes_{k>j} \mathbf{1}, \quad j > s_i, \end{aligned}$$

where  $m_i \phi_i + m_{s_i} \phi_{s_i} = 0 \pmod{2\pi}$ ,  $x_{s_i} \in \sigma_{y_{s_i}}$  is fixed, and

$$h_j(t, x_{s_i}) = -(1 - \alpha_{s_i}^2) x_{s_i} + \alpha_j^2 t.$$

If  $i = d$ , then  $\mathcal{H} = \bigotimes_{k=1}^{d-1} l_2(\mathbb{Z}_+)$  and

$$\begin{aligned} C_j^2 &= \bigotimes_{k<j} d(\alpha_k^2) \otimes d(f_j) \otimes \bigotimes_{k>j} \mathbf{1}, \quad U_j^* = \bigotimes_{k<j} \mathbf{1} \otimes S \otimes \bigotimes_{k>j} \mathbf{1}, \quad j = 1, \dots, d-1, \\ C_d^2 &= \frac{1}{1 - \alpha_d^2} \bigotimes_{k<d} d(\alpha_k^2), \quad U_d^* = e^{i\phi_d} \bigotimes_{k<d} \mathbf{1}. \end{aligned}$$

In the *third* type representations, the generators  $C_j^2, U_j^*, j = 1, \dots, i-1$ , are as in the Fock representation and

$$\sigma(C_i^2) = \left\{ \alpha_1^{2n_1} \dots \alpha_{i-1}^{2n_{i-1}} \left( \frac{1 - \alpha_i^{2n_i}}{1 - \alpha_i^2} + \alpha_i^{2n_i} x_i \right), n_1, \dots, n_{i-1} \in \mathbb{Z}_+, n_i \in \mathbb{Z} \right\},$$

where  $x_i \in \tau_{y_i} = (1 + \alpha_i^2 y_i, y_i]$ ,  $y_i > \frac{1}{1 - \alpha_i^2}$  is fixed.

In this case we have

$$\mathcal{H} = \bigotimes_{k<i} l_2(\mathbb{Z}_+) \otimes l_2(\mathbb{Z}) \otimes \bigotimes_{k>i} l_2(\mathbb{Z}_-)$$

and

$$\begin{aligned} C_j^2 &= \bigotimes_{k<j} d(\alpha_k^2) \otimes d(f_j) \otimes \bigotimes_{k>j} \mathbf{1}, \quad U_j^* = \bigotimes_{k<j} \mathbf{1} \otimes S \otimes \bigotimes_{k>j} \mathbf{1}, \quad j < i, \\ C_i^2 &= \bigotimes_{k<i} d(\alpha_k^2) \otimes D(f_i, x_i) \otimes \bigotimes_{k>i} \mathbf{1}, \quad U_i^* = \bigotimes_{k<i} \mathbf{1} \otimes E \otimes \bigotimes_{k>i} \mathbf{1}, \\ C_j^2 &= \bigotimes_{k<i} d(\alpha_k^2) \otimes D(\alpha_i^2) \otimes \bigotimes_{i < k < j} \widehat{d}(\alpha_k^2) \widehat{d}(u_j(0, x_i)) \otimes \bigotimes_{k>j} \mathbf{1}, \quad j > i, \\ U_j^* &= \bigotimes_{k<j} \mathbf{1} \otimes \widehat{S} \otimes \bigotimes_{k>j} \mathbf{1}, \quad j > i, \end{aligned}$$

where  $u_j(t, x_i) = 1 - x_i + \alpha_j^2 t$ .

Combining the results of Propositions 1,2,3 we get the following theorem.

**Theorem 1.** *Any irreducible representation of  $A_\alpha$  belongs to one of the types described above. Representations corresponding to the different types or to the different parameters within the same type are non-equivalent.*

*Acknowledgments.* The author expresses his gratitude to Prof. Yu. Samoïlenko and Dr. D. Proskurin for a setting of the problem and useful discussions.

#### REFERENCES

1. P. E. T. Jørgensen, D. P. Proskurin, and Yu. S. Samoïlenko, *The kernel of Fock representations of Wick algebras with braided operator of coefficients*, Pacific J. Math. **198** (2001), 109–122.
2. P. E. T. Jørgensen, D. P. Proskurin, and Yu. S. Samoïlenko, *Generalized canonical commutation relations: Representations and stability of universal enveloping  $C^*$ -algebra*, Symmetry in Nonlinear Mathematical Physics, Part 2, Proceedings of the Institute of Mathematics of the Ukrainian National Academy of Sciences: Mathematics and Its Applications, vol. 43, Kyiv, 2002, pp. 456–460.
3. V. L. Ostrovskyĭ and D. P. Proskurin, *Operator relations, dynamical systems, and representations of a class of Wick algebras*, Oper. Theory Adv. Appl., vol. 118, Birkhauser Verlag, Basel, 2000, pp. 335–345.
4. V. L. Ostrovskyĭ and Yu. Samoïlenko, *Introduction to the Theory of Representations of Finitely Presented  $*$ -Algebras, I: Representations by Bounded Operators*, The Gordon and Breach Publishing Group, London, 1999.
5. W. Pusz and S. L. Woronowicz, *Twisted second quantization*, Rep. Math. Phys. **27** (1989), 251–263.

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS'KA,  
KYIV, 01601, UKRAINE

*E-mail address:* sukretniy@gmail.com

Received 10/04/2009; Revised 15/07/2009