

## ON THE RANGE AND KERNEL OF TOEPLITZ AND LITTLE HANKEL OPERATORS

NAMITA DAS AND PABITRA KUMAR JENA

**ABSTRACT.** In this paper we study the interplay between the range and kernel of Toeplitz and little Hankel operators on the Bergman space. Let  $T_\phi$  denote the Toeplitz operator on  $L_a^2(\mathbb{D})$  with symbol  $\phi \in L^\infty(\mathbb{D})$  and  $S_\psi$  denote the little Hankel operator with symbol  $\psi \in L^\infty(\mathbb{D})$ . We have shown that if  $\text{Ran}(T_\phi) \subseteq \text{Ran}(S_\psi)$  then  $\phi \equiv 0$  and find necessary and sufficient conditions for the existence of a positive bounded linear operator  $X$  defined on the Bergman space such that  $T_\phi X = S_\psi$  and  $\text{Ran}(S_\psi) \subseteq \text{Ran}(T_\phi)$ . We also obtain necessary and sufficient conditions on  $\psi \in L^\infty(\mathbb{D})$  such that  $\text{Ran}(T_\psi)$  is closed.

### 1. INTRODUCTION

Let  $dA(z)$  be the Lebesgue area measure on the open unit disk  $\mathbb{D}$  normalized so that the measure of the disk  $\mathbb{D}$  equals 1. The Bergman space  $L_a^2(\mathbb{D})$  is the Hilbert space consisting of analytic functions on  $\mathbb{D}$  that are also in  $L^2(\mathbb{D}, dA)$ . For  $z \in \mathbb{D}$ , the Bergman reproducing kernel is the function  $K_z \in L_a^2(\mathbb{D})$  such that  $f(z) = \langle f, K_z \rangle$  for every  $f \in L_a^2(\mathbb{D})$ . The normalized reproducing kernel  $k_z$  is the function  $\frac{K_z}{\|K_z\|_2}$ . Here the norm  $\|\cdot\|_2$  and the inner product  $\langle \cdot, \cdot \rangle$  are taken in the space  $L^2(\mathbb{D}, dA)$ . For any  $n \geq 0, n \in \mathbb{Z}$ , let  $e_n(z) = \sqrt{n+1}z^n$ . The sequence  $\{e_n\}_{n=0}^\infty$  forms an orthonormal basis for  $L_a^2(\mathbb{D})$ . Let

$$K(z, \bar{w}) = \overline{K_z(w)} = \frac{1}{(1 - z\bar{w})^2} = \sum_{n=0}^{\infty} e_n(z) \overline{e_n(w)}.$$

For  $\phi \in L^\infty(\mathbb{D})$ , the Toeplitz operator  $T_\phi$  with symbol  $\phi$  is the operator on  $L_a^2(\mathbb{D})$  defined by  $T_\phi f = P(\phi f)$ ; here  $P$  is the orthogonal projection from  $L^2(\mathbb{D}, dA)$  onto  $L_a^2(\mathbb{D})$ . For  $\phi \in L^\infty(\mathbb{D})$ , the big Hankel operator  $H_\phi$  is a mapping from  $L_a^2(\mathbb{D})$  into  $(L_a^2(\mathbb{D}))^\perp$  defined by  $H_\phi f = (I - P)(\phi f)$  for  $f \in L_a^2(\mathbb{D})$ . Let  $\overline{L_a^2(\mathbb{D})}$  be the space of conjugate analytic functions in  $L^2(\mathbb{D}, dA)$ . Clearly,  $\overline{L_a^2(\mathbb{D})} = \{\bar{f} : f \in L_a^2(\mathbb{D})\}$  is closed in  $L^2(\mathbb{D}, dA)$ . For  $\phi \in L^\infty(\mathbb{D})$ , the little Hankel operator  $h_\phi$  is a mapping from  $L_a^2(\mathbb{D})$  into  $\overline{L_a^2(\mathbb{D})}$  defined by  $h_\phi f = \overline{P(\phi f)}$ ,  $f \in L_a^2(\mathbb{D})$  where  $\overline{P}$  is the orthogonal projection from  $L^2(\mathbb{D}, dA)$  onto  $\overline{L_a^2(\mathbb{D})}$ .

There are also many equivalent ways of defining little Hankel operators. For example, define the map  $S_\phi$  from  $L_a^2(\mathbb{D})$  into  $L_a^2(\mathbb{D})$  by  $S_\phi f = PJ(\phi f)$ , where  $J$  is the self-adjoint, unitary mapping from  $L^2(\mathbb{D}, dA)$  into itself given by  $Jh(z) = h(\bar{z})$ . Notice that  $JS_\phi = h_\phi$ . Thus  $S_\phi$  is unitarily equivalent to  $h_\phi$ .

Let  $H^\infty(\mathbb{D})$  be the space of bounded analytic functions on  $\mathbb{D}$ . Let  $\text{Aut}(\mathbb{D})$  be the Lie group of all automorphisms (biholomorphic mappings) of  $\mathbb{D}$ . We can define for each  $a \in \mathbb{D}$ , an automorphism  $\phi_a$  in  $\text{Aut}(\mathbb{D})$  such that

(i)  $(\phi_a \circ \phi_a)(z) \equiv z$ ;

---

2000 *Mathematics Subject Classification.* 47B38, 47B35.

*Key words and phrases.* Toeplitz operators, little Hankel operators, Bergman space, inner functions, range and kernel of operators.

(ii)  $\phi_a(0) = a, \phi_a(a) = 0$ ;

(iii)  $\phi_a$  has a unique fixed point in  $\mathbb{D}$ .

In fact,  $\phi_a(z) = \frac{a-z}{1-\bar{a}z}$  for all  $a$  and  $z$  in  $\mathbb{D}$ . An easy calculation shows that the derivative of  $\phi_a$  at  $z$  is equal to  $-k_a(z)$ . It follows that the real Jacobian determinant of  $\phi_a$  at  $z$  is  $J_{\phi_a(z)} = |k_a(z)|^2 = \frac{(1-|a|^2)^2}{|1-\bar{a}z|^4}$ . Let  $h^\infty(\mathbb{D})$  be the space of bounded harmonic functions on  $\mathbb{D}$ . Then  $h^\infty(\mathbb{D}) \subset L^\infty(\mathbb{D})$ . For  $H$  a nonzero complex Hilbert space, let  $\mathcal{L}(H)$  denote the algebra of all bounded linear operators from the Hilbert space  $H$  into itself. Hence  $\mathcal{L}(L_a^2(\mathbb{D}))$  is the space of all bounded linear operators from  $L_a^2(\mathbb{D})$  into itself. Let  $C(\mathbb{D})$  be the algebra of complex-valued, continuous functions on  $\mathbb{D}$ , the Euclidean closure of  $\mathbb{D}$  and  $C_0(\mathbb{D})$  be the subalgebra of  $C(\mathbb{D})$  consisting of functions  $f$  with  $f(z) \rightarrow 0$  as  $|z| \rightarrow 1^-$ .

In this paper we study the interplay between the range and kernel of little Hankel and Toeplitz operators defined on the Bergman space. In Section 2, we show that  $\text{Ran}(S_\phi) = L_a^2(\mathbb{D})$  if and only if  $S_\phi$  is bounded below and  $\ker(S_\phi) = \{0\}$  if and only if  $\ker(S_\phi^2) = \{0\}$ . Further if  $S_\phi$  is normal then  $\ker(S_\phi) = \ker(S_{\phi^+}) = \ker(S_\phi S_{\phi^+})$  where  $\phi^+(z) = \overline{\phi(\bar{z})}$ . We then establish that if  $T_\phi S_\psi, T_\phi S_\psi^2, T_\phi^2 S_\psi$  are positive then  $\ker(S_\psi) = \{0\}$  implies  $T_\phi \geq 0$  and  $\ker(T_\phi) = \{0\}$  implies  $S_\psi \geq 0$  and if further  $T_\phi S_\psi$  is invertible then  $T_\phi \geq 0$  and  $S_\psi \geq 0$ . Thus if we know a priori that the products  $T_\phi S_\psi, T_\phi S_\psi^2, T_\phi^2 S_\psi$  are positive and if the kernel of the operators  $T_\phi, S_\psi$  are trivial then the operators  $T_\phi, S_\psi$  are positive. This gives us the motivation to investigate when the products  $T_\phi S_\psi$  and  $S_\psi T_\phi$  are positive.

We then proceed to show that there is no nonzero Toeplitz and Hankel operators  $T$  such that  $T^k$  is compact for some  $k \in \mathbb{Z}_+$  and  $\ker(T) = \ker(T^2)$  and  $\text{Ran}(T) = \text{Ran}(T^2)$ . On the other hand, if there is a little Hankel operator  $S_\psi$  such that  $S_\psi^k$  is compact for some  $k \in \mathbb{Z}_+$  and  $\ker(S_\psi) = \ker(S_\psi^2)$  and  $\text{Ran}(S_\psi) = \text{Ran}(S_\psi^2)$  then the symbol  $\psi$  admits a particular form. More precisely, in this case  $\psi = \phi + \bar{\chi}$  where  $\chi \in (\overline{L_a^2})^\perp \cap L^\infty(\mathbb{D})$  and  $\bar{\phi}$  is a linear combination of the Bergman kernels and some of their derivatives.

We further establish that there is no nonzero little Hankel operator whose range contains the range of a Toeplitz operator and we obtain sufficient conditions on  $\phi$  and  $\psi$  such that  $\ker(T_\phi) \subset \ker(S_\psi)$ . We obtain conditions on  $\phi, \psi \in L^\infty(\mathbb{D})$  such that  $T_\phi$  commutes with  $S_\psi$  and showed that if  $S_\psi$  intertwines  $T_\phi$  and  $T_{\bar{\phi}}$  and  $\ker(S_\psi)$  is trivial then  $T_\phi$  and  $T_{\bar{\phi}}$  are unitarily equivalent.

In Section 3, we show that if  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$  then  $\ker(T_{1-\bar{\phi}}) = \{0\}$  if and only if  $T_{\frac{1+\phi}{2}}^n$  converges to 0 weakly. We further prove that if  $\|\phi\|_\infty \leq 1$  then  $T_{\frac{1+\phi}{2}}^n$  converges to 0 in norm if and only if  $\ker(T_{1-\bar{\phi}}) = \{0\}$  and  $\text{Ran}(T_{1-\bar{\phi}})$  is closed. We find necessary and sufficient conditions for the existence of a positive bounded linear operator  $X$  defined on the Bergman space such that  $T_\phi X = S_\psi$  and  $\text{Ran}(S_\psi) \subseteq \text{Ran}(T_\phi)$ . We also obtain necessary and sufficient conditions on  $\psi \in L^\infty(\mathbb{D})$  such that  $\text{Ran}(T_\psi)$  is closed.

## 2. KERNEL OF LITTLE HANKEL OPERATORS

Let  $\mathbb{T}$  denote the unit circle in the complex plane  $\mathbb{C}$ . It is well known in case of Hardy space [20] that the Hankel operator  $S$  has a nontrivial kernel if and only if the range of  $S$  is not dense and in this case the symbol of the Hankel operator  $S$  is of the form  $z\bar{u}h$  where  $u$  is an inner function in  $H^\infty(\mathbb{T})$  and  $h \in H^\infty(\mathbb{T})$ . These results play important roles in deriving the algebraic and asymptotic properties of Toeplitz and Hankel operators. In this paper we investigate whether similar results are possible for operators on the Bergman space.

In the following lemma, we show that  $\text{Ran}(S_\phi) = L_a^2(\mathbb{D})$  if and only if  $S_\phi$  is bounded below and  $\ker(S_\phi)$  is trivial if and only if  $\ker(S_\phi^2)$  is trivial.

**Lemma 2.1.** *Let  $S_\phi$  be a little Hankel operator on  $L_a^2(\mathbb{D})$  with symbol  $\phi \in L^\infty(\mathbb{D})$ . Then the following holds:*

- (a):  $\ker(S_\phi) = \{0\}$  if and only if  $\ker(S_\phi^2) = \{0\}$ .
- (b):  $\text{Ran}(S_\phi) = L_a^2(\mathbb{D})$  if and only if  $S_\phi$  is bounded below (i.e., there exists an  $\epsilon > 0$  such that  $\|S_\phi f\| \geq \epsilon \|f\|$  for all  $f \in L_a^2(\mathbb{D})$ ).

*Proof.* (a) To prove (a) the points to note are the following:

- (i)  $S_\phi^* = S_{\phi^+}$  where  $\phi^+(z) = \overline{\phi(\bar{z})}$ .
- (ii)  $\ker(S_\phi) = \{0\}$  if and only if  $\ker(S_{\phi^+}) = \{0\}$ .
- (iii)  $\ker(S_\phi^2) = \{0\}$  if and only if  $\ker(S_{\phi^+}^2) = \{0\}$ .

These can be verified as follows :

- (i) For  $f, g \in L_a^2(\mathbb{D})$ ,

$$\begin{aligned} \langle S_\phi^* f, g \rangle &= \langle f, S_\phi g \rangle = \langle f, PJ(\phi g) \rangle = \langle f, (J\phi)Jg \rangle = \langle \overline{J\phi}f, Jg \rangle = \langle \phi^+ f, Jg \rangle \\ &= \langle J(\phi^+ f), g \rangle = \langle PJ(\phi^+ f), g \rangle = \langle S_{\phi^+} f, g \rangle. \end{aligned}$$

Thus  $S_\phi^* = S_{\phi^+}$  where  $\phi^+(z) = \overline{\phi(\bar{z})}$ .

(ii) Let  $f \in \ker S_\phi$ . Then  $S_\phi f = PJ(\phi f) = 0$ . This implies  $\phi f \in (\overline{L_a^2(\mathbb{D})})^\perp$ . That is,  $\int_{\mathbb{D}} \phi(z)f(z)g(z)dA(z) = 0$  for all  $g \in L_a^2(\mathbb{D})$ . Thus  $\int_{\mathbb{D}} \phi^+(z)f^+(z)g^+(z)dA(z) = 0$  for all  $g^+ \in L_a^2(\mathbb{D})$  and therefore  $\int_{\mathbb{D}} \phi^+(z)f^+(z)g(z)dA(z) = 0$  for all  $g \in L_a^2(\mathbb{D})$ . Hence  $\langle \phi^+ f^+, g \rangle = 0$  for all  $g \in L_a^2(\mathbb{D})$ . That is,  $f^+ \in \ker(S_{\phi^+})$ . Similarly one can verify that if  $f^+ \in \ker(S_{\phi^+})$  then  $f \in \ker(S_\phi)$ . Thus  $f \in \ker(S_\phi)$  if and only if  $f^+ \in \ker(S_{\phi^+})$ . It hence follows that  $\ker(S_\phi) = \{0\}$  if and only if  $\ker(S_{\phi^+}) = \{0\}$ . This proves (ii).

(iii) Now let  $f \in \ker(S_\phi^2)$ . Then either  $f \in \ker(S_\phi)$  or  $f \notin \ker(S_\phi)$ .

Case 1: If  $f \in \ker(S_\phi)$  then  $f^+ \in \ker(S_{\phi^+})$ .

Case 2: If  $f \notin \ker(S_\phi)$  then since  $S_\phi^2 f = 0$  hence  $(S_\phi f)^+ \in \ker(S_{\phi^+})$ . Thus  $\ker(S_{\phi^+}) = \{0\}$  implies  $\ker(S_\phi^2) = \{0\}$ .

Conversely, if  $g \in \ker(S_{\phi^+})$ , then  $g^+ \in \ker(S_\phi)$ . Hence  $S_\phi^2 g^+ = 0$ . Thus  $\ker(S_\phi^2) = \{0\}$  implies  $\ker(S_{\phi^+}) = \{0\}$ .

To prove (iii) suppose  $\ker(S_\phi^2) = \{0\}$ . This happens if and only if  $\ker(S_{\phi^+}) = \{0\}$ . But from (ii) this is true if and only if  $\ker(S_\phi) = \{0\}$ . Proceeding similarly one can show that  $\ker(S_\phi) = \{0\}$  if and only if  $\ker(S_{\phi^+}^2) = \{0\}$ .

Thus (a) follows.

(b) Suppose  $\text{Ran}(S_\phi) = L_a^2(\mathbb{D})$ . Then  $\ker(S_\phi^*) = \{0\}$  and hence from (a) it follows that  $\ker(S_\phi) = \{0\}$ . Thus from bounded inverse theorem [21] and [5], it follows that  $S_\phi$  is bounded below. Conversely, if  $S_\phi$  is bounded below then  $\ker(S_\phi) = \{0\}$  and hence from (a) it follows that  $\ker(S_\phi^*) = \{0\}$ . This implies  $\overline{\text{Ran}(S_\phi)} = L_a^2(\mathbb{D})$ . Since  $S_\phi$  is bounded below and it has dense range, hence from [5], it follows that  $S_\phi$  is invertible and  $\text{Ran}(S_\phi) = L_a^2(\mathbb{D})$ .

This proves (b). □

In Lemma 2.2 we establish that if  $S_\phi$  is normal then  $\ker(S_\phi)$  coincides with  $\ker(S_\phi^*) = \ker(S_{\phi^+}) = \ker(S_\phi S_{\phi^+})$ .

**Lemma 2.2.** *If  $S_\phi$  is normal then  $\ker(S_\phi) = \ker(S_{\phi^+}) = \ker(S_\phi S_{\phi^+})$  where  $\phi^+(z) = \overline{\phi(\bar{z})}$ .*

*Proof.* Suppose  $S_\phi$  is normal. Let  $f \in \ker(S_\phi)$ . Then  $S_\phi S_{\phi^+} f = S_{\phi^+} S_\phi f = 0$ . Thus  $S_{\phi^+} f \in \ker(S_\phi) = (\overline{\text{Ran}(S_{\phi^+})})^\perp$ . But  $S_{\phi^+} f \in \text{Ran}(S_{\phi^+}) \subseteq \overline{\text{Ran}(S_{\phi^+})}$ . Hence  $S_{\phi^+} f = 0$ . That is,  $f \in \ker(S_{\phi^+})$ . Therefore

$$(2.1) \quad \ker(S_\phi) \subset \ker(S_{\phi^+}).$$

Conversely, let  $f \in \ker(S_{\phi+})$ . This implies,  $f^+ \in \ker(S_\phi)$ . From (2.1) it follows that,  $f^+ \in \ker(S_{\phi+})$ . Therefore  $f \in \ker(S_\phi)$ . Hence

$$(2.2) \quad \ker(S_{\phi+}) \subset \ker(S_\phi).$$

From (2.1) and (2.2), we obtain  $\ker(S_\phi) = \ker(S_{\phi+})$ .

We shall now show that  $\ker(S_{\phi+}) = \ker(S_\phi S_{\phi+})$ . Let  $f \in \ker(S_{\phi+})$ . This implies,  $S_{\phi+}f = 0$ . That is,  $S_\phi S_{\phi+}f = 0$ . So  $f \in \ker(S_\phi S_{\phi+})$ . Therefore,  $\ker(S_{\phi+}) \subseteq \ker(S_\phi S_{\phi+})$ .

Now let  $f \in \ker(S_\phi S_{\phi+})$ . This implies,  $S_\phi S_{\phi+}f = 0$ . Then  $S_{\phi+}f \in \ker(S_\phi) = \overline{(\text{Ran}(S_{\phi+}))}^\perp$ . But  $S_{\phi+}f \in \text{Ran}(S_{\phi+}) \subseteq \overline{\text{Ran}(S_{\phi+})}$ . Hence  $S_{\phi+}f = 0$ . That is,  $f \in \ker(S_{\phi+})$ . Therefore,  $\ker(S_\phi S_{\phi+}) \subseteq \ker(S_{\phi+})$ . Hence  $\ker(S_{\phi+}) = \ker(S_\phi S_{\phi+})$ .  $\square$

Using Lemma 2.1 and Lemma 2.2, we now prove the following proposition which gives us sufficient conditions for  $T_\phi$  and  $S_\psi$  to be positive.

**Proposition 2.3.** *Let  $\phi, \psi \in L^\infty(\mathbb{D})$ . Suppose  $T_\phi S_\psi \geq 0, T_\phi^2 S_\psi \geq 0, T_\phi S_\psi^2 \geq 0$ .*

(i) *If  $\ker(S_\psi) = \{0\}$  then  $T_\phi \geq 0$ . Similarly if  $\ker(T_\phi) = \{0\}$  then  $S_\psi \geq 0$ .*

(ii) *If  $T_\phi S_\psi$  is invertible then  $S_\psi$  is similar to  $S_{\psi+}$ ,  $T_\phi \geq 0$  and  $S_\psi \geq 0$ . Here  $\psi^+(z) = \overline{\psi(\bar{z})}$ .*

*Proof.* (i) From Lemma 2.1 it follows that  $\ker(S_\psi) = \{0\}$  if and only if  $\ker(S_\psi^*) = \{0\}$ . Since  $T_\phi S_\psi \geq 0$ , hence  $T_\phi S_\psi^2 = (T_\phi S_\psi^2)^* = S_\psi^*(T_\phi S_\psi)^* = S_\psi^* T_\phi S_\psi$ . Thus we have, for all  $f \in L_a^2(\mathbb{D})$ ,

$$\langle T_\phi S_\psi f, S_\psi f \rangle = \langle S_\psi^* T_\phi S_\psi f, f \rangle = \langle T_\phi S_\psi^2 f, f \rangle \geq 0.$$

Now since  $\ker(S_\psi) = \{0\}$ , we obtain  $\ker(S_\psi^*) = \{0\}$ . Hence  $\overline{\text{Ran}(S_\psi)} = (\ker(S_\psi^*))^\perp = \{0\}^\perp = L_a^2(\mathbb{D})$ . It therefore follows that  $T_\phi \geq 0$ .

Further  $T_\phi^2 S_\psi = (T_\phi^2 S_\psi)^* = (T_\phi T_\phi S_\psi)^* = (T_\phi S_\psi)^* T_\phi^* = T_\phi S_\psi T_\phi^*$ . Hence

$$\langle S_\psi T_\phi^* f, T_\phi^* f \rangle = \langle T_\phi S_\psi T_\phi^* f, f \rangle = \langle T_\phi^2 S_\psi f, f \rangle \geq 0.$$

Now if  $\ker(T_\phi) = \{0\}$  then  $\overline{\text{Ran}(T_\phi^*)} = L_a^2(\mathbb{D})$ . Thus it follows that,  $S_\psi \geq 0$ .

To prove (ii) assume  $T_\phi S_\psi$  is invertible. This implies  $\ker(T_\phi S_\psi) = \{0\}$ . Notice that  $\ker(S_\psi) \subset \ker(T_\phi S_\psi)$ . Hence  $\ker(S_\psi) = \{0\}$ . Therefore  $\ker(S_\psi^*) = \{0\}$ . This implies  $\overline{\text{Ran}(S_\psi)} = L_a^2(\mathbb{D})$ . Now  $S_\psi = (T_\phi S_\psi)^{-1} T_\phi S_\psi^2 = (T_\phi S_\psi)^{-1} S_\psi^* T_\phi S_\psi$ . Thus  $S_\psi$  is similar to  $S_\psi^* = S_{\psi+}$  and  $\overline{\text{Ran}(S_\psi)} = (\ker(S_\psi^*))^\perp = (\ker(S_\psi))^\perp = L_a^2(\mathbb{D})$ . From (i) it follows that  $T_\phi \geq 0$ . Since  $\ker(T_\phi^*) \subset \ker(S_\psi^* T_\phi^*) = \{0\}$ , we obtained that  $T_\phi^*$  is injective. Now  $T_\phi^* = (T_\phi S_\psi)^{-1} (T_\phi S_\psi) T_\phi^* = (T_\phi S_\psi)^{-1} T_\phi S_\psi T_\phi^* = (T_\phi S_\psi)^{-1} T_\phi^2 S_\psi = (T_\phi S_\psi)^{-1} T_\phi (T_\phi S_\psi)$ . Hence  $T_\phi^*$  is similar to  $T_\phi$ . Since  $\ker(T_\phi^*) = \{0\}$ , hence  $\ker(T_\phi) = \{0\}$ . This implies  $\overline{\text{Ran}(T_\phi^*)} = L_a^2(\mathbb{D})$ . From (i) it follows that  $S_\psi \geq 0$ .  $\square$

In the following theorem we show that there is no nonzero Toeplitz and Hankel operators  $T$  such that  $T^k$  is compact for some  $k \in \mathbb{Z}_+$  and  $\ker(T) = \ker(T^2)$  and  $\text{Ran}(T) = \text{Ran}(T^2)$ . On the other hand, if there is a little Hankel operator  $S_\psi$  such that  $S_\psi^k$  is compact for some  $k \in \mathbb{Z}_+$  and  $\ker(S_\psi) = \ker(S_\psi^2)$  and  $\text{Ran}(S_\psi) = \text{Ran}(S_\psi^2)$  then the symbol  $\psi$  admits a particular form. More precisely, in this case  $\psi = \phi + \bar{\chi}$  where  $\chi \in (\overline{L_a^2})^\perp \cap L^\infty(\mathbb{D})$  and  $\bar{\phi}$  is a linear combination of the Bergman kernels and some of their derivatives.

**Theorem 2.4.** *If  $T \in \mathcal{L}(L_a^2(\mathbb{D}))$  is such that  $\ker(T) = \ker(T^2)$ ,  $\text{Ran}(T) = \text{Ran}(T^2)$  and  $T^k$  is compact for some  $k \in \mathbb{Z}_+$ , then the following holds:*

(i): *If  $T = T_\phi$  for some  $\phi \in L^\infty(\mathbb{D})$  then  $\phi \equiv 0$ .*

(ii): *If  $T = H_\phi$  for some  $\phi \in L^\infty(\mathbb{D})$  then  $\phi \in H^\infty(\mathbb{D})$ .*

(iii): *If  $T = S_\psi$  for some  $\psi \in L^\infty(\mathbb{D})$  then  $\psi = \phi + \bar{\chi}$  where  $\chi \in (\overline{L_a^2})^\perp \cap L^\infty(\mathbb{D})$  and  $\bar{\phi}$  is a linear combination of the Bergman kernels and some of their derivatives.*

*Proof.* (i) Suppose  $T = T_\phi$  for some  $\phi \in L^\infty(\mathbb{D})$ . By our hypothesis it follows from [21] that  $L_a^2(\mathbb{D}) = \ker(T_\phi) \oplus \text{Ran}(T_\phi)$ . Since  $\ker(T_\phi)$  is always a closed subspace of  $L_a^2(\mathbb{D})$ , we obtain from [9] that  $\text{Ran}(T_\phi)$  is closed. But  $\text{Ran}(T_\phi) = \text{Ran}(T_\phi^m)$  for all integers  $m \geq 1$ . This can be verified by Mathematical induction.

We shall only verify for  $m = 3$ . Let  $g \in \text{Ran}(T_\phi^3)$ . That implies,  $g = T_\phi^3 f = T_\phi(T_\phi^2 f)$  for some  $f \in L_a^2(\mathbb{D})$ . That is,  $g \in \text{Ran}(T_\phi) = \text{Ran}(T_\phi^2)$ . Therefore,  $\text{Ran}(T_\phi^3) \subset \text{Ran}(T_\phi^2)$ . Conversely, let  $g \in \text{Ran}(T_\phi^2)$ . That implies,  $g = T_\phi^2 h = T_\phi(T_\phi h)$ . Since  $T_\phi h \in \text{Ran}(T_\phi) = \text{Ran}(T_\phi^2)$ . This implies,  $T_\phi h = T_\phi^2 k$  for some  $k \in L_a^2(\mathbb{D})$ . That is,  $g = T_\phi(T_\phi h) = T_\phi(T_\phi^2 k) = T_\phi^3 k$ . So  $g \in \text{Ran}(T_\phi^3)$ . Therefore,  $\text{Ran}(T_\phi^2) \subset \text{Ran}(T_\phi^3)$ . Hence  $\text{Ran}(T_\phi^3) = \text{Ran}(T_\phi^2)$ .

Using induction one can show that  $\text{Ran}(T_\phi^m) = \text{Ran}(T_\phi)$  for all  $m \geq 1$ . But  $T_\phi^k$  is compact for some  $k \in \mathbb{Z}_+$ . Hence [5]  $\text{Ran}(T_\phi^k)$  shall not contain any closed infinite dimensional subspace of  $L_a^2(\mathbb{D})$ . Thus  $\text{Ran}(T_\phi)$  is a finite dimensional subspace of  $L_a^2(\mathbb{D})$  and  $T_\phi$  is a finite rank operator. It then follows from [17] that  $\phi \equiv 0$ .

(ii) Let  $T = H_\phi$ , the big Hankel operator with symbol  $\phi \in L^\infty(\mathbb{D})$ . Proceeding similarly as in (i), one can show that  $H_\phi$  is a finite rank operator. Thus  $\ker(H_\phi) \neq \{0\}$  and hence it is clear from [7] that  $\ker(H_\phi) = L_a^2(\mathbb{D})$  and  $\phi \in H^\infty(\mathbb{D})$ .

(iii) Let  $T = S_\psi$ , the little Hankel operator with symbol  $\psi \in L^\infty(\mathbb{D})$ . Proceeding similarly as in (i), one can show that  $S_\psi$  is a finite rank operator. Therefore [7]  $\psi = \phi + \chi$  where  $\chi \in (\overline{L_a^2})^\perp \cap L^\infty(\mathbb{D})$  and  $\bar{\phi}$  is a linear combination of the Bergman kernels and some of their derivatives.  $\square$

In Theorem 2.5, we show that the range of a nonzero little Hankel operator can never contain the range of a Toeplitz operator and if  $\text{Ran}(S_\phi) \subseteq \text{Ran}(T_\phi)$  then  $\|P(\bar{\phi}Jf)\|^2 \leq c\|P(\bar{\phi}f)\|^2$  for some constant  $c > 0$  and for all  $f \in L_a^2(\mathbb{D})$ .

**Theorem 2.5.** *Let  $\phi, \psi \in L^\infty(\mathbb{D})$ . Then the following holds:*

- (i):  $\text{Ran}(T_\phi) \subseteq \text{Ran}(S_\psi)$  if and only if  $\phi \equiv 0$ .
- (ii): If  $\text{Ran}(S_\phi) \subseteq \text{Ran}(T_\phi)$  then  $\|P(\bar{\phi}Jf)\|^2 \leq c\|P(\bar{\phi}f)\|^2$  for all  $f \in L_a^2(\mathbb{D})$  and for some constant  $c > 0$ .

*Proof.* (i) If  $\phi = 0$ , then  $T_\phi = 0$ . Hence  $\text{Ran}(T_\phi) = \{0\} \subseteq \text{Ran}(S_\psi)$ .

Suppose  $\text{Ran}(T_\phi) \subseteq \text{Ran}(S_\psi)$ . By [6] there is a constant  $c > 0$  such that  $T_\phi T_\phi^* \leq cS_\psi S_\psi^*$ . Hence  $\langle T_\phi T_\phi^* f, f \rangle \leq c \langle S_\psi S_\psi^* f, f \rangle$  for all  $f \in L_a^2(\mathbb{D})$ . That is,

$$\begin{aligned} \|T_\phi^* f\|^2 &\leq c \|S_\psi^* f\|^2 = c \|S_{\psi^+} f\|^2 = c \|Jh_{\psi^+} f\|^2 = c \|h_{\psi^+} f\|^2 \\ &\leq c \|H_{\psi^+} f\|^2 \quad \text{for all } f \in L_a^2(\mathbb{D}). \end{aligned}$$

Thus

$$\begin{aligned} c^{-1} \|P(\bar{\phi}f)\|^2 &\leq \|H_{\psi^+} f\|^2 = \|(I - P)(\psi^+ f)\|^2 \\ &= \langle \psi^+ f - P(\psi^+ f), \psi^+ f - P(\psi^+ f) \rangle = \|\psi^+ f\|^2 - \|P(\psi^+ f)\|^2. \end{aligned}$$

Hence  $c^{-1} \|P(\bar{\phi}f)\|^2 + \|P(\psi^+ f)\|^2 \leq \|\psi^+ f\|^2 \leq \|\psi^+\|_\infty^2 \|f\|^2$ . This implies

$$c^{-1} \frac{\|P(\bar{\phi}f)\|^2}{\|f\|^2} + \frac{\|P(\psi^+ f)\|^2}{\|f\|^2} \leq \|\psi^+\|_\infty^2 \quad \text{for all } f \in L_a^2(\mathbb{D}).$$

Thus  $c^{-1} \|\bar{\phi}\|_\infty^2 + \|\psi^+\|_\infty^2 \leq \|\psi^+\|_\infty^2$ . Hence  $\|\bar{\phi}\|_\infty^2 = 0$  and  $\phi \equiv 0$ .

(ii) If  $\text{Ran}(S_\phi) \subseteq \text{Ran}(T_\phi)$  then by [6] we have  $S_\phi S_\phi^* \leq cT_\phi T_\phi^*$  for some constant  $c > 0$ . That is,

$$\|S_\phi f\|^2 = \langle S_\phi S_\phi^* f, f \rangle \leq c \langle T_\phi T_\phi^* f, f \rangle = c \|T_\phi^* f\|^2 \quad \text{for all } f \in L_a^2(\mathbb{D}).$$

Hence  $\|PJ(\phi^+ f)\|^2 \leq c\|P(\bar{\phi}f)\|^2$  for all  $f \in L_a^2(\mathbb{D})$ . Thus  $\|P(\bar{\phi}Jf)\|^2 \leq c\|P(\bar{\phi}f)\|^2$  for all  $f \in L_a^2(\mathbb{D})$ .  $\square$

**Definition 1.** A function  $G \in L_a^2(\mathbb{D})$  is called an inner function in  $L_a^2(\mathbb{D})$  if  $|G|^2 - 1$  is orthogonal to  $H^\infty(\mathbb{D})$ .

For more details about Bergman space inner divisors, see [14]. In the following theorem, we find conditions on  $\phi, \psi \in L^\infty(\mathbb{D})$  such that  $\ker(T_\phi) \subseteq \ker(S_\psi)$ .

**Theorem 2.6.** *Suppose  $\phi \in L^\infty(\mathbb{D})$  is such that  $\ker(T_\phi) = \{0\}$ . Let*

$$\bar{\psi}(z) = \sum_{j=1}^N \sum_{v=0}^{m_j-1} c_{jv} \frac{\partial^v}{\partial \bar{b}_j^v} K_{b_j}(z),$$

where  $\mathbf{b} = \{b_j\}_{j=1}^N$  is a finite set of points in  $\mathbb{D}$ ,  $c_{jv} \neq 0$  for all  $j, v$  and  $m_j$  is the number of times  $b_j$  appears in  $\mathbf{b}$ . Let  $S_\psi T_\phi = T_\phi S_\psi$ . Then there exists an inner function  $G \in H^\infty(\mathbb{D})$  such that  $\ker(T_\phi^*) \subseteq \ker(S_\psi^*) = GL_a^2(\mathbb{D})$ .

*Proof.* Since  $\bar{\psi}(z) = \sum_{j=1}^N \sum_{v=0}^{m_j-1} c_{jv} \frac{\partial^v}{\partial \bar{b}_j^v} K_{b_j}(z)$  where  $\mathbf{b} = \{b_j\}_{j=1}^N$  is a finite set of points in  $\mathbb{D}$ ,  $c_{jv} \neq 0$  for all  $j, v$  and  $m_j$  is the number of times  $b_j$  appears in  $\mathbf{b}$ , hence the operator  $S_\psi$  is a [7] finite rank operator on  $L_a^2(\mathbb{D})$  and there exists an inner function  $G \in H^\infty(\mathbb{D})$  such that  $\ker(S_\psi^*) = GL_a^2(\mathbb{D})$ . Thus there exists a system of linearly independent vectors  $\zeta_i, i = 1, 2, \dots, n$  and a system of nonzero bounded linear functionals  $\phi_i$  for  $i = 1, 2, \dots, n$  on  $L_a^2(\mathbb{D})$  such that

$$S_\psi f = \sum_{i=1}^n \phi_i(f) \zeta_i, f \in L_a^2(\mathbb{D}).$$

Moreover,

$$\sum_{i=1}^n \phi_i(f) T_\phi \zeta_i = T_\phi S_\psi f = S_\psi T_\phi f = \sum_{i=1}^n \phi_i(T_\phi f) \zeta_i, f \in L_a^2(\mathbb{D}).$$

On the other hand, since  $T_\phi$  is injective, it is clear that the vectors  $T_\phi \zeta_i, i = 1, 2, \dots, n$  are linearly independent. Hence  $S_\psi f \in \text{span}\{\zeta_1, \zeta_2, \dots, \zeta_n\} = \text{span}\{T_\phi \zeta_1, \dots, T_\phi \zeta_n\}$  for all  $f \in L_a^2(\mathbb{D})$ . Thus  $\overline{\text{Ran}(S_\psi)} = \text{Ran}(S_\psi) \subseteq \text{Ran}(T_\phi) \subseteq \overline{\text{Ran}(T_\phi)}$  and therefore  $\ker(T_\phi^*) = (\overline{\text{Ran}(T_\phi)})^\perp \subseteq (\overline{\text{Ran}(S_\psi)})^\perp = \ker(S_\psi^*) = GL_a^2(\mathbb{D})$  for some inner [7] function  $G \in H^\infty(\mathbb{D})$ .  $\square$

**Definition 2.** An operator  $A$  defined on a Hilbert space  $H$  is said to be hyponormal if and only if  $A^*A - AA^* \geq 0$ .

In the following theorem we find conditions on  $\phi \in L^\infty(\mathbb{D})$  such that  $T_\phi$  commutes with  $S_\psi$  where  $S_\psi$  is a positive little Hankel operator on  $L_a^2(\mathbb{D})$  with trivial kernel.

**Theorem 2.7.** *Let  $\phi \in C_0(\mathbb{D}), \|\phi\|_\infty \leq 1$ . Suppose  $\psi \in L^\infty(\mathbb{D})$  and  $S_\psi$  is a positive little Hankel operator on  $L_a^2(\mathbb{D})$  such that  $\ker(S_\psi) = \{0\}$  and  $S_\psi \leq T_\phi S_\psi T_\phi$ . Then  $T_\phi$  is unitary and  $T_\phi S_\psi = S_\psi T_\phi$ .*

*Proof.* The operator  $S_\psi^{\frac{1}{2}} T_\phi$  is compact [23] since  $\phi \in C_0(\mathbb{D})$ . By Lemma 2.1,  $\ker(S_\psi^*) = \{0\}$ . Let  $S_\psi^{\frac{1}{2}} T_\phi = A$ . Then

$$AA^* = S_\psi^{\frac{1}{2}} T_\phi T_\phi^* S_\psi^{\frac{1}{2}} \leq S_\psi.$$

Now

$$0 \leq T_\phi S_\psi T_\phi - S_\psi \leq T_\phi S_\psi T_\phi - S_\psi^{\frac{1}{2}} T_\phi T_\phi^* S_\psi^{\frac{1}{2}} = A^*A - AA^*.$$

Thus the operator  $A$  is hyponormal and as  $A$  is compact, the [10] operator  $A$  is normal. Therefore,

$$S_\psi = T_\phi S_\psi T_\phi = S_\psi^{\frac{1}{2}} T_\phi T_\phi^* S_\psi^{\frac{1}{2}}$$

and  $T_{\bar{\phi}}$  is an isometry in  $\overline{\text{Ran}(S_{\psi})} = (\ker(S_{\psi}^*))^{\perp} = \{0\}^{\perp} = L_a^2(\mathbb{D})$ . Further,  $S_{\psi}$  commutes with  $T_{\phi}$  and  $T_{\bar{\phi}}$ . It follows therefore that

$$T_{\bar{\phi}}T_{\phi}S_{\psi} = T_{\bar{\phi}}S_{\psi}T_{\phi} = S_{\psi} = S_{\psi}T_{\phi}T_{\bar{\phi}}.$$

Hence  $T_{\phi}$  is unitary.  $\square$

In Theorem 2.8, we show that if  $S_{\psi}$  is a little Hankel operator on  $L_a^2(\mathbb{D})$  with trivial kernel and  $S_{\psi}$  intertwines  $T_{\phi}$  and  $T_{\bar{\phi}}$  then  $T_{\phi}$  and  $T_{\bar{\phi}}$  are unitarily equivalent.

**Theorem 2.8.** *Suppose  $T_{\phi}S_{\psi} = S_{\psi}T_{\phi}^*$  and  $T_{\phi}^*S_{\psi} = S_{\psi}T_{\phi}$  and  $\ker(S_{\psi}) = \{0\}$ ,  $\phi, \psi \in L^{\infty}(\mathbb{D})$ . Then  $T_{\phi}$  and  $T_{\bar{\phi}}$  are unitarily equivalent. Further,  $T_{\phi}^*T_{\phi}$  and  $T_{\phi}T_{\phi}^*$  are unitarily equivalent.*

*Proof.*  $T_{\phi}S_{\psi}S_{\psi}^* = S_{\psi}T_{\phi}^*S_{\psi}^* = S_{\psi}S_{\psi}^*T_{\phi}$ . Thus  $T_{\phi}$  commutes with  $S_{\psi}S_{\psi}^*$ . Proceeding similarly as in Lemma 2.2, one can verify that

$$\overline{\text{Ran}(S_{\psi})} = (\ker(S_{\psi}^*))^{\perp} = (\ker(S_{\psi}S_{\psi}^*))^{\perp} = \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}.$$

This can also be verified as follows: Let  $g \in \ker(S_{\psi+})$ . Then  $S_{\psi}S_{\psi+}g = 0$  and  $g \in \ker(S_{\psi}S_{\psi+})$ . Further, let  $h \in \ker(S_{\psi}S_{\psi+})$ . Then  $S_{\psi+}h \in \ker(S_{\psi})$ . But  $S_{\psi+}h \in \text{Ran}(S_{\psi+}) = \text{Ran}(S_{\psi}^*) \subset \overline{\text{Ran}(S_{\psi}^*)} = (\ker(S_{\psi}))^{\perp}$ . Hence  $S_{\psi+}h = 0$  and  $h \in \ker(S_{\psi+})$ . Thus we verify that  $\ker(S_{\psi+}) = \ker(S_{\psi}S_{\psi+})$ .

We now show that  $T_{\phi}(\overline{\text{Ran}(S_{\psi}S_{\psi}^*)}) \subset \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}$ . Let  $g \in \text{Ran}(S_{\psi}S_{\psi}^*)$ . Then  $g = S_{\psi}S_{\psi}^*f$  for some  $f \in L_a^2(\mathbb{D})$ . Hence

$$T_{\phi}g = T_{\phi}S_{\psi}S_{\psi}^*f = S_{\psi}S_{\psi}^*T_{\phi}f \in \text{Ran}(S_{\psi}S_{\psi}^*).$$

Thus  $T_{\phi}(\overline{\text{Ran}(S_{\psi}S_{\psi}^*)}) \subset \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}$ . Now let  $g \in \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}$  and  $g = \lim_{n \rightarrow \infty} g_n$ ,  $g_n \in \text{Ran}(S_{\psi}S_{\psi}^*)$ . Then  $T_{\phi}g = \lim_{n \rightarrow \infty} T_{\phi}g_n$  and  $T_{\phi}g_n \in \text{Ran}(S_{\psi}S_{\psi}^*)$ . Therefore,  $T_{\phi}g \in \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}$ . Thus  $T_{\phi}(\overline{\text{Ran}(S_{\psi}S_{\psi}^*)}) \subset \overline{\text{Ran}(S_{\psi}S_{\psi}^*)}$ .

We now proceed to verify that  $T_{\phi}(\ker(S_{\psi}S_{\psi}^*)) \subset \ker(S_{\psi}S_{\psi}^*)$ . Let  $g \in \ker(S_{\psi}S_{\psi}^*)$ . Then  $S_{\psi}S_{\psi}^*g = 0$ . Hence  $T_{\phi}S_{\psi}S_{\psi}^*g = 0$ . This implies,  $S_{\psi}S_{\psi}^*T_{\phi}g = 0$ . That is,  $T_{\phi}g \in \ker(S_{\psi}S_{\psi}^*)$ . Thus  $\overline{\text{Ran}(S_{\psi})}$  is a reducing subspace of  $T_{\phi}$ . Proceeding similarly one can show that  $T_{\phi}$  commutes  $S_{\psi}^*S_{\psi}$  and  $\ker(S_{\psi}) = \ker(S_{\psi}^*S_{\psi})$  reduces  $T_{\phi}$ .

Further, let  $S_{\psi} = VQ$  be the polar decomposition of  $S_{\psi}$  such that  $\ker(V) = \ker(Q)$ . Here  $V$  is the partial isometry and  $Q$  is the positive operator. Let  $f \in \ker(Q)$ . Then  $Qf = 0$  and therefore  $S_{\psi}f = VQf = 0$ . Hence  $f \in \ker(S_{\psi}) = \{0\}$ . Thus  $f = 0$  and  $\ker(V) = \ker(Q) = \{0\}$ . Since  $V^*Vf = f$ ,  $f \in (\ker(V))^{\perp} = \{0\}^{\perp} = L_a^2(\mathbb{D})$ , hence  $V$  is an isometry.

Since  $T_{\phi}S_{\psi}^*S_{\psi} = S_{\psi}^*S_{\psi}T_{\phi}$ , we obtain  $T_{\phi}QV^*VQ = QV^*VQT_{\phi}$ . Thus  $T_{\phi}Q^2 = Q^2T_{\phi}$  and therefore [5], we have  $T_{\phi}Q = QT_{\phi}$  and  $QT_{\phi}^* = T_{\phi}^*Q$ . Now  $T_{\phi}S_{\psi} = S_{\psi}T_{\phi}^*$  implies  $T_{\phi}VQ = VQT_{\phi}^* = VT_{\phi}^*Q$ . Thus

$$(2.3) \quad (T_{\phi}V - VT_{\phi}^*)Qf = 0 \quad \text{for all } f \in L_a^2(\mathbb{D}).$$

Notice that  $(\overline{\text{Ran}(S_{\psi})})^{\perp} = \ker(S_{\psi}^*) = \{0\}$ , hence  $\overline{\text{Ran}(S_{\psi})} = L_a^2(\mathbb{D})$  and  $\overline{\text{Ran}(Q)} = (\ker(Q))^{\perp} = (\ker(S_{\psi}))^{\perp} = \{0\}^{\perp} = L_a^2(\mathbb{D})$ . From equation (2.3) it follows that  $\text{Ran}(Q) \subset \ker(T_{\phi}V - VT_{\phi}^*)$ .

Let  $h \in \overline{\text{Ran}(Q)}$  and  $h = \lim_{n \rightarrow \infty} h_n$  where  $h_n \in \text{Ran}(Q)$ . Then

$$(T_{\phi}V - VT_{\phi}^*)h = (T_{\phi}V - VT_{\phi}^*)(\lim_{n \rightarrow \infty} h_n) = \lim_{n \rightarrow \infty} (T_{\phi}V - VT_{\phi}^*)h_n = 0.$$

Thus  $L_a^2(\mathbb{D}) = \overline{\text{Ran}(Q)} \subset \ker(T_{\phi}V - VT_{\phi}^*)$  and  $T_{\phi}V = VT_{\phi}^*$ . Similarly, since  $T_{\phi}^*S_{\psi} = S_{\psi}T_{\phi}$  we obtain  $T_{\phi}^*V = VT_{\phi}$ . Thus  $V^*T_{\phi}V = T_{\bar{\phi}}$ .

Since  $S_{\psi} = VQ$ , we have  $S_{\psi}^* = S_{\psi+} = QV^*$ . Let  $f \in \ker V^*$ . Then  $V^*f = 0$  and hence  $S_{\psi}^*f = QV^*f = 0$ . Thus by Lemma 2.1,  $f \in \ker(S_{\psi}^*) = \{0\}$  and  $f = 0$ . Thus

$\ker(V^*) = \{0\}$ . Therefore  $\{0\} = \ker(V^*) = (\text{Ran}(V))^\perp$  and  $\overline{\text{Ran}(V)} = L_a^2(\mathbb{D})$ . Since  $\|Vf\| = \|f\|$ , hence  $V$  is bounded below with dense range. By [5],  $V$  is invertible. Therefore  $T_\phi$  and  $T_\phi^*$  are unitarily equivalent. Further,  $T_\phi^*T_\phi = V^*T_\phi T_\phi^*V$  and  $T_\phi^*T_\phi$  and  $T_\phi T_\phi^*$  are unitarily equivalent.  $\square$

### 3. RANGE AND KERNEL OF TOEPLITZ OPERATORS

A well known lemma attributed to Coburn [5] states that a bounded Toeplitz operator with nontrivial kernel acting on the Hardy space must have dense range. That is, if  $\phi$  is a function in  $L^\infty(\mathbb{T})$  not almost everywhere zero, then either  $\ker(T_\phi) = \{0\}$  or  $\ker(T_\phi^*) = \{0\}$ . Vukotic [22] showed that the range of a nonzero Toeplitz operator with  $\ker(T_\phi) \neq \{0\}$  must contain all polynomials. Further, if  $\ker(T_\phi) \neq \{0\}$  then [11]  $\ker(T_\phi) = g(H^2(\mathbb{T}) \ominus z\theta H^2(\mathbb{T}))$  where  $g$  is an outer function and  $\theta$  is an inner function in  $H^\infty(\mathbb{T})$ .

In this section we show that if  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$  then  $\ker(T_{1-\bar{\phi}}) = \{0\}$  if and only if  $T_{\frac{1+\phi}{2}}^n$  converges to 0 weakly. We further prove that if  $\|\phi\|_\infty \leq 1$  then  $T_{\frac{1+\phi}{2}}^n$  converges to 0 in norm if and only if  $\ker(T_{1-\bar{\phi}}) = \{0\}$  and  $\text{Ran}(T_{1-\phi})$  is closed. We find necessary and sufficient conditions for the existence of a positive bounded linear operator  $X$  defined on the Bergman space such that  $T_\phi X = S_\psi$  and  $\text{Ran}(S_\psi) \subseteq \text{Ran}(T_\phi)$ . We also obtain necessary and sufficient conditions on  $\psi \in L^\infty(\mathbb{D})$  such that  $\text{Ran}(T_\psi)$  is closed.

**Theorem 3.1.** *Let  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$  and  $\text{Ran}(T_{1-\phi})$  be closed. If  $\ker(T_{1-\phi}) \oplus \text{Ran}(T_{1-\phi})$  is closed then there exists a closed complementary subspace  $M$  of  $\ker(T_{1-\phi})$  containing  $\text{Ran}(T_{1-\phi})$ .*

*Proof.* First we shall show that  $\ker(T_{1-\phi}) \cap \text{Ran}(T_{1-\phi}) = \{0\}$ . Let  $V = T_{\frac{1+\phi}{2}}$ . Then the range and the kernel of the operator  $I - V$  coincide with those of  $I - T_\phi$ . Let  $g \in \ker(I - V) \cap \text{Ran}(I - V)$ . Since  $(I - V)g = 0$ , that is,  $Vg = g$ , we have  $V^n g = g$  for every  $n$ . Further  $(I - V)f = g$  for some  $f \in L_a^2(\mathbb{D})$ , that is,  $g = f - Vf$ . Hence  $g = V^n f - V^{n+1} f$ . By [12],  $\|V^n f - V^{n+1} f\| \rightarrow 0$  as  $n \rightarrow \infty$ , which implies that  $g = 0$ . So  $\ker(I - V) \cap \text{Ran}(I - V) = \{0\}$ . Thus  $\ker(I - T_\phi) \cap \text{Ran}(I - T_\phi) = \{0\}$ .

Let  $L = (\ker(T_{1-\phi}) \oplus \text{Ran}(T_{1-\phi}))^\perp$  be the orthogonal complement of the closed subspace  $\ker(T_{1-\phi}) \oplus \text{Ran}(T_{1-\phi})$ . Then  $L_a^2(\mathbb{D}) = (\ker(T_{1-\phi}) \oplus \text{Ran}(T_{1-\phi})) \oplus L$ . Hence  $\text{Ran}(T_{1-\phi}) \oplus L$  is closed and  $M = \text{Ran}(T_{1-\phi}) \oplus L$  is the desired complementary subspace of  $\ker(T_{1-\phi})$ .  $\square$

Let  $L^2(\mathbb{R})$  be the usual Lebesgue space considered with the Lebesgue measure. Since both the infinite dimensional Hilbert spaces  $L_a^2(\mathbb{D})$  and  $L^2(\mathbb{R})$  are separable, they are isomorphic. Therefore, there exists a unitary map  $U$  from  $L_a^2(\mathbb{D})$  onto  $L^2(\mathbb{R})$ .

For each  $n \in \mathbb{N}$ , define the operator  $\check{L}_n$  on  $L^2(\mathbb{R})$  by  $(\check{L}_n f)(s) := e^{\frac{iq(s)}{n}} f(s)$ ,  $s \in \mathbb{R}$ ,  $f \in L^2(\mathbb{R})$  where  $q : \mathbb{R} \rightarrow [0, 1]$  is strictly monotone. It is not difficult to see that

$$\begin{aligned} \|\check{L}_n - I_{\mathcal{L}(L^2(\mathbb{R}))}\| &= \sup_{s \in \mathbb{R}} |e^{\frac{iq(s)}{n}} - 1| \\ &\leq |e^{\frac{i}{n}} - 1| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Let  $L_n = U^* \check{L}_n U$ . Then  $L_n \in \mathcal{L}(L_a^2(\mathbb{D}))$  and  $\|L_n - I_{\mathcal{L}(L_a^2(\mathbb{D}))}\| \rightarrow 0$  as  $n \rightarrow \infty$ . That is,  $L_n \rightarrow I_{\mathcal{L}(L_a^2(\mathbb{D}))}$  in norm operator topology.

In Theorem 3.2, we show that if  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$  then  $T_{\frac{1+\phi}{2}}^n \xrightarrow{w} 0$  if and only if  $T_{1-\bar{\phi}}$  has trivial kernel. We also present necessary and sufficient conditions such that  $T_{\frac{1+\phi}{2}}^n \rightarrow 0$  in norm.

**Theorem 3.2.** *Let  $\phi \in L^\infty(\mathbb{D})$  be such that  $\|\phi\|_\infty \leq 1$ . Then the following holds:*

- (i):  $\ker(T_{1-\bar{\phi}}) = \{0\}$  if and only if  $T_{\frac{1+\phi}{2}}^n$  converges to zero weakly as  $n \rightarrow \infty$ .

- (ii): If  $\ker(T_{1-\bar{\phi}}) = \{0\}$  then  $T_{\frac{1+\phi}{2}}^n L_n \xrightarrow{w} 0$ .  
 (iii):  $\ker(T_{1-\bar{\phi}}) = \{0\}$  and  $\text{Ran}(T_{1-\bar{\phi}})$  is closed if and only if  $T_{\frac{1+\phi}{2}}^n$  converges to zero in norm as  $n \rightarrow \infty$ .

*Proof.* Since  $\|\phi\|_\infty \leq 1$  we have  $\|T_{\frac{1+\phi}{2}}\| \leq 1$ . Hence the sequence  $\{T_{\frac{1+\phi}{2}}^{*n}\}_{n=0}^\infty$  is bounded. So by [1] the sequence  $\{T_{\frac{1+\phi}{2}}^{*n}\}_{n=0}^\infty$  has a subsequence which converges to an operator  $K \in \mathcal{L}(L_a^2(\mathbb{D}))$  in the weak operator topology. Without loss of generality, we shall assume the original sequence  $\{T_{\frac{1+\phi}{2}}^{*n}\}_{n=0}^\infty$  converges to an operator  $K \in \mathcal{L}(L_a^2(\mathbb{D}))$  in the weak operator topology. Hence  $\langle (T_{\frac{1+\phi}{2}}^{*n+1} f - T_{\frac{1+\phi}{2}}^{*n} f), g \rangle \rightarrow 0$  for every  $f, g \in L_a^2(\mathbb{D})$  and  $\{\langle T_{\frac{1+\phi}{2}}^{*n+1} f, g \rangle\}_{n=0}^\infty$  converges to  $\langle Kf, g \rangle$  as  $n$  tends to  $\infty$  for all  $f, g \in L_a^2(\mathbb{D})$ . This implies

$$\langle T_{\frac{1+\phi}{2}}^{*n} f, T_{\frac{1+\phi}{2}} g \rangle \rightarrow \langle Kf, g \rangle \quad \text{for all } f, g \in L_a^2(\mathbb{D}).$$

Thus  $\langle Kf, T_{\frac{1+\phi}{2}} g \rangle = \langle Kf, g \rangle$  for all  $f, g \in L_a^2(\mathbb{D})$  and therefore  $T_{\frac{1+\phi}{2}}^* K = K$ . Further since  $\{\langle T_{\frac{1+\phi}{2}}^{*n} T_{\frac{1+\phi}{2}}^* f, g \rangle\}_{n=0}^\infty$  converges to  $\langle Kf, g \rangle$  for all  $f, g \in L_a^2(\mathbb{D})$ , hence

$$\langle KT_{\frac{1+\phi}{2}}^* f, g \rangle = \langle Kf, g \rangle \quad \text{for all } f, g \in L_a^2(\mathbb{D}).$$

Thus  $KT_{\frac{1+\phi}{2}}^* = K$  and  $T_{\frac{1+\phi}{2}}^{*n} K = K$  for all  $n \in \mathbb{Z}_+$ . That is,

$$\langle T_{\frac{1+\phi}{2}}^{*n} Kf, g \rangle = \langle Kf, g \rangle \quad \text{for all } f, g \in L_a^2(\mathbb{D}).$$

Taking limit both the sides, we obtain  $K^2 = K$ . This proves that the operator  $K$  is an idempotent. Moreover,  $T_{\frac{1+\phi}{2}}^* K = K$  implies  $T_\phi^* K = K$  and  $KT_{\frac{1+\phi}{2}}^* = K$  implies  $KT_\phi^* = K$ . So  $\text{Ran}(K) \subseteq \ker(T_{1-\bar{\phi}})$ .

On the other hand, for  $f \in \ker(T_{1-\bar{\phi}})$ , we have  $T_\phi^* f = f$ , so  $T_{\frac{1+\phi}{2}}^* f = f$ . Hence  $T_{\frac{1+\phi}{2}}^{*n} f = f$  for all  $n \in \mathbb{Z}_+$  and this implies  $Kf = f$ . Hence  $\text{Ran}(K) = \ker(T_{1-\bar{\phi}})$ .

To prove the inclusion  $\text{Ran}(T_{1-\bar{\phi}}) \subseteq \ker(K)$ , let  $f \in L_a^2(\mathbb{D})$  be an arbitrary element and  $g = f - T_\phi^* f$ . Then we have  $Kg = Kf - KT_\phi^* f = Kf - Kf = 0$ . Hence  $g \in \ker(K)$ .

Thus we have shown that, if  $\|\phi\|_\infty \leq 1$  then there exists an idempotent  $K \in \mathcal{L}(L_a^2(\mathbb{D}))$  whose range is  $\ker(T_{1-\bar{\phi}})$  and kernel contains  $\text{Ran}(T_{1-\bar{\phi}})$  and there exists a subsequence of  $\{T_{\frac{1+\phi}{2}}^{*n}\}$  which converges to  $K$  weakly.

To prove (i), assume that  $\langle T_{\frac{1+\phi}{2}}^{*n} f, g \rangle \rightarrow 0$  for every  $f, g \in L_a^2(\mathbb{D})$ . Then

$$(3.1) \quad \langle f, T_{\frac{1+\phi}{2}}^{*n} g \rangle \rightarrow 0 \quad \text{for all } f, g \in L_a^2(\mathbb{D}).$$

That is, the sequence  $\{\langle f, T_{\frac{1+\phi}{2}}^{*n} g \rangle\}_{n=0}^\infty$  is a Cauchy sequence. Thus if any subsequence of  $\{\langle f, T_{\frac{1+\phi}{2}}^{*n} g \rangle\}_{n=0}^\infty$  converges to some  $r \in \mathbb{C}$ , then the sequence  $\{\langle f, T_{\frac{1+\phi}{2}}^{*n} g \rangle\}_{n=0}^\infty$  itself converges to  $r$ . We have already seen in the first part that there exists a subsequence of  $\{\langle f, T_{\frac{1+\phi}{2}}^{*n} g \rangle\}_{n=0}^\infty$  which converges to  $\langle f, Kg \rangle$  for all  $f, g \in L_a^2(\mathbb{D})$ . Thus

$$T_{\frac{1+\phi}{2}}^{*n} \xrightarrow{w} K$$

in  $\mathcal{L}(L_a^2(\mathbb{D}))$  and the operator  $K$  is an idempotent. Thus it follows from (3.1) that  $K = 0$ ,  $T_{\frac{1+\phi}{2}}^{*n} \xrightarrow{w} 0$  and  $\ker(T_{1-\bar{\phi}}) = \text{Ran}(K) = \{0\}$ .

Conversely, assume  $\ker(T_{1-\bar{\phi}}) = \{0\}$ . Then since  $T_{\frac{1+\phi}{2}}^{*n} \xrightarrow{w} K$  and the operator  $K \in \mathcal{L}(L_a^2(\mathbb{D}))$  is an idempotent and  $\text{Ran } K = \ker T_{1-\bar{\phi}} = \{0\}$  we obtain  $K = 0$  and  $T_{\frac{1+\phi}{2}}^{*n} \xrightarrow{w} 0$ .

To prove (ii) suppose  $\ker(T_{1-\bar{\phi}}) = \{0\}$ . Then by (i),  $T_{\frac{1+\phi}{2}}^n$  converges to zero weakly as  $n \rightarrow \infty$ . Since  $L_n \rightarrow I_{\mathcal{L}(L_a^2(\mathbb{D}))}$  in norm operator topology, hence it follows from [8] that  $T_{\frac{1+\phi}{2}}^n L_n \xrightarrow{w} 0$ .

To prove (iii), suppose  $\ker(T_{1-\phi}^*) = \{0\}$  and  $\text{Ran}(T_{1-\phi})$  is closed. Then  $\text{Ran}(T_{1-\phi}) = L_a^2(\mathbb{D})$  and from Theorem 3.1, it follows that,  $\ker(T_{1-\phi}) = \{0\}$ . Thus  $T_{1-\phi}$  is invertible and  $1 \notin \sigma(T_\phi)$ . But

$$I - T_{\frac{1+\phi}{2}} = I - \frac{I}{2} - \frac{T_\phi}{2} = \frac{I - T_\phi}{2}.$$

Hence  $1 \notin \sigma(T_{\frac{1+\phi}{2}})$ . It therefore follows from [13] that if  $\text{Ran}(T_{1-\phi}) = L_a^2(\mathbb{D})$  then  $\sigma(T_{\frac{1+\phi}{2}}) \cap \{z \in \mathbb{C} : |z| = 1\} = \emptyset$ .

Notice that  $\sigma(T_{\frac{1+\phi}{2}})$  is a compact subset [5] of  $\mathbb{C}$  and since  $T_{\frac{1+\phi}{2}}$  is a contraction, the spectral radius  $r(T_{\frac{1+\phi}{2}}) \leq \|T_{\frac{1+\phi}{2}}\| \leq 1$ . Further, as  $\sigma(T_{\frac{1+\phi}{2}}) \cap \{z \in \mathbb{C} : |z| = 1\} = \emptyset$ , hence  $\sigma(T_{\frac{1+\phi}{2}})$  is a compact subset of  $\mathbb{D}$  and therefore  $r(T_{\frac{1+\phi}{2}}) < 1$ . By [15],

$$\|T_{\frac{1+\phi}{2}}^n\| \rightarrow 0.$$

Conversely, assume that  $\|T_{\frac{1+\phi}{2}}^n\| \rightarrow 0$ . Then by [15],  $r(T_{\frac{1+\phi}{2}}) < 1$ . This implies that  $\sigma(T_{\frac{1+\phi}{2}}) \cap \{z \in \mathbb{C} : |z| = 1\} = \emptyset$ . Therefore  $1 \notin \sigma(T_{\frac{1+\phi}{2}})$ . Since

$$I - T_{\frac{1+\phi}{2}} = \frac{I - T_\phi}{2}.$$

So  $1 \notin \sigma(T_\phi)$ . Hence  $T_{1-\phi}$  is invertible. Hence  $\text{Ran}(T_{1-\phi}) = L_a^2(\mathbb{D})$ .  $\square$

It is not difficult to find examples of operators  $T \in \mathcal{L}(H)$  such that  $\text{Ran}(T)$  is closed but  $\text{Ran}(T^2)$  is not closed.

Let  $\{u_j, f_j, h_j, j = 1, 2, \dots\}$  be an orthonormal basis for  $H$ . Let  $\{a_n\}_{n=0}^\infty$  be a sequence of nonnegative numbers converging to zero such that  $a_n \leq 1$  for all  $n$ . For each  $n$ , let  $b_n = (1 - a_n^2)^{\frac{1}{2}}$  and set  $v_n = a_n u_n + b_n h_n, w_n = a_n f_n + b_n h_n$ .

Let  $M^\perp$  be the closed span of  $\{v_j\}$  and define  $N$  to be the closed span of  $\{w_j\}$ . Then  $M^\perp \cap N = \{0\}$  and the angle between  $M^\perp$  and  $N$  is zero since

$$\langle v_j, w_j \rangle = b_n^2 = 1 - a_n^2.$$

Let  $T$  be a partially isometry with initial space  $M$  and the final space  $N$ . Then  $\text{Ran}(T)$  is closed but  $\text{Ran}(T^2)$  is not closed. For more details see [4].

In the following lemma we shall show that if  $\|\phi\|_\infty \leq 1$  and  $\text{Ran}(T_{1-\bar{\phi}})$  is closed then  $\text{Ran}(T_{1-\bar{\phi}}^2)$  is closed.

**Lemma 3.3.** *Let  $\phi \in L^\infty(\mathbb{D})$  be such that  $\|\phi\|_\infty \leq 1$  and  $\ker(T_{1-\bar{\phi}}) = \{0\}$ . Then the following conditions are equivalent:*

- (i):  $\text{Ran}(T_{1-\bar{\phi}})$  is closed.
- (ii):  $\text{Ran}(T_{1-\bar{\phi}} T_{1-\phi})$  is closed.
- (iii):  $\text{Ran}(T_{1-\bar{\phi}}^2)$  is closed.

*Proof.* To show (i)  $\iff$  (ii), suppose  $T_{1-\bar{\phi}}$  has closed range. From [18], it follows that  $\text{Ran}(T_{1-\bar{\phi}} T_{1-\phi})$  is closed and  $\text{Ran}(T_{1-\bar{\phi}}) = \text{Ran}(T_{1-\bar{\phi}} T_{1-\phi})$ . Conversely, if  $T_{1-\bar{\phi}} T_{1-\phi}$

has closed range then

$$\begin{aligned} F &= \text{Ran } (T_{1-\bar{\phi}}T_{1-\phi}) \oplus \ker(T_{1-\bar{\phi}}T_{1-\phi}) = \text{Ran } (T_{1-\bar{\phi}}T_{1-\phi}) \oplus \ker(T_{1-\phi}) \\ &\subset \text{Ran } (T_{1-\bar{\phi}}) \oplus \ker(T_{1-\phi}) \subset F, \end{aligned}$$

which implies  $T_{1-\bar{\phi}}$  has closed range.

To prove (i)  $\implies$  (iii) suppose  $\text{Ran } (T_{1-\bar{\phi}})$  is closed. It is enough [16] to prove that the space  $\text{Ran } (T_{1-\bar{\phi}}) + \ker(T_{1-\bar{\phi}})$  is closed. Let  $\{T_{1-\bar{\phi}}f_n + g_n\}$  be a sequence in  $\text{Ran } (T_{1-\bar{\phi}}) + \ker(T_{1-\bar{\phi}})$  that converges to  $f \in L_a^2(\mathbb{D})$ . Since  $\|\phi\|_\infty \leq 1$ , from the first part of the proof of Theorem 3.2, it follows that there exists an idempotent operator  $K \in \mathcal{L}(L_a^2(\mathbb{D}))$  such that

$$T_\phi^* K = K = K T_\phi^*, \text{Ran } (T_{1-\bar{\phi}}) \subset \ker(K)$$

and

$$\ker(T_{1-\bar{\phi}}) = \text{Ran } (K).$$

Thus we obtain

$$K(T_{1-\bar{\phi}}f_n + g_n) = K g_n \longrightarrow K f.$$

As  $g_n \in \ker(T_{1-\bar{\phi}}) = \text{Ran } (K)$ , we obtain  $g_n = K x_n$  for some  $x_n \in L_a^2(\mathbb{D})$  and  $K g_n = K K x_n = K x_n = g_n$ . Since  $K g_n \longrightarrow K f$ , we obtain  $g_n \longrightarrow K f$ . Thus the sequence  $\{T_{1-\bar{\phi}}f_n\}$  converges to  $f - K f$  which must be in  $\text{Ran } (T_{1-\bar{\phi}})$ , as the space  $\text{Ran } (T_{1-\bar{\phi}})$  is closed.

To prove (iii)  $\implies$  (i) suppose  $\text{Ran } (T_{1-\bar{\phi}}^2)$  is closed. We have to show  $\text{Ran } (T_{1-\bar{\phi}})$  is closed. Let  $\{h_n\} \in \text{Ran } (T_{1-\bar{\phi}})$ , suppose  $h_n \rightarrow h$  and  $h_n = T_{1-\bar{\phi}}f_n, f_n \in L_a^2(\mathbb{D})$ .

Now  $T_{1-\bar{\phi}}f_n \rightarrow h$  implies  $T_{1-\bar{\phi}}^2 f_n \rightarrow T_{1-\bar{\phi}}h$ . That is,  $T_{1-\bar{\phi}}h \in \text{Ran } (T_{1-\bar{\phi}}^2)$ . Thus  $T_{1-\bar{\phi}}h = T_{1-\bar{\phi}}^2 k$  for some  $k \in L_a^2(\mathbb{D})$ . Therefore,  $T_{1-\bar{\phi}}^2 k - T_{1-\bar{\phi}}h = 0$ . That is,  $T_{1-\bar{\phi}}(T_{1-\bar{\phi}}k - h) = 0$ . So  $h = T_{1-\bar{\phi}}k$  as  $\ker(T_{1-\bar{\phi}}) = \{0\}$ . Hence  $h \in \text{Ran } (T_{1-\bar{\phi}})$ . Therefore  $\text{Ran } (T_{1-\bar{\phi}})$  is closed.  $\square$

*Remark 1.* It follows from Lemma 3.3 that if  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$  then  $\text{Ran } (T_{1-\bar{\phi}})$  is closed implies  $\text{Ran } (T_{1-\bar{\phi}}^2)$  is closed. We do not need the condition  $\ker(T_{1-\bar{\phi}}) = \{0\}$  in this case.

**Theorem 3.4.** *Let  $\psi = 1 - \phi$  where  $\phi \in L^\infty(\mathbb{D})$  and  $\|\phi\|_\infty \leq 1$ . Then  $\text{Ran } (T_\psi)$  is closed if and only if there exists an invertible operator  $S \in \mathcal{L}(L_a^2(\mathbb{D}))$  which commutes with  $T_\psi$  and  $S T_\psi$  is a projection operator.*

*Proof.* If  $\text{Ran } (T_\psi)$  is closed, then from [4] it follows that  $\text{Ran } (T_{\bar{\psi}})$  is closed and hence by Lemma 3.3 and Remark 1, the space  $\text{Ran } (T_{\bar{\psi}}^2)$  is also closed. Again from [4], it follows that the space  $\text{Ran } (T_\psi^2)$  is closed. Since  $\|\phi\|_\infty \leq 1$ , we obtain (see the proof of Theorem 3.1) that  $\ker(T_\psi) \cap \text{Ran } (T_\psi) = \{0\}$ .

Let  $f \in \ker(T_\psi^2)$ . Then the element  $g = T_{\bar{\psi}}f \in \ker(T_{\bar{\psi}}) \cap \text{Ran } (T_{\bar{\psi}}) = \{0\}$ . Hence  $f \in \ker(T_{\bar{\psi}})$ . Thus  $\ker(T_\psi^2) \subseteq \ker(T_{\bar{\psi}})$ . The inclusion relation  $\ker(T_{\bar{\psi}}) \subseteq \ker(T_\psi^2)$  is always true. Hence  $\ker(T_{\bar{\psi}}) = \ker(T_\psi^2)$ .

Now,

$$(\text{Ran } (T_\psi))^\perp = \ker(T_{\bar{\psi}}) = \ker(T_\psi^2) = (\text{Ran } (T_\psi^2))^\perp$$

and  $\text{Ran } (T_\psi)$  is closed. Thus we obtain

$$\text{Ran } (T_\psi^2) = \overline{\text{Ran } (T_\psi^2)} = \text{Ran } (T_\psi).$$

So for every  $f \in L_a^2(\mathbb{D})$ , there exists  $g \in L_a^2(\mathbb{D})$  such that  $T_\psi f = T_\psi^2 g$ . Thus  $f - T_\psi g$  is in  $\ker(T_\psi)$  and from Theorem 3.1, we obtain

$$\text{Ran } (T_\psi) \oplus \ker(T_\psi) = L_a^2(\mathbb{D}).$$

Now, define  $R : L_a^2(\mathbb{D}) \longrightarrow L_a^2(\mathbb{D})$  such that  $R(h + g) = T_\psi h + g$  where  $h \in \text{Ran}(T_\psi)$  and  $g \in \ker(T_\psi)$ . The mapping  $R$  is well-defined, linear. Now suppose  $h_n + g_n \rightarrow h + g$  and  $R(h_n + g_n) \rightarrow l$ . Then  $l = R(h + g)$ . This can be verified as follows:

$$\begin{aligned} l &= \lim_{n \rightarrow \infty} R(h_n + g_n) = \lim_{n \rightarrow \infty} (T_\psi h_n + g_n) \\ &= T_\psi \left( \lim_{n \rightarrow \infty} h_n \right) + \left( \lim_{n \rightarrow \infty} g_n \right) = T_\psi h + g = R(h + g). \end{aligned}$$

Thus  $R$  is a closed operator. By the closed graph theorem,  $R$  is bounded. We claim that  $R$  is invertible. That is,  $R$  is onto and one-one.

Let  $f = h + g \in L_a^2(\mathbb{D})$  where  $h \in \text{Ran}(T_\psi) = \text{Ran}(T_\psi^2)$  and  $g \in \ker(T_\psi)$ . Thus  $h = T_\psi p$  for some  $p \in \text{Ran}(T_\psi)$  and  $R(p + g) = T_\psi p + g = h + g = f$ . Hence  $R$  is onto.

Again let  $f = h + g \in \ker(R)$  where  $h \in \text{Ran}(T_\psi)$  and  $g \in \ker(T_\psi)$ . Then  $R(h + g) = 0$ . That is,  $T_\psi h = -g$ . From Theorem 3.1, it follows that  $T_\psi h = g = 0$ . This implies  $h$  is in the intersection of the spaces  $\ker(T_\psi)$  and  $\text{Ran}(T_\psi)$  and so it is 0 as well. Hence  $f = h + g = 0$  and  $R$  is one-one.

Let  $\Theta$  be the projection with range  $\text{Ran}(T_\psi)$  and kernel  $\ker(T_\psi)$ . Now let  $f = h + g, h \in \text{Ran}(T_\psi), g \in \ker(T_\psi)$ . Then

$$R\Theta f = Rh = T_\psi h = T_\psi(h + g) = T_\psi f$$

and

$$\Theta Rf = \Theta(T_\psi h + g) = T_\psi h = T_\psi(h + g) = T_\psi f.$$

Thus  $T_\psi = R\Theta = \Theta R$ . Hence  $R^{-1}T_\psi = \Theta = T_\psi R^{-1}$ . Let  $S = R^{-1}$ . Then  $ST_\psi = T_\psi S$  and  $ST_\psi = \Theta$  is a projection operator.

To prove the converse, suppose there exists an invertible operator  $S \in \mathcal{L}(L_a^2(\mathbb{D}))$  such that  $ST_\psi = T_\psi S$  and  $ST_\psi = \Theta$ , a projection operator. Let  $S^{-1} = R$ . Then  $T_\psi = R\Theta = \Theta R$ ,  $\Theta$  is a projection and  $R$  is invertible. We shall show that  $\text{Ran}(T_\psi)$  is closed. Let  $M = \text{Ran}(\Theta)$ . Then  $M$  is a closed subspace of  $L_a^2(\mathbb{D})$ . The map  $R$  is linear, one-one, onto, bounded and  $R^{-1}$  is also bounded. Hence  $R$  is a homeomorphism. Thus  $\text{Ran}(T_\psi) = R(M)$  is a closed subspace of  $L_a^2(\mathbb{D})$  as  $R$  is a closed map.  $\square$

Recall that, if  $H, K$  are two Hilbert spaces and  $C \in \mathcal{L}(H, K)$  has closed range, then there exists a unique  $C^\dagger \in \mathcal{L}(K, H)$  such that  $CC^\dagger C = C, C^\dagger CC^\dagger = C^\dagger$  and  $CC^\dagger, C^\dagger C$  are Hermitian,  $C^\dagger$  is called the Moore-Penrose inverse of  $C$  (For more details see [3]). If  $C$  is positive then  $C^\dagger \geq 0$ .

**Theorem 3.5.** *Let  $\phi \in L^\infty(\mathbb{D})$  and  $\text{Ran}(T_\phi)$  is closed. Suppose  $\psi \in L^\infty(\mathbb{D})$  and  $\text{Ran}(S_\psi)$  is a closed subspace of  $L_a^2(\mathbb{D})$  of finite codimension. Then  $\text{Ran}(S_\psi) \subset \text{Ran}(T_\phi)$  and there exists a positive operator  $X \in \mathcal{L}(L_a^2(\mathbb{D}))$  such that  $T_\phi X = S_\psi$  if and only if  $S_\psi T_\phi^\dagger \geq 0$  and  $S_\psi = S_\psi T_\phi^\dagger C$  for some  $C \in \mathcal{L}(L_a^2(\mathbb{D}))$ . The operator  $X$  is unique if  $\ker(X) = \ker(T_\phi) = \ker(S_\psi)$ . If  $X$  is invertible then  $\phi \equiv 0$ .*

*Proof.* Suppose  $\psi \in L^\infty(\mathbb{D})$  and  $\text{Ran}(S_\psi)$  is a closed subspace of  $L_a^2(\mathbb{D})$  of finite codimension. Then  $\ker(S_\psi^*)$  is finite dimensional. Hence by Lemma 2.1,  $\ker(S_\psi)$  is of finite dimension. Since  $\text{Ran}(T_\phi)$  is closed and  $\ker(S_\psi)$  is finite dimensional hence from [4] it follows that  $\text{Ran}(T_\phi^*)$  is closed and  $\text{Ran}(S_\psi T_\phi^\dagger)$  is closed. Now suppose  $S_\psi T_\phi^\dagger \geq 0$  and  $S_\psi = S_\psi T_\phi^\dagger C$  for some  $C \in \mathcal{L}(L_a^2(\mathbb{D}))$ . Then  $\text{Ran}(S_\psi) \subset \text{Ran}(S_\psi T_\phi^\dagger)$  and  $(S_\psi T_\phi^\dagger)(S_\psi T_\phi^\dagger)^\dagger S_\psi = S_\psi$ . Set  $X = S_{\psi+} (S_\psi T_\phi^\dagger)^\dagger S_\psi$ . Then  $X \geq 0$  and

$$T_\phi X = (T_\phi S_{\psi+})(S_\psi T_\phi^\dagger)^\dagger S_\psi = (S_\psi T_\phi^\dagger)(S_\psi T_\phi^\dagger)^\dagger S_\psi = S_\psi.$$

From [5] and [2], it follows that  $\text{Ran}(S_\psi) \subseteq \text{Ran}(T_\phi)$ .

We now prove the converse. Since  $T_\phi X = S_\psi$  and  $X \geq 0$ , hence  $S_\psi T_\phi^\dagger = T_\phi X T_\phi^\dagger \geq 0$ . We shall show that  $\ker(S_\psi T_\phi^\dagger) \subset \ker(S_{\psi+})$ . Suppose  $f \in L_a^2(\mathbb{D})$  be such that  $S_\psi T_\phi^\dagger f =$

$T_\phi X T_{\bar{\phi}} f = 0$ . Then by Reid's inequality [19], we have

$$\begin{aligned} \|S_{\psi+} f\|^2 &= \|X T_{\bar{\phi}} f\|^2 \\ &\leq \|X\| \langle X T_{\bar{\phi}} f, T_{\bar{\phi}} f \rangle = \|X\| \langle T_{\phi} X T_{\bar{\phi}} f, f \rangle = 0. \end{aligned}$$

Hence  $\ker(S_{\psi} T_{\bar{\phi}}) \subset \ker(S_{\psi+})$  and therefore  $\text{Ran}(S_{\psi}) = \overline{\text{Ran } S_{\psi}} \subset \text{Ran}(S_{\psi} T_{\bar{\phi}})$ . Thus by [6], we have  $S_{\psi} = S_{\psi} T_{\bar{\phi}} C$  for some  $C \in \mathcal{L}(L_a^2(\mathbb{D}))$ .

From [6] it follows that the operator  $X$  is unique if  $\ker(X) = \ker(T_{\phi}) = \ker(S_{\psi})$ . Further if the operator  $X$  is invertible and  $T_{\phi} X = S_{\psi}$  then it follows from [6], [2] that  $\text{Ran}(T_{\phi}) = \text{Ran}(S_{\psi})$ . From Theorem 2.5, it follows that  $\phi \equiv 0$ .  $\square$

#### REFERENCES

1. N. I. Akhiezer and I. M. Glazman, *Theory of Linear Operators in Hilbert Space*, Monographs and Studies in Mathematics, vol. 9, Pitman, Boston—London, 1981.
2. B. A. Barnes, *Majorization, range inclusion and factorization for bounded linear operators*, Proc. Amer. Math. Soc. **133** (2004), no. 1, 155–162.
3. A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory and Applications*, Robert E. Krieger publishing Co., Inc., Huntington, New York, 1980.
4. R. Bouldin, *The product of operators with closed range*, Tohoku Math. J. **25** (1973), 359–363.
5. R. G. Douglas, *Banach Algebra Techniques in Operator Theory*, Academic Press, New York, 1972.
6. R. G. Douglas, *On majorization, factorization and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc. **17** (1966), 413–415.
7. N. Das, *The kernel of a Hankel operator on the Bergman space*, Bull. London Math. Soc. **31** (1999), 75–80.
8. D. Deckard, R. G. Douglas, C. Pearcy, *On invariant subspaces of quasitriangular operators*, Amer. J. Math. **91** (1969), no. 3, 637–647.
9. S. Goldberg, *Unbounded Linear Operators*, McGraw-Hill, New York, 1966.
10. P. R. Halmos, *A Hilbert Space Problem Book*, 2nd ed., Springer-Verlag, New York—Heidelberg—Berlin, 1982.
11. E. Hayashi, *The kernel of a Toeplitz operator*, Integr. Equ. Oper. Theory **9** (1986), 588–591.
12. S. Ishikawa, *Fixed points and iteration of a nonexpansive mapping in a Banach space*, Proc. Amer. Math. Soc. **59** (1976), 65–71.
13. Y. Katznelson and L. Tzafriri, *On power bounded operators*, J. Funct. Anal. **68** (1986), 313–328.
14. B. Korenblum and M. Stessin, *On Toeplitz-invariant subspaces of the Bergman space*, J. Funct. Anal. **111** (1983), 76–96.
15. C. S. Kubrusly, *Similarity to contractions and weak stability*, Adv. Math. Sci. Appl. **2** (1993), 335–343.
16. T. Kato, *Perturbation theory for nullity, deficiency, and other quantities of linear operators*, J. Analyse Math. **6** (1958), 261–322.
17. D. Luecking, *Finite rank Toeplitz operators on the Bergman space*, Proc. Amer. Math. Soc. **136** (2008), no. 5, 1717–1723.
18. E. C. Lance, *Hilbert  $C^*$ -Modules*, LMS Lecture Note Series 210, Cambridge University Press, 1995.
19. C. S. Lin, *Inequalities of Reid type and Furuta*, Proc. Amer. Math. Soc. **129** (2001), no. 3, 855–859.
20. S. C. Power, *Hankel operators on Hilbert space*, Bull. Lond. Math. Soc. **12** (1980), 422–442.
21. A. E. Taylor, *Introduction to Functional Analysis*, Wiley, New York, 1958.
22. D. Vukotic, *A note on the range of Toeplitz operators*, Integr. Equ. Oper. Theory **50** (2004), 565–567.
23. K. H. Zhu, *Operator Theory in Function Spaces*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 139, Marcel Dekker Inc., New York, 1990.

P. G. DEPARTMENT OF MATHEMATICS, UTKAL UNIVERSITY, VANI VIHAR, BHUBANESWAR, 751004, ORISSA, INDIA

*E-mail address:* namitadas440@yahoo.co.in

P. G. DEPARTMENT OF MATHEMATICS, UTKAL UNIVERSITY, VANI VIHAR, BHUBANESWAR, 751004, ORISSA, INDIA

*E-mail address:* pabitrmath@gmail.com

Received 23/02/2012