

ELLIPTIC PROBLEMS IN THE SENSE OF B. LAWYUK ON TWO-SIDED REFINED SCALES OF SPACES

IRYNA S. CHEPURUKHINA AND ALEKSANDR A. MURACH

To the blessed memory of Professor V. V. Sharko

ABSTRACT. We investigate elliptic boundary-value problems with additional unknown functions on the boundary of a Euclidean domain. These problems were introduced by Lawruk. We prove that the operator corresponding to such a problem is bounded and Fredholm on two-sided refined scales built on the base of inner product isotropic Hörmander spaces. The regularity of the distributions forming these spaces are characterized by a real number and an arbitrary function that varies slowly at infinity in the sense of Karamata. For the generalized solutions to the problem, we prove theorems on a priori estimates and local regularity in these scales. As applications, we find new sufficient conditions under which the solutions have continuous classical derivatives of a prescribed order.

1. INTRODUCTION

The paper is devoted to an investigation of operators generated by elliptic boundary-value problems with additional unknown functions on the boundary of a Euclidean domain. Such problems were introduced by B. Lawruk [13, 14, 15]. They appear naturally if we pass from a general (nonregular) elliptic boundary-value problem to its formally adjoint problem. Moreover, the class of these problems is closed with respect to this passage. As to applications, various problems in hydrodynamics and the theory of elasticity proved to belong to this class [3, 8, 25].

This class has been investigated completely enough in Sobolev spaces; see the monographs by V. A. Kozlov, V. G. Maz'ya, J. Rossmann [12, Chapt. 3] and Ya. Roitberg [30, Chapt. 2]. Among the proved results are theorems on the Fredholm property of the operators corresponding to the problem under investigation, on the isomorphisms generated by these operators, on a priori estimates for solutions to the problem, theorems on the increase in local regularity of these solutions. These results have been established for the two-sided scales (of function spaces) that modify the classical Sobolev scale in the sense of Ya. Roitberg [27, 28] (see also his monograph [29, Sect. 2], where this modification is systematically used in the theory of usual elliptic boundary-value problems).

The purpose of this paper is to prove versions of these theorems for two-sided refined scales built on the base of the isotropic inner product Hörmander spaces

$$H^{s,\varphi}(\mathbb{R}^n) := \{w \in \mathcal{S}'(\mathbb{R}^n) : \langle \xi \rangle^s \varphi(\langle \xi \rangle) \widehat{w}(\xi) \in L_2(\mathbb{R}^n, d\xi)\},$$

parametrized with a number $s \in \mathbb{R}$ and a function $\varphi : [1, \infty) \rightarrow (0, \infty)$ that varies slowly at infinity in the sense of J. Karamata. Here, \widehat{w} is the Fourier transform of the tempered distribution w , whereas $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$. These spaces form a refined Sobolev scale, which was selected and investigated by V. A. Mikhailets and A. A. Murach [16, 17, 21].

2010 *Mathematics Subject Classification.* Primary 35J40, 46E35.

Key words and phrases. Elliptic boundary-value problem, slowly varying function, Hörmander space, two-sided refined scale, Fredholm operator, a priori estimate for solutions, local regularity of solutions.

This scale has an important interpolation property; namely, each space $H^{s,\varphi}(\mathbb{R}^n)$ is a result of the interpolation with an appropriate function parameter of the inner product Sobolev spaces $H^{s-\varepsilon}(\mathbb{R}^n)$ and $H^{s+\delta}(\mathbb{R}^n)$ with $\varepsilon, \delta > 0$.

V. A. Mikhailets and A. A. Murach built a theory of solvability of general elliptic boundary-value problems in the one-sided refined Sobolev scale and its two-sided modifications [16–20, 22–24]. The mentioned interpolation is a key method in their theory. In this connection, we also note papers [1, 2, 32]. However, this theory does not involve the important class of elliptic boundary-value problems in the sense of B. Lawruk.

This paper consists of seven sections. Section 1 is Introduction. In Section 2, we state an elliptic boundary-value problem in the sense of B. Lawruk and consider the corresponding Green formula and formally adjoint problem. In Section 3, we give definitions of function spaces that form the two-sided refined scales. The main results of the paper are formulated in Section 4. They are theorems on the character of solvability of the problem under investigation and on properties of its generalized solutions. Section 5 is devoted to a method of interpolation with a function parameter of Hilbert spaces and its applications to the refined scales. The main results are proved in Section 6. In Section 7, we give some applications of the refined scales to an investigation of classical smoothness of the generalized solutions.

2. STATEMENT OF THE PROBLEM

Let Ω be a bounded domain in the Euclidean space \mathbb{R}^n , with $n \geq 2$. Suppose that the boundary $\Gamma := \partial\Omega$ of Ω is an infinitely smooth closed manifold of dimension $n - 1$. As usual, $\bar{\Omega} := \Omega \cup \Gamma$. Let $\nu(x)$ denote the unit vector of the inner normal to Γ at a point $x \in \Gamma$.

We arbitrarily choose integers $q \geq 1$, $\varkappa \geq 1$, $m_1, \dots, m_{q+\varkappa} \in [0, 2q - 1]$ and r_1, \dots, r_\varkappa . We consider the linear boundary-value problem in the domain Ω with \varkappa additional unknown functions on the boundary Γ

$$(1) \quad Au = f \quad \text{in } \Omega,$$

$$(2) \quad B_j u + \sum_{k=1}^{\varkappa} C_{j,k} v_k = g_j \quad \text{on } \Gamma, \quad j = 1, \dots, q + \varkappa.$$

Here,

$$A := A(x, D) := \sum_{|\mu| \leq 2q} a_\mu(x) D^\mu$$

is a linear differential operator on $\bar{\Omega}$ of the even order $2q$. Moreover, every

$$B_j(x, D) := \sum_{|\mu| \leq m_j} b_{j,\mu}(x) D^\mu$$

is a linear boundary differential operator on Γ of order m_j , and each $C_{j,k} := C_{j,k}(x, D_\tau)$ is a linear tangent differential operator on Γ with $\text{ord } C_{j,k} \leq m_j + r_k$. All the coefficients of these differential operators are infinitely smooth complex-valued functions given on $\bar{\Omega}$ and Γ respectively.

We use the following standard notation: $\mu := (\mu_1, \dots, \mu_n)$ is a multi-index with $|\mu| := \mu_1 + \dots + \mu_n$, and $D^\mu := D_1^{\mu_1} \dots D_n^{\mu_n}$ with $D_\ell := i\partial/\partial x_\ell$, where i is the imaginary unit and $x = (x_1, \dots, x_n)$ is an arbitrary point in \mathbb{R}^n . Moreover, we put $D_\nu := i\partial/\partial \nu(x)$ and $\xi^\mu := \xi_1^{\mu_1} \dots \xi_n^{\mu_n}$ for $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$.

The function u given on Ω and all the functions v_1, \dots, v_\varkappa given on Γ are unknown in the boundary-value problem (1), (2). Note that all functions and distributions are assumed in the paper to be complex-valued.

We suppose that the boundary-value problem (1), (2) is elliptic in Ω in the sense of B. Lawruk [13]. We recall the corresponding definition (see also [12, Subsect. 3.1.3]).

Let $A^{(0)}(x, \xi)$, $B_j^{(0)}(x, \xi)$, and $C_{j,k}^{(0)}(x, \tau)$ denote the principal symbols of the differential operators $A(x, D)$, $B_j(x, D)$, and $C_{j,k}(x, D_\tau)$, respectively. Recall that

$$A^{(0)}(x, \xi) := \sum_{|\mu|=2q} a_\mu(x) \xi^\mu, \quad \text{with } x \in \overline{\Omega} \text{ and } \xi \in \mathbb{C}^n,$$

is a homogeneous polynomial of order $2q$ in ξ and that

$$B_j^{(0)}(x, \xi) := \sum_{|\mu|=m_j} b_{j,\mu}(x) \xi^\mu, \quad \text{with } x \in \Gamma \text{ and } \xi \in \mathbb{C}^n,$$

is a homogeneous polynomial of order m_j in ξ . Moreover, for each point $x \in \Gamma$, the expression $C_{j,k}^{(0)}(x, \tau)$ is a homogeneous polynomial of order $m_j + r_k$ in τ , where τ is a tangent vector to the boundary Γ at the point x . (If $\text{ord } C_{j,k} < m_j + r_k$, then $C_{j,k}^{(0)}(x, \tau) := 0$.)

The boundary-value problem (1), (2) is called elliptic in the domain Ω if the following three conditions are satisfied:

- (i) The differential operator $A(x, D)$ is elliptic at each point $x \in \overline{\Omega}$; i.e., $A^{(0)}(x, \xi) \neq 0$ for an arbitrary vector $\xi \in \mathbb{R}^n \setminus \{0\}$.
- (ii) The differential operator $A(x, D)$ is properly elliptic at each point $x \in \Gamma$; i.e., for an arbitrary tangent vector $\tau \neq 0$ to Γ at x , the polynomial $A^{(0)}(x, \tau + \zeta \nu(x))$ in $\zeta \in \mathbb{C}$ has q roots with positive imaginary part and q roots with negative imaginary part (of course, these roots are calculated with regard for their multiplicity).
- (iii) The system of boundary-value conditions (2) covers equation (1) at each point $x \in \Gamma$. This means that, for an arbitrary vector $\tau \neq 0$ from condition (ii), the boundary-value problem

$$A^{(0)}(x, \tau + D_t \nu(x)) \theta(t) = 0 \quad \text{for } t > 0,$$

$$B_j^{(0)}(x, \tau + D_t \nu(x)) \theta(t) \Big|_{t=0} + \sum_{k=1}^{\varkappa} C_{j,k}^{(0)}(x, \tau) \lambda_k = 0,$$

$$j = 1, \dots, q + \varkappa,$$

has only the trivial (zero) solution. This problem is posed relative to the unknown function $\theta \in C^\infty([0, \infty))$ with $\theta(t) \rightarrow 0$ as $t \rightarrow \infty$ and the unknown complex-valued numbers $\lambda_1, \dots, \lambda_\varkappa$. Here, $A^{(0)}(x, \tau + D_t \nu(x))$ and $B_j^{(0)}(x, \tau + D_t \nu(x))$ are differential operators with respect to $D_t := i\partial/\partial t$. We obtain them if we put $\zeta := D_t$ in the polynomials $A^{(0)}(x, \tau + \zeta \nu(x))$ and $B_j^{(0)}(x, \tau + \zeta \nu(x))$, in ζ , respectively.

Note that condition (ii) follows from condition (i) in the case of $n \geq 3$.

Various examples of the elliptic boundary-value problem (1), (2) are given in [12, Subsect. 3.1.5]. Specifically, the boundary-value problem

$$\Delta u = f \text{ in } \Omega, \quad u + v = g_1 \text{ and } D_\nu u + D_\tau v = g_2 \text{ on } \Gamma$$

is elliptic in Ω . Here, $n = 2$, $\varkappa = 1$, Δ is the Laplace operator, and $D_\tau := i\partial/\partial\tau$, where $\partial/\partial\tau$ is the derivative along the curve Γ .

With the problem (1), (2), we connect the linear mapping

$$(3) \quad \Lambda : (u, v_1, \dots, v_\varkappa) \rightarrow \left(Au, B_1 u + \sum_{k=1}^{\varkappa} C_{1,k} v_k, \dots, B_{q+\varkappa} u + \sum_{k=1}^{\varkappa} C_{q+\varkappa,k} v_k \right),$$

$$\text{with } u \in C^\infty(\overline{\Omega}) \text{ and } v_1, \dots, v_\varkappa \in C^\infty(\Gamma).$$

We will investigate properties of the extension (by continuity) of this mapping in appropriate couples of Hilbert spaces that form certain two-sided refined scales.

To describe the range of this extension, we need the following Green formula (see [12, Theorem 3.1.2]):

$$\begin{aligned} (Au, w)_\Omega + \sum_{j=1}^{q+\varkappa} \left(B_j u + \sum_{k=1}^{\varkappa} C_{j,k} v_k, h_j \right)_\Gamma \\ = (u, A^+ w)_\Omega + \sum_{j=1}^{2q} \left(D_\nu^{j-1} u, K_j w + \sum_{k=1}^{q+\varkappa} Q_{k,j}^+ h_k \right)_\Gamma + \sum_{j=1}^{\varkappa} \left(v_j, \sum_{k=1}^{q+\varkappa} C_{k,j}^+ h_k \right)_\Gamma, \end{aligned}$$

where $u, w \in C^\infty(\bar{\Omega})$, $v = (v_1, \dots, v_\varkappa) \in (C^\infty(\Gamma))^\varkappa$, $h = (h_1, \dots, h_{q+\varkappa}) \in (C^\infty(\Gamma))^{q+\varkappa}$ are arbitrary, whereas $(\cdot, \cdot)_\Omega$ and $(\cdot, \cdot)_\Gamma$ are the inner products in the Hilbert spaces $L_2(\Omega)$ and $L_2(\Gamma)$ of functions square integrable over Ω and Γ respectively.

Here, A^+ denotes the differential operator that is formally adjoint to A with respect to $(\cdot, \cdot)_\Omega$. Moreover, $C_{k,j}^+$ and $Q_{k,j}^+$ denote the tangent differential operators that are formally adjoint to $C_{k,j}$ and $Q_{k,j}$ respectively relative to $(\cdot, \cdot)_\Gamma$, the tangent differential operators $Q_{k,j}$ appearing in the representation of the boundary differential operators B_j in the form

$$B_j(x, D) = \sum_{k=1}^{2q} Q_{j,k}(x, D_\tau) D_\nu^{k-1}.$$

(Of course, if $k > m_j$, then $Q_{j,k} = 0$). Finally,

$$K_j := K_j(x, D) := \sum_{k=1}^{2q-j+1} T_{j,k}(x, D_\tau) D_\nu^{k-1},$$

where every $T_{j,k}(D_\tau)$ is a certain tangent differential operator on Γ of the order $\text{ord } T_{j,k} \leq 2q + 1 - j - k$.

In view of the Green formula, we consider the following boundary-value problem in Ω with $q + \varkappa$ additional unknown function on Γ :

$$(4) \quad A^+ w = \omega \quad \text{in } \Omega,$$

$$(5) \quad K_j w + \sum_{k=1}^{q+\varkappa} Q_{k,j}^+ h_k = \chi_j \quad \text{on } \Gamma, \quad j = 1, \dots, 2q,$$

$$(6) \quad \sum_{k=1}^{q+\varkappa} C_{k,j}^+ h_k = \chi_{2q+j} \quad \text{on } \Gamma, \quad j = 1, \dots, \varkappa.$$

This problem is formally adjoint to the problem (1), (2) with respect to the Green formula. Note that the problem (1), (2) is elliptic if and only if the problem (4), (5), (6) is elliptic (see [12, Theorem 3.1.2]).

3. REFINED SCALES OF SPACES

We will investigate properties of the elliptic boundary-value problem (1), (2) in appropriate couples of Hilbert spaces that form the two-sided refined scale introduced in [20]. This scale is constructed on the base of some Hörmander isotropic inner product spaces. In this section, we give the definitions of these spaces and the scale and, moreover, discuss some of their properties. We follow [20] (see also [22, 24, Subsect. 4.2.2]).

We start with the definition of the Hörmander spaces $H^{s,\varphi}(\mathbb{R}^n)$, with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. Here and below, \mathcal{M} denotes the set of all Borel measurable functions $\varphi : [1, \infty) \rightarrow (0, \infty)$ such that both the functions φ and $1/\varphi$ are bounded on each compact

interval $[1, b]$, with $1 < b < \infty$, and that the function φ is slowly varying at infinity in the sense of J. Karamata [11], i.e., $\varphi(\lambda t)/\varphi(t) \rightarrow 1$ as $t \rightarrow \infty$ for each $\lambda > 0$.

Note that the slowly varying functions are well investigated and have various applications [5, 31]. The standard example of such functions is

$$\varphi(t) := (\log t)^{r_1} (\log \log t)^{r_2} \dots \underbrace{(\log \dots \log t)^{r_k}}_{k \text{ times}}, \quad t \gg 1,$$

where $k \in \mathbb{Z}$, with $k \geq 1$, and $r_1, \dots, r_k \in \mathbb{R}$ are arbitrary parameters.

Let $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. By definition, the complex linear space $H^{s,\varphi}(\mathbb{R}^n)$, with $n \geq 1$, consists of all tempered distributions w on \mathbb{R}^n that the Fourier transform \widehat{w} of w is locally Lebesgue integrable over \mathbb{R}^n and satisfies the condition

$$\int \langle \xi \rangle^{2s} \varphi^2(\langle \xi \rangle) |\widehat{w}(\xi)|^2 d\xi < \infty.$$

Here, the integral is taken over \mathbb{R}^n , and $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$. The inner product in this space is defined by the formula

$$(w_1, w_2)_{H^{s,\varphi}(\mathbb{R}^n)} := \int \langle \xi \rangle^{2s} \varphi^2(\langle \xi \rangle) \widehat{w}_1(\xi) \overline{\widehat{w}_2(\xi)} d\xi$$

and induces the norm $\|w\|_{H^{s,\varphi}(\mathbb{R}^n)} := (w, w)_{H^{s,\varphi}(\mathbb{R}^n)}^{1/2}$.

The space $H^{s,\varphi}(\mathbb{R}^n)$ is a special isotropic Hilbert case of the spaces $\mathcal{B}_{p,\mu}$ introduced and investigated by L. Hörmander [9, Sect. 2.2] (see also [10, Sect. 10.1]). Namely, $H^{s,\varphi}(\mathbb{R}^n) = \mathcal{B}_{p,\mu}$ provided that $p = 2$ and $\mu(\xi) = \langle \xi \rangle^s \varphi(\langle \xi \rangle)$ for all $\xi \in \mathbb{R}^n$. The Hilbert spaces $\mathcal{B}_{2,\mu}$ were also investigated by L. R. Volevich and B. P. Paneah [33, § 2].

In the special case where $\varphi \equiv 1$, the space $H^{s,\varphi}(\mathbb{R}^n)$ becomes the inner product Sobolev space $H^s(\mathbb{R}^n)$ of order $s \in \mathbb{R}$. Generally, we have the continuous and dense embeddings

$$(7) \quad H^{s+\varepsilon}(\mathbb{R}^n) \hookrightarrow H^{s,\varphi}(\mathbb{R}^n) \hookrightarrow H^{s-\varepsilon}(\mathbb{R}^n) \quad \text{for every } \varepsilon > 0.$$

This means that, in the class of Hilbert separable spaces $\{H^{s,\varphi}(\mathbb{R}^n) : s \in \mathbb{R}, \varphi \in \mathcal{M}\}$, the function parameter φ refines the main regularity (of distributions) characterized by the number s . This class is selected by V. A. Mikhailets and A. A. Murach [16] and is called the refined Sobolev scale over \mathbb{R}^n (see [24, Subsect. 1.3.3]).

We need analogs of the space $H^{s,\varphi}(\mathbb{R}^n)$ for Ω and Γ . They are constructed in a standard way. Let us give the corresponding definitions [22, 24, Sect. 2.1 and 3.1].

By definition, the linear space $H^{s,\varphi}(\Omega)$ consists of the restrictions $u = w \upharpoonright \Omega$ of all distributions $w \in H^{s,\varphi}(\mathbb{R}^n)$ to Ω . It is endowed with the norm

$$\|u\|_{H^{s,\varphi}(\Omega)} := \inf \{ \|w\|_{H^{s,\varphi}(\mathbb{R}^n)} : w \in H^{s,\varphi}(\mathbb{R}^n), u = w \upharpoonright \Omega \}.$$

The space $H^{s,\varphi}(\Omega)$ is Hilbert and separable with respect to this norm. The set $C^\infty(\overline{\Omega})$ is dense in this space.

The space $H^{s,\varphi}(\Gamma)$ consists of all distributions on Γ that belong, in local coordinates, to $H^{s,\varphi}(\mathbb{R}^{n-1})$. Let us give a detailed definition. From C^∞ -structure on Γ , we arbitrarily choose a finite collection of local charts $\alpha_j : \mathbb{R}^{n-1} \leftrightarrow \Gamma_j$, with $j = 1, \dots, \lambda$, that the open sets $\Gamma_1, \dots, \Gamma_\lambda$ form a covering of the manifold Γ . Moreover, we arbitrarily choose functions $\chi_j \in C^\infty(\Gamma)$, with $j = 1, \dots, \lambda$, that form a partition of unity on Γ which satisfies the condition $\text{supp } \chi_j \subset \Gamma_j$. Then, by definition, the linear space $H^{s,\varphi}(\Gamma)$ consists of all distributions h on Γ such that $(\chi_j h) \circ \alpha_j \in H^{s,\varphi}(\mathbb{R}^{n-1})$ for every $j \in \{1, \dots, \lambda\}$. Here, $(\chi_j h) \circ \alpha_j$ is the representation of $\chi_j h$ in the local chart α_j . This space

is endowed with the inner product

$$(h_1, h_2)_{H^{s,\varphi}(\Gamma)} := \sum_{j=1}^{\lambda} ((\chi_j h_1) \circ \alpha_j, (\chi_j h_2) \circ \alpha_j)_{H^{s,\varphi}(\mathbb{R}^{n-1})}.$$

The space $H^{s,\varphi}(\Gamma)$ is Hilbert and separable. It does not depend (up to equivalence of norms) on the indicated choice of the local charts and partition of unity [22, 24, Theorem 2.3]. The set $C^\infty(\Gamma)$ is dense in $H^{s,\varphi}(\Gamma)$.

The spaces $H^{s,\varphi}(\Omega)$ and $H^{s,\varphi}(\Gamma)$, with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$, form the refined Sobolev scales over Ω and Γ respectively. In the case of $\varphi(t) \equiv 1$, these spaces become the inner product Sobolev spaces.

Note the following connection between these spaces [22, 24, Subsect. 3.2.1]. If $s > 1/2$, then the trace mapping $u \mapsto u \upharpoonright \Gamma$, with $u \in C^\infty(\Gamma)$, extends uniquely (by continuity) to a bounded surjective operator $R_\Gamma : H^{s,\varphi}(\Omega) \rightarrow H^{s-1/2,\varphi}(\Gamma)$. Thus, for each distribution $u \in H^{s,\varphi}(\Omega)$, its trace $R_\Gamma u$ on Γ is well defined provided that $s > 1/2$. But it is impossible to define this trace reasonably in the $s < 1/2$ case. Hence, we cannot investigate the boundary-value problem (1), (2) with $u \in H^{s,\varphi}(\Omega)$, where s is an arbitrary real number.

Therefore, we use a certain modification of the space $H^{s,\varphi}(\Omega)$ to be able to investigate this problem for arbitrary real s . This modification is denoted by $H^{s,\varphi,(2q)}(\Omega)$ and introduced in [20] by analogy with Ya. A. Roitberg's [27, 28] construction applied to the Sobolev scale (see also monographs [4, Chapt. III, Sect. 6] and [29, Sect. 2.1]).

Firstly, let us define the space $H^{s,\varphi,(0)}(\Omega)$. We put $H^{s,\varphi,(0)}(\Omega) := H^{s,\varphi}(\Omega)$ in the case of $s \geq 0$. But, in the case of $s < 0$, we let $H^{s,\varphi,(0)}(\Omega)$ define the dual Hilbert space to $H^{-s,1/\varphi}(\Omega)$ with respect to the inner product in $L_2(\Omega)$. Namely, if $s < 0$, then $H^{s,\varphi,(0)}(\Omega)$ is the completion of $C^\infty(\overline{\Omega})$ with respect to the Hilbert norm

$$\|u\|_{H^{s,\varphi,(0)}(\Omega)} := \sup \left\{ \frac{|(u, w)_\Omega|}{\|w\|_{H^{-s,1/\varphi}(\Omega)}} : w \in H^{-s,1/\varphi}(\Omega), w \neq 0 \right\}.$$

Note that the mapping $u \mapsto \mathcal{O}u$, with $u \in C^\infty(\overline{\Omega})$, extends uniquely (by continuity) to an isometric isomorphism of the space $H^{s,\varphi,(0)}(\Omega)$, with $s < 0$, onto the subspace $\{w \in H^{s,\varphi}(\mathbb{R}^n) : \text{supp } w \subseteq \overline{\Omega}\}$ of $H^{s,\varphi}(\mathbb{R}^n)$. Here, $\mathcal{O}u := u$ on $\overline{\Omega}$, and $\mathcal{O}u := 0$ on $\mathbb{R}^n \setminus \overline{\Omega}$.

We can now define the Hilbert space $H^{s,\varphi,(2q)}(\Omega)$. Let $E_{2q} := \{k-1/2 : k = 1, \dots, 2q\}$. In the case of $s \in \mathbb{R} \setminus E_{2q}$, the space $H^{s,\varphi,(2q)}(\Omega)$ is defined to be the completion of $C^\infty(\overline{\Omega})$ with respect to the Hilbert norm

$$\|u\|_{H^{s,\varphi,(2q)}(\Omega)} := \left(\|u\|_{H^{s,\varphi,(0)}(\Omega)}^2 + \sum_{k=1}^{2q} \|(D_\nu^{k-1} u) \upharpoonright \Gamma\|_{H^{s-k+1/2,\varphi}(\Gamma)}^2 \right)^{1/2}.$$

In the case of $s \in E_{2q}$, the space $H^{s,\varphi,(2q)}(\Omega)$ is defined with the help of interpolation between Hilbert spaces, namely,

$$H^{s,\varphi,(2q)}(\Omega) := [H^{s-\varepsilon,\varphi,(2q)}(\Omega), H^{s+\varepsilon,\varphi,(2q)}(\Omega)]_{1/2}, \quad \text{with } 0 < \varepsilon < 1.$$

We will give the definition of this interpolation in Section 5. Note [20, Theorem 7.1] that the right-hand side of the last equality does not depend (up to equivalence of norms) on the choice of ε . In the Sobolev case of $\varphi \equiv 1$, the space $H^{s,\varphi,(2q)}(\Omega)$ was introduced by Ya. A. Roitberg [27, 28].

For arbitrary $s \in \mathbb{R}$, the linear mapping $T_{2q} : u \mapsto (u, u \upharpoonright \Gamma, \dots, (D_\nu^{2q-1} u) \upharpoonright \Gamma)$, with $u \in C^\infty(\overline{\Omega})$, extends uniquely (by continuity) to a bounded operator

$$T_{2q} : H^{s,\varphi,(2q)}(\Omega) \rightarrow H^{s,\varphi,(0)}(\Omega) \oplus \bigoplus_{k=1}^{2q} H^{s-k+1/2,\varphi}(\Gamma) =: \Pi_{s,\varphi,(2q)}(\Omega, \Gamma).$$

If $s \notin E_{2q}$, then this operator is isometric and then its range consists of all vectors

$$(u_0, u_1, \dots, u_{2q}) \in \Pi_{s, \varphi, (2q)}(\Omega, \Gamma)$$

such that $u_k = R_\Gamma D_\nu^{k-1} u_0$ for all integers $k \in \{1, \dots, 2q\}$ satisfying $s > k - 1/2$ (see [20, Theorem 4.2]).

The Hilbert spaces $H^{s, \varphi, (2q)}(\Omega)$ and $H^{s, \varphi, (0)}(\Omega)$ are separable. They form the refined scales

$$(8) \quad \{H^{s, \varphi, (2q)}(\Omega) : s \in \mathbb{R}, \varphi \in \mathcal{M}\} \quad \text{and} \quad \{H^{s, \varphi, (0)}(\Omega) : s \in \mathbb{R}, \varphi \in \mathcal{M}\}$$

over Ω . These scales are two-sided because the parameter s runs through the whole real line. Note that $H^{s, \varphi, (2q)}(\Omega) = H^{s, \varphi}(\Omega)$ for $s > 2q - 1/2$ and that $H^{s, \varphi, (0)}(\Omega) = H^{s, \varphi}(\Omega)$ for $s > -1/2$, the equalities of Hilbert spaces being fulfilled up to equivalence of norms [22, 24, Theorems 3.9 and 4.12].

In the Sobolev case of $\varphi \equiv 1$, we will omit φ in the designations of the spaces introduced in the paper. It follows from (7) that

$$(9) \quad H^{s+\varepsilon, (2q)}(\Omega) \hookrightarrow H^{s, \varphi, (2q)}(\Omega) \hookrightarrow H^{s-\varepsilon, (2q)}(\Omega),$$

$$(10) \quad H^{s+\varepsilon, (0)}(\Omega) \hookrightarrow H^{s, \varphi, (0)}(\Omega) \hookrightarrow H^{s-\varepsilon, (0)}(\Omega),$$

$$(11) \quad H^{s+\varepsilon}(\Gamma) \hookrightarrow H^{s, \varphi}(\Gamma) \hookrightarrow H^{s-\varepsilon}(\Gamma)$$

for every $\varepsilon > 0$. These embeddings are compact and dense.

4. THE MAIN RESULTS

Here, we will formulate our results concerning the properties of the elliptic boundary-value problem (1), (2) on the two-sided refined scales (8) and $\{H^{s, \varphi}(\Gamma) : s \in \mathbb{R}, \varphi \in \mathcal{M}\}$.

With the problem (1), (2) and its formally adjoint problem (4), (5), (6), we associate the linear spaces N and N^+ of infinitely smooth solutions to these problems in the homogeneous case. Namely, let N consist of all solutions $(u, v_1, \dots, v_\varkappa) \in C^\infty(\overline{\Omega}) \times (C^\infty(\Gamma))^\varkappa$ to the problem (1), (2) in the case where $f = 0$ in Ω and all $g_j = 0$ on Γ . Besides, let N^+ consist of all solutions $(w, h_1, \dots, h_{q+\varkappa}) \in C^\infty(\overline{\Omega}) \times (C^\infty(\Gamma))^{q+\varkappa}$ to the problem (4), (5), (6) in the case where $\omega = 0$ in Ω and all $\chi_j = 0$ and all $\chi_{2q+j} = 0$ on Γ . Since these problems are elliptic in Ω , the spaces N and N^+ are finite-dimensional [12, Lemma 3.4.2].

Theorem 1. *For arbitrary $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$, the mapping (3) extends uniquely (by continuity) to a bounded operator*

$$(12) \quad \Lambda : \mathcal{D}^{s, \varphi, (2q)}(\Omega, \Gamma) := H^{s, \varphi, (2q)}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{s+r_k-1/2, \varphi}(\Gamma) \\ \rightarrow H^{s-2q, \varphi, (0)}(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2, \varphi}(\Gamma) =: \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma).$$

This operator is Fredholm. Its kernel coincides with N , and its range consists of all vectors

$$(13) \quad (f, g) := (f, g_1, \dots, g_{q+\varkappa}) \in \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma)$$

such that

$$(14) \quad (f, w)_\Omega + \sum_{j=1}^{q+\varkappa} (g_j, h_j)_\Gamma = 0 \quad \text{for each} \quad (w, h_1, \dots, h_{q+\varkappa}) \in N^+.$$

The index of the operator (12) is equal to $\dim N - \dim N^+$ and does not depend on s and φ .

Here and below, we let $(\cdot, \cdot)_\Omega$ and $(\cdot, \cdot)_\Gamma$ denote the extension by continuity of the inner products in the Hilbert spaces $L_2(\Omega)$ and $L_2(\Gamma)$ respectively.

In connection with this theorem, we recall that a linear bounded operator $T : E_1 \rightarrow E_2$ between some Banach spaces E_1 and E_2 is said to be Fredholm if its kernel $\ker T$ and co-kernel $E_2/T(E_1)$ are finite-dimensional. If this operator is Fredholm, then its range $T(E_1)$ is closed in E_2 , and its index $\text{ind } T := \dim \ker T - \dim(E_2/T(E_1))$ is finite.

It is useful to note that, in the case of $s > 2q - 1/2$, the bounded and Fredholm operator (12) acts between the spaces

$$\Lambda : H^{s,\varphi}(\Omega) \oplus \bigoplus_{k=1}^{\varkappa} H^{s+r_k-1/2,\varphi}(\Gamma) \rightarrow H^{s-2q,\varphi}(\Omega) \oplus \bigoplus_{j=1}^{q+\varkappa} H^{s-m_j-1/2,\varphi}(\Gamma),$$

which consist of distributions on Ω or Γ .

In the case where $N = \{0\}$ and $N^+ = \{0\}$, the operator (12) is an isomorphism of the space $\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)$ onto the space $\mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma)$. Generally, this operator induces an isomorphism between some of their (closed) subspaces, which have finite co-dimension. In this connection, we consider the following decompositions of these spaces in direct sums of subspaces:

$$(15) \quad \begin{aligned} \mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma) &= N \dot{+} \left\{ (u, v_1, \dots, v_\varkappa) \in \mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma) : \right. \\ &\left. (u_0, u^{(0)})_\Omega + \sum_{k=1}^{\varkappa} (v_k, v_k^{(0)})_\Gamma = 0 \text{ for all } (u^{(0)}, v_1^{(0)}, \dots, v_\varkappa^{(0)}) \in N \right\} \end{aligned}$$

with u_0 being the initial component of the vector $T_{2q}u = (u_0, u_1, \dots, u_{2q})$, and

$$(16) \quad \mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma) = N^+ \dot{+} \Lambda(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)).$$

Each decomposition is well defined because the subspaces have the intersection $\{0\}$ and because the finite dimension of the first subspace is equal to the co-dimension of the second. Let P and P^+ denote respectively the projectors of the spaces $\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)$ and $\mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma)$ onto the second term in the sums (15) and (16) parallel to the first term. These projectors do not depend, as mappings, on s and φ .

Theorem 2. *For arbitrary $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$, the restriction of the mapping (12) to the subspace $P(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma))$ is an isomorphism*

$$(17) \quad \Lambda : P(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)) \leftrightarrow P^+(\mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma)).$$

We will discuss properties of the generalized solutions to the elliptic boundary-value problem (1), (2). Beforehand, we give the definition of such solutions.

Let $\mathcal{D}^{-\infty,(2q)}(\Omega, \Gamma)$ denote the union of all the spaces $\mathcal{D}^{s,\varphi,(2q)}(\Omega)$ with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. Similarly, let $\mathcal{E}^{-\infty,(0)}(\Omega, \Gamma)$ denote the union of all the spaces $\mathcal{E}^{s,\varphi,(0)}(\Omega)$ with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. The linear spaces $\mathcal{D}^{-\infty,(2q)}(\Omega, \Gamma)$ and $\mathcal{E}^{-\infty,(0)}(\Omega, \Gamma)$ are well defined in view of the embeddings (9)–(11) and are endowed with the topology of inductive limit.

Consider an arbitrary vector

$$(18) \quad (u, v) := (u, v_1, \dots, v_\varkappa) \in \mathcal{D}^{-\infty,(2q)}(\Omega, \Gamma).$$

Since $(u, v) \in \mathcal{D}^{s,\varphi,(2q)}(\Omega)$ for some $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$, its image (13) is defined by the formula $\Lambda(u, v) = (f, g)$ by means of the bounded operator (12). The image (f, g) does not depend on $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. The vector (18) is said to be a generalized solution (in the sense of Ya. A. Roitberg) of the boundary-value problem (1), (2) with the right-hand sides (13). Such a solution satisfies the following a priori estimate.

Theorem 3. *Let $s \in \mathbb{R}$, $\varphi \in \mathcal{M}$, and a real number $\sigma > 0$. Then there exists a number $c > 0$ such that*

$$(19) \quad \|(u, v)\|_{\mathcal{D}^{s, \varphi, (2q)}(\Omega, \Gamma)} \leq c \left(\|\Lambda(u, v)\|_{\mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma)} + \|(u, v)\|_{\mathcal{D}^{s-\sigma, \varphi, (2q)}(\Omega, \Gamma)} \right)$$

for an arbitrary vector $(u, v) \in \mathcal{D}^{s, \varphi, (2q)}(\Omega, \Gamma)$. Here, the number $c = c(s, \varphi)$ does not depend on (u, v) .

Let us now focus our attention on the regularity properties of the generalized solutions to the elliptic problem under investigation. Let V be an arbitrary open subset of \mathbb{R}^n . We put $\Omega_0 := \Omega \cap V \neq \emptyset$ and $\Gamma_0 := \Gamma \cap V$, the $\Gamma_0 = \emptyset$ case being possible. Introduce a local analogs of the spaces $\mathcal{D}^{s, \varphi, (2q)}(\Omega)$ and $\mathcal{E}^{s, \varphi, (0)}(\Omega)$, with $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. We let $\mathcal{D}_{\text{loc}}^{s, \varphi, (2q)}(\Omega_0, \Gamma_0)$ denote the linear space of all vectors (18) such that $\chi(u, v) \in \mathcal{D}^{s, \varphi, (2q)}(\Omega)$ for an arbitrary function $\chi \in C^\infty(\bar{\Omega})$ with $\text{supp } \chi \subset \Omega_0 \cup \Gamma_0$. Similarly, we let $\mathcal{E}_{\text{loc}}^{s, \varphi, (0)}(\Omega_0, \Gamma_0)$ denote the linear space of all vectors $(f, g) \in \mathcal{E}^{-\infty, (0)}(\Omega, \Gamma)$ such that $\chi(f, g) \in \mathcal{E}^{s, \varphi, (0)}(\Omega)$ for an arbitrary function χ indicated just above. Of course, the multiplication of the vectors by the scalar function χ is understood to be component-wise, the products χu and χf are well defined by closure, and the product χh , where h is a distribution on Γ , means $(\chi \upharpoonright \Gamma)h$. We introduce topologies in the spaces $\mathcal{D}_{\text{loc}}^{s, \varphi, (2q)}(\Omega_0, \Gamma_0)$ and $\mathcal{E}_{\text{loc}}^{s, \varphi, (0)}(\Omega_0, \Gamma_0)$ by means of the seminorms $(u, v) \mapsto \|\chi(u, v)\|_{\mathcal{D}^{s, \varphi, (2q)}(\Omega)}$ and $(f, g) \mapsto \|\chi(f, g)\|_{\mathcal{E}^{s, \varphi, (0)}(\Omega)}$ respectively, where χ is an arbitrary function from the definitions of these spaces.

Theorem 4. *Let a vector $(u, v) \in \mathcal{D}^{-\infty, (2q)}(\Omega, \Gamma)$ be a generalized solution to the elliptic problem (1), (2) whose right-hand sides satisfy the condition $(f, g) \in \mathcal{E}_{\text{loc}}^{s-2q, \varphi, (0)}(\Omega_0, \Gamma_0)$ for some parameters $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. Then $(u, v) \in \mathcal{D}_{\text{loc}}^{s, \varphi, (2q)}(\Omega_0, \Gamma_0)$.*

We can see that the solution inherits the (supplementary) regularity φ of the right-hand sides of the elliptic problem.

In the case where $\Omega_0 = \Omega$ and $\Gamma_0 = \Gamma$, we have the equalities $\mathcal{D}_{\text{loc}}^{s, \varphi, (2q)}(\Omega_0, \Gamma_0) = \mathcal{D}^{s, \varphi, (2q)}(\Omega, \Gamma)$ and $\mathcal{E}_{\text{loc}}^{s-2q, \varphi, (0)}(\Omega_0, \Gamma_0) = \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma)$. Therefore, Theorem 4 says about the global regularity in this case, i.e., the regularity on the whole closed domain $\bar{\Omega}$.

If $\Gamma_0 = \emptyset$, then this theorem states that the local regularity of the component u of the generalized solution increases in neighborhoods of internal points of the domain Ω .

In the Sobolev case of $\varphi(t) \equiv 1$, Theorem 1–4 are proved in monographs [12, Sect. 3.2 and 3.4] (for an arbitrary integer s) and [30, Sect. 2.4] (for arbitrary real s and general elliptic systems). For arbitrary $\varphi \in \mathcal{M}$ and real $s > 2q$, Theorems 1 and 2 with analogs of Theorems 3 and 4 are proved in [7].

5. AUXILIARY RESULTS

The refined scales of spaces introduced in Section 3 possess an important interpolation property, which will play a key role in the proof of Theorem 1. Namely, each of these spaces can be obtained by the interpolation, with a function parameter, of an appropriate couple of Hilbert spaces corresponding to the Sobolev case of $\varphi(t) \equiv 1$. Therefore, we will recall the definition of this interpolation in the case of arbitrary Hilbert spaces and discuss some of its properties that are necessary for us. Among them, we will formulate the mentioned interpolation property for Hilbert spaces appearing in (12).

The method of interpolation with a function parameter of Hilbert spaces was introduced by C. Foias and J.-L. Lions in [6, p. 278]. We follow monographs [22, 24, Sect. 1.1], which systematically expound this method. It is sufficient for our purposes to restrict ourselves to separable Hilbert spaces.

Let $X := [X_0, X_1]$ be a given ordered couple of separable complex Hilbert spaces X_0 and X_1 such that the continuous and dense embedding $X_1 \hookrightarrow X_0$ holds. This couple is called admissible. For X , there exists a self-adjoint positive-definite operator J on X_0 which has the domain X_1 and satisfies the condition $\|Jw\|_{X_0} = \|w\|_{X_1}$ for all $w \in X_1$. This operator is uniquely determined by the couple X and is called a generating operator for X . It sets an isometric isomorphism $J : X_1 \leftrightarrow X_0$.

Let \mathcal{B} denote the set of all Borel measurable functions $\psi : (0, \infty) \rightarrow (0, \infty)$ such that ψ is bounded on each compact interval $[a, b]$, with $0 < a < b < \infty$, and that $1/\psi$ is bounded on every set $[r, \infty)$, with $r > 0$.

For arbitrary $\psi \in \mathcal{B}$, we consider the (generally, unbounded) operator $\psi(J)$, which is defined on X_0 as the Borel function ψ of J . Let $[X_0, X_1]_\psi$ or, simply, X_ψ denote the domain of the operator $\psi(J)$ endowed with the inner product $(u_1, u_2)_{X_\psi} := (\psi(J)u_1, \psi(J)u_2)_{X_0}$ and the corresponding norm $\|u\|_{X_\psi} = \|\psi(J)u\|_{X_0}$. The space X_ψ is Hilbert and separable.

A function $\psi \in \mathcal{B}$ is called an interpolation parameter if the following condition is fulfilled for all admissible couples $X = [X_0, X_1]$ and $Y = [Y_0, Y_1]$ of Hilbert spaces and for an arbitrary linear mapping T given on X_0 : if the restriction of T to X_j is a bounded operator $T : X_j \rightarrow Y_j$ for each $j \in \{0, 1\}$, then the restriction of T to X_ψ is also a bounded operator $T : X_\psi \rightarrow Y_\psi$. In this case we say that X_ψ is obtained by the interpolation with the function parameter ψ of the couple X .

The function $\psi \in \mathcal{B}$ is an interpolation parameter if and only if ψ is pseudoconcave on a neighborhood of infinity, i.e. $\psi(t) \asymp \psi_1(t)$ with $t \gg 1$ for a certain positive concave function $\psi_1(t)$. (As usual, the designation $\psi \asymp \psi_1$ means that both the functions ψ/ψ_1 and ψ_1/ψ are bounded on the indicated set). This fundamental fact follows from J. Peetre's [26] description of all interpolation functions of positive order. Specifically, the power function $\psi(t) \equiv t^s$ is an interpolation parameter if and only if $0 \leq s \leq 1$. In this case, the interpolation space X_ψ is denoted by X_s .

We need the following interpolation properties of the refined scales [22, 24, Theorems 1.14, 2.2, 3.10, and 4.22].

Proposition 1. *Let a function $\varphi \in \mathcal{M}$ and positive real numbers ε, δ be given. Define a function $\psi \in \mathcal{B}$ by the formulas $\psi(t) := t^{\varepsilon/(\varepsilon+\delta)}\varphi(t^{1/(\varepsilon+\delta)})$ for $t > 1$, and $\psi(t) := \varphi(1)$ if $0 < t < 1$. Then ψ is an interpolation parameter, and, for arbitrary $\sigma \in \mathbb{R}$, we have*

$$\begin{aligned} [H^{\sigma-\varepsilon, (2q)}(\Omega), H^{\sigma+\delta, (2q)}(\Omega)]_\psi &= H^{\sigma, \varphi, (2q)}(\Omega), \\ [H^{\sigma-\varepsilon, (0)}(\Omega), H^{\sigma+\delta, (0)}(\Omega)]_\psi &= H^{\sigma, \varphi, (0)}(\Omega), \\ [H^{\sigma-\varepsilon}(\Gamma), H^{\sigma+\delta}(\Gamma)]_\psi &= H^{\sigma, \varphi}(\Gamma) \end{aligned}$$

up to equivalence of norms.

We will also use the following two general properties of the interpolation [22, 24, Theorems 1.7 and 1.5].

Proposition 2. *Let $X = [X_0, X_1]$ and $Y = [Y_0, Y_1]$ be admissible couples of Hilbert spaces, and let a linear mapping T be given on X_0 . Suppose that we have the bounded and Fredholm operators $T : X_j \rightarrow Y_j$, with $j = 0, 1$, that possess the common kernel and the common index. Then, for an arbitrary interpolation parameter $\psi \in \mathcal{B}$, the bounded operator $T : X_\psi \rightarrow Y_\psi$ is Fredholm, has the same kernel and the same index, and, moreover, its range $T(X_\psi) = Y_\psi \cap T(X_0)$.*

Proposition 3. *Let $[X_0^{(j)}, X_1^{(j)}]$, with $j = 1, \dots, r$, be a finite collection of admissible couples of Hilbert spaces. Then, for every function $\psi \in \mathcal{B}$, we have*

$$\left[\bigoplus_{j=1}^r X_0^{(j)}, \bigoplus_{j=1}^r X_1^{(j)} \right]_{\psi} = \bigoplus_{j=1}^r [X_0^{(j)}, X_1^{(j)}]_{\psi}$$

with equality of norms.

6. PROOFS OF THE MAIN RESULTS

Let us prove Theorems 1–4.

Proof of Theorem 1. In the Sobolev case of $\varphi(t) \equiv 1$, this theorem is proved in [12, Theorem 3.4.1] (for integer-valued s) and [30, Theorem 2.4.1] (for real s and elliptic systems). We will deduce Theorem 1 from the Sobolev case with the help of the interpolation with a function parameter.

Let $s \in \mathbb{R}$ and $\varphi \in \mathcal{M}$. We choose positive numbers ε and δ such that $s - \varepsilon$ and $s + \delta$ are integers. According to [12, Theorem 3.4.1] the mapping (3) extends by continuity to the bounded and Fredholm operators

$$(20) \quad \Lambda : \mathcal{D}^{s-\varepsilon, (2q)}(\Omega, \Gamma) \rightarrow \mathcal{E}^{s-\varepsilon-2q, (0)}(\Omega, \Gamma), \quad \Lambda : \mathcal{D}^{s+\delta, (2q)}(\Omega, \Gamma) \rightarrow \mathcal{E}^{s+\delta-2q, (0)}(\Omega, \Gamma).$$

(Recall that we omit the index φ in the designations of the spaces if $\varphi \equiv 1$.) These operators have the common kernel N and the same index equaled to $\dim N - \dim N^+$. The range of the first operator satisfies the equality

$$(21) \quad \Lambda(\mathcal{D}^{s-\varepsilon, (2q)}(\Omega, \Gamma)) = \{(f, g) \in \mathcal{E}^{s-\varepsilon-2q, (0)}(\Omega, \Gamma) : (14) \text{ is true}\}$$

and analogous formula holds for the second operator.

Let us apply the interpolation with the function parameter ψ from Proposition 1 to (20). According to Proposition 2, we get the bounded and Fredholm operator

$$(22) \quad \Lambda : [\mathcal{D}^{s-\varepsilon, (2q)}(\Omega, \Gamma), \mathcal{D}^{s+\delta, (2q)}(\Omega, \Gamma)]_{\psi} \rightarrow [\mathcal{E}^{s-\varepsilon-2q, (0)}(\Omega, \Gamma), \mathcal{E}^{s+\delta-2q, (0)}(\Omega, \Gamma)]_{\psi}.$$

This operator is an extension of the mapping (3) by continuity.

Let us describe the interpolation spaces appearing in (22). Applying Propositions 3 and 1 successively, we obtain

$$\begin{aligned} & [\mathcal{D}^{s-\varepsilon, (2q)}(\Omega, \Gamma), \mathcal{D}^{s+\delta, (2q)}(\Omega, \Gamma)]_{\psi} \\ &= [H^{s-\varepsilon, (2q)}(\Omega), H^{s+\delta, (2q)}(\Omega)]_{\psi} \oplus \bigoplus_{k=1}^{\infty} [H^{s+r_k-1/2-\varepsilon}(\Gamma), H^{s+r_k-1/2+\delta}(\Gamma)]_{\psi} \\ &= H^{s, \varphi, (2q)}(\Omega) \oplus \bigoplus_{k=1}^{\infty} H^{s+r_k-1/2, \varphi}(\Gamma) = \mathcal{D}^{s, \varphi, (2q)}(\Omega, \Gamma). \end{aligned}$$

Analogously,

$$[\mathcal{E}^{s-\varepsilon-2q, (0)}(\Omega, \Gamma), \mathcal{E}^{s+\delta-2q, (0)}(\Omega, \Gamma)]_{\psi} = \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma).$$

These equalities of Hilbert spaces are fulfilled up to equivalence of norms.

Thus, the bounded and Fredholm operator (22) is the operator (12) from Theorem 1. By virtue of Proposition 2, the kernel and index of the operator (12) coincide respectively with the common kernel N and the common index $\dim N - \dim N^+$ of the operators (20). Moreover, the range of (12) is equal to

$$\mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma) \cap \Lambda(\mathcal{D}^{s-\varepsilon, (2q)}(\Omega, \Gamma)) = \{(f, g) \in \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma) : (14) \text{ is true}\}$$

in view of (21). \square

Proof of Theorem 2. According to Theorem 1, the restriction of the operator (12) to $P(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma))$ is a continuous and one-to-one mapping from the whole subspace $P(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma))$ to the whole subspace $P^+(\mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma))$. Then, by the Banach theorem on inverse operator, this linear mapping is the isomorphism (17). \square

Proof of Theorem 3. Choose vector $(u, v) \in \mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)$ arbitrarily. According to Theorem 2, we have the inequalities

$$(23) \quad \begin{aligned} & \|(u, v)\|_{\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)} \\ & \leq \|P(u, v)\|_{\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)} + \|(u, v) - P(u, v)\|_{\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)} \\ & \leq c_1 \|\Lambda P(u, v)\|_{\mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma)} + c_2 \|(u, v) - P(u, v)\|_{\mathcal{D}^{s-\sigma,\varphi,(2q)}(\Omega, \Gamma)}. \end{aligned}$$

Here, c_1 is the norm of the inverse operator to the isomorphism (17), whereas c_2 is a certain positive number not depending on (u, v) . This number exists because the vector $(u, v) - P(u, v)$ belongs to the finite-dimensional space N , where all norms are equivalent, specifically, the norms in $\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)$ and $\mathcal{D}^{s-\sigma,\varphi,(2q)}(\Omega, \Gamma)$. Besides,

$$(24) \quad \|P(u, v)\|_{\mathcal{D}^{s-\sigma,\varphi,(2q)}(\Omega, \Gamma)} \leq c_3 \|(u, v)\|_{\mathcal{D}^{s-\sigma,\varphi,(2q)}(\Omega, \Gamma)},$$

where c_3 is the norm of the projector P acting on the space $\mathcal{D}^{s-\sigma,\varphi,(2q)}(\Omega, \Gamma)$. Now, the formulas (23) and (24) directly imply the required estimate (19). \square

Proof of Theorem 4. First, let us prove this theorem in the case of global regularity, where $\Omega_0 = \Omega$ and $\Gamma_0 = \Gamma$. According to Theorem 1, the vector $(f, g) := \Lambda(u, v)$ satisfies (14). Therefore, by virtue of the same theorem and the condition $(f, g) \in \mathcal{E}^{s-2q,\varphi,(0)}(\Omega, \Gamma)$, we get the inclusion $(f, g) \in \Lambda(\mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma))$. Hence, together with $\Lambda(u, v) = (f, g)$, we have the equality $\Lambda(u', v') = (f, g)$ for a certain vector $(u', v') \in \mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma)$. Therefore, $\Lambda(u - u', v - v') = 0$, that implies

$$(u - u', v - v') \in N \subset C^\infty(\bar{\Omega}) \times (C^\infty(\Gamma))^\varkappa$$

in view of Theorem 1. Thus,

$$(u, v) = (u', v') + (u - u', v - v') \in \mathcal{D}^{s,\varphi,(2q)}(\Omega, \Gamma).$$

Theorem 4 is proved in the case of global regularity.

We now pass to the general case of local regularity and deduce the theorem in this case from the specific case of global regularity just considered. We will previously prove that the following implication holds true for each $p \geq 1$:

$$(25) \quad (u, v) \in \mathcal{D}_{\text{loc}}^{s-p,\varphi,(2q)}(\Omega_0, \Gamma_0) \Rightarrow (u, v) \in \mathcal{D}_{\text{loc}}^{s-p+1,\varphi,(2q)}(\Omega_0, \Gamma_0)$$

under the conditions of the theorem.

Suppose that the premise of this implication is valid for a certain number $p \geq 1$. We arbitrarily choose a function $\chi \in C^\infty(\bar{\Omega})$ such that $\text{supp } \chi \subset \Omega_0 \cup \Gamma_0$. We also choose a certain function $\eta \in C^\infty(\bar{\Omega})$ that satisfies the conditions $\text{supp } \eta \subset \Omega_0 \cup \Gamma_0$ and $\eta = 1$ on a neighborhood of $\text{supp } \chi$.

Interchanging the operator of the multiplication by χ with each of the differential operators A , B_j , and $C_{j,k}$, we can write down the following equalities:

$$(26) \quad A(\chi u) = A(\chi \eta u) = \chi A(\eta u) + A'(\eta u) = \chi A u + A'(\eta u),$$

$$(27) \quad B_j(\chi u) = B_j(\chi \eta u) = \chi B_j(\eta u) + B'_j(\eta u) = \chi B_j u + B'_j(\eta u),$$

$$(28) \quad C_{j,k}(\chi v_k) = C_{j,k}(\chi \eta v_k) = \chi C_{j,k}(\eta v_k) + C'_{j,k}(\eta v_k) = \chi C_{j,k} v_k + C'_{j,k}(\eta v_k).$$

Here, A' is a certain differential operator on $\bar{\Omega}$, B'_j is a certain boundary differential operator on Γ , and $C'_{j,k}$ is a certain tangent differential operator on Γ . These operators

are linear, all of their coefficients are infinitely smooth on the indicated sets, and the orders of these operators satisfy the conditions

$$(29) \quad \text{ord } A' \leq 2q - 1, \quad \text{ord } B'_j \leq m_j - 1, \quad \text{ord } C'_{j,k} \leq m_j + r_k - 1.$$

We put

$$\Lambda'(u, v) := \left(A'u, B'_1 u + \sum_{k=1}^{\infty} C'_{1,k} v_k, \dots, B'_{q+\infty} u + \sum_{k=1}^{\infty} C'_{q+\infty,k} v_k \right).$$

It follows from (26)–(28) that

$$(30) \quad \Lambda(\chi u, \chi v) = \chi \Lambda(u, v) + \Lambda'(\eta u, \eta v) = (\chi f, \chi g) + \Lambda'(\eta u, \eta v).$$

Here, by the condition of the theorem,

$$(31) \quad (\chi f, \chi g) \in \mathcal{E}^{s-2q, \varphi, (0)}(\Omega, \Gamma) \subseteq \mathcal{E}^{s-p+1-2q, \varphi, (0)}(\Omega, \Gamma).$$

According to the premise of the implication, we have $(\eta u, \eta v) \in \mathcal{D}^{s-p, \varphi, (2q)}(\Omega, \Gamma)$. It follows from the latter inclusion and the inequalities (29) that

$$(32) \quad \Lambda'(\eta u, \eta v) \in \mathcal{E}^{s-p-(2q-1), \varphi, (0)}(\Omega, \Gamma) = \mathcal{E}^{s-p+1-2q, \varphi, (0)}(\Omega, \Gamma)$$

in view of [22, 24, Theorem 4.13 and Lemma 2.5].

The formulas (30)–(32) gives the inclusion

$$(33) \quad \Lambda(\chi u, \chi v) \in \mathcal{E}^{s-p+1-2q, \varphi, (0)}(\Omega, \Gamma).$$

By virtue of Theorem 4 in the global case proved above, we conclude that (33) implies the inclusion $(\chi u, \chi v) \in \mathcal{D}^{s-p+1, \varphi, (2q)}(\Omega, \Gamma)$. In view of arbitrariness of the indicated choice of χ , this inclusion means that the inference of the implication (25) is true. Thus, we have proved this implication for each $p \geq 1$.

Now, using the implication (25), we can prove Theorem 4 in the general case. By the condition of this theorem, $(u, v) \in \mathcal{D}^{-\infty, (2q)}(\Omega, \Gamma)$. It follows from this inclusion, in view of (9) and (11), that

$$(u, v) \in \mathcal{D}^{s-\lambda, \varphi, (2q)}(\Omega, \Gamma) \subseteq \mathcal{D}_{\text{loc}}^{s-\lambda, \varphi, (2q)}(\Omega_0, \Gamma_0)$$

for a certain integer $\lambda \geq 1$. Using the implication (25) successively for values $p = \lambda$, $p = \lambda - 1, \dots$, and $p = 1$, we conclude that

$$\begin{aligned} (u, v) \in \mathcal{D}_{\text{loc}}^{s-\lambda, \varphi, (2q)}(\Omega_0, \Gamma_0) &\Rightarrow (u, v) \in \mathcal{D}_{\text{loc}}^{s-\lambda+1, \varphi, (2q)}(\Omega_0, \Gamma_0) \\ &\Rightarrow \dots \Rightarrow (u, v) \in \mathcal{D}_{\text{loc}}^{s-1, \varphi, (2q)}(\Omega_0, \Gamma_0) \Rightarrow (u, v) \in \mathcal{D}_{\text{loc}}^{s, \varphi, (2q)}(\Omega_0, \Gamma_0). \end{aligned}$$

Thus, we have proved Theorem 4 in the general case. \square

7. APPLICATIONS

In this section, we will apply Theorem 4 to obtain new sufficient conditions under which a generalized solution to the elliptic problem (1), (2) has continuous classical derivatives of a prescribed order, specifically, is a classical solution. To this end, we also use the following version [22, 24, Theorems 2.8 and 3.4] of the Hörmander embedding theorem [9, Theorem 2.2.7].

Proposition 4. *Let an integer $\ell \geq 0$ and function $\varphi \in \mathcal{M}$ be arbitrarily chosen. Then each of the embeddings $H^{\ell+n/2, \varphi}(\Omega) \subset C^\ell(\bar{\Omega})$ and $H^{\ell+(n-1)/2, \varphi}(\Gamma) \subset C^\ell(\Gamma)$ is equivalent to the condition*

$$(34) \quad \int_1^\infty \frac{dt}{t \varphi^2(t)} < \infty.$$

These embeddings are compact.

In connection with this proposition, we recall that, by the Sobolev embedding theorem, the embedding $H^\sigma(\Omega) \hookrightarrow C^\ell(\overline{\Omega})$ is equivalent to the condition $\sigma > \ell + n/2$. An analogous result is true for the Sobolev spaces over Γ . Thus, Proposition 4 refines, so to say, the Sobolev embedding theorem in the limiting case.

Applying Theorem 4 and Proposition 4, we can obtain the following result.

Theorem 5. *Let an integer ℓ meet the inequalities $\ell \geq 0$ and $\ell > 2q - (n+1)/2$. Suppose that the condition of Theorem 4 is fulfilled for $s = \ell + n/2$ and a certain function $\varphi \in \mathcal{M}$ that satisfies (34). Then $u \in C^\ell(\Omega_0 \cup \Gamma_0)$.*

Proof. Let a point $x_0 \in \Omega_0 \cup \Gamma_0$ be arbitrary. We choose a function $\chi \in C^\infty(\overline{\Omega})$ such that $\text{supp } \chi \subset \Omega_0 \cup \Gamma_0$ and $\chi = 1$ in a certain neighborhood V_0 of the point x_0 . (Of course, the neighborhood is considered in the topology of $\overline{\Omega}$.) According to Theorem 4, we have the inclusion $\chi u \in H^{\ell+n/2, \varphi, (2q)}(\Omega)$. Applying Proposition 4, we deduce that

$$\chi u \in H^{\ell+n/2, \varphi, (2q)}(\Omega) = H^{\ell+n/2, \varphi}(\Omega) \subset C^\ell(\overline{\Omega}).$$

Here, the equality holds true because $\ell + n/2 > 2q - 1/2$ by the condition of this theorem. Therefore, $u \in C^\ell(V_0)$. This means the inclusion $u \in C^\ell(\Omega_0 \cup \Gamma_0)$ in view of arbitrariness of the choice of $x_0 \in \Omega_0 \cup \Gamma_0$. \square

Remark 1. The conclusion of Theorem 5 remains valid if we omit the condition $\ell > 2q - (n+1)/2$ and additionally suppose that $u \in H^\sigma(\Omega)$ for certain $\sigma > 2q - 1/2$. Indeed, if $\ell \leq 2q - (n+1)/2$, then $H^\sigma(\Omega) \subset C^\ell(\overline{\Omega})$ by virtue of the Sobolev embedding theorem and the assumption $\sigma > 2q - 1/2$.

Let us formulate a version of Theorem 5 for the function v_k , with $k \in \{1, \dots, \varkappa\}$.

Theorem 6. *Let $\ell \in \mathbb{Z}$ and $\ell \geq 0$. Suppose that the condition of Theorem 4 is fulfilled for $s = \ell - r_k + n/2$ and a certain function $\varphi \in \mathcal{M}$ that satisfies (34). Then $v_k \in C^\ell(\Gamma_0)$.*

Proof. Let a point $x_0 \in \Gamma_0$ be arbitrary, and let a function χ and neighborhood V_0 be those as in the proof of Theorem 5. According to Theorem 4, we have the inclusion $\chi v_k \in H^{\ell+(n-1)/2, \varphi}(\Gamma)$. Hence, $\chi v_k \in C^\ell(\Gamma)$ by Proposition 4. Therefore, $v_k \in C^\ell(V_0 \cap \Gamma)$. This means the inclusion $v_k \in C^\ell(\Gamma_0)$ in view of arbitrariness of the choice of $x_0 \in \Gamma_0$. \square

Remark 2. It follows from Proposition 4 that the condition (34) not only is sufficient in Theorems 5 and 6 but also is necessary on the class of all the solutions considered.

Using Theorems 5 and 6 we can deduce the following sufficient condition under which a generalized solution (u, v) to the elliptic problem (1), (2) is classical, i.e., $u \in C^{2q}(\Omega) \cap C^m(\overline{\Omega})$ and $v_k \in C^{m+r_k}(\Gamma)$ for all $k \in \{1, \dots, \varkappa\}$. Here, we denote $m := \max\{m_1, \dots, m_{q+\varkappa}\}$.

Theorem 7. *Let $(u, v) \in \mathcal{D}^{-\infty, (2q)}(\Omega, \Gamma)$, and let $u \in H^\sigma(\Omega)$ for certain $\sigma > 2q - 1/2$. Suppose that the vector (u, v) is a solution to the elliptic problem (1), (2) whose right-hand sides satisfy the condition*

$$(f, g) \in \mathcal{E}_{\text{loc}}^{n/2, \varphi_1}(\Omega, \emptyset) \cap \mathcal{E}^{m+n/2-2q, \varphi_2}(\Omega)$$

for some functions $\varphi_1, \varphi_2 \in \mathcal{M}$ that meet the condition (34) with $\varphi := \varphi_1$ and $\varphi := \varphi_2$ respectively. Then the solution (u, v) is classical.

Proof. The inclusion $u \in C^{2q}(\Omega)$ follows from the condition $(f, g) \in \mathcal{E}_{\text{loc}}^{n/2, \varphi_1}(\Omega, \emptyset)$ by virtue of Theorem 5 with $\ell := 2q$, $\Omega_0 := \Omega$, and $\Gamma_0 := \emptyset$. Besides, the inclusion $u \in C^m(\overline{\Omega})$ follows from the condition $(f, g) \in \mathcal{E}^{m+n/2-2q, \varphi_2}(\Omega)$ in view of the same theorem with $\ell := m$, $\Omega_0 := \Omega$, and $\Gamma_0 := \Gamma$ if we take into account Remark 1. Finally, the inclusion $v_k \in C^{m+r_k}(\Gamma)$ for every $k \in \{1, \dots, \varkappa\}$ follows from the latter condition by virtue of Theorem 6 with $\ell := m + r_k$ and $\Gamma_0 := \Gamma$. \square

REFERENCES

1. A. V. Anop, A. A. Murach, *Parameter-elliptic problems and interpolation with a function parameter*, Methods Funct. Anal. Topology **20** (2014), no. 2, 103–116.
2. A. V. Anop, A. A. Murach, *Regular elliptic boundary-value problems in the extended Sobolev scale*, Ukrainian Math. J. **66** (2014), no. 7, 969–985.
3. A. G. Aslanyan, D. G. Vassiliev, V. B. Lidskii, *Frequencies of free oscillations of thin shell interacting with fluid*, Functional Anal. Appl. **15** (1981), no. 3, 157–164.
4. Ju. M. Berezanskii, *Expansions in Eigenfunctions of Selfadjoint Operators*, Amer. Math. Soc., Providence, RI, 1968. (Russian edition: Naukova Dumka, Kiev, 1965)
5. N. H. Bingham, C. M. Goldie, J. L. Teugels, *Regular Variation*, Cambridge University Press, Cambridge, 1989.
6. C. Foias, J.-L. Lions, *Sur certains théorèmes d'interpolation*, Acta Sci. Math. (Szeged) **22** (1961), no. 3–4, 269–282.
7. I. S. Chepurukhina, *On some classes of elliptic boundary-value problems in spaces of generalized smoothness*, Differential equations and related topics, Zb. prac' Inst. mat. NAN Ukr., Kyiv **11** (2014), no. 2, pp. 284–304. (Ukrainian)
8. P. G. Ciarlet, *Plates and Junctions in Elastic Multistructures. An Asymptotic Analysis*, Mayson, Paris, 1990.
9. L. Hörmander, *Linear Partial Differential Operators*, Springer, Berlin, 1963.
10. L. Hörmander, *The Analysis of Linear Partial Differential Operators*, vol. 2, Springer, Berlin, 1983.
11. J. Karamata, *Sur certains "Tauberian theorems" de M. M. Hardy et Littlewood*, Mathematica (Cluj) **3** (1930), 33–48.
12. V. A. Kozlov, V. G. Maz'ya, J. Rossmann, *Elliptic Boundary Value Problems in Domains with Point Singularities*, Amer. Math. Soc., Providence, RI, 1997.
13. B. Lawruk, *Parametric boundary-value problems for elliptic systems of linear differential equations. I. Construction of conjugate problems*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **11** (1963), no. 5, 257–267. (Russian)
14. B. Lawruk, *Parametric boundary-value problems for elliptic systems of linear differential equations. II. A boundary-value problem for a half-space*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **11** (1963), no. 5, 269–278. (Russian)
15. B. Lawruk, *Parametric boundary-value problems for elliptic systems of linear differential equations. III. Conjugate boundary problem for a half-space*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **13** (1965), no. 2, 105–110. (Russian)
16. V. A. Mikhailets, A. A. Murach, *Elliptic operators in a refined scale of function spaces*, Ukrainian Math. J. **57** (2005), no. 5, 817–825.
17. V. A. Mikhailets, A. A. Murach, *Refined scales of spaces and elliptic boundary-value problems. II*, Ukrainian Math. J. **58** (2006), no. 3, 398–417.
18. V. A. Mikhailets, A. A. Murach, *A regular elliptic boundary-value problem for a homogeneous equation in a two-sided refined scale of spaces*, Ukrainian Math. J. **58** (2006), no. 11, 1748–1767.
19. V. A. Mikhailets, A. A. Murach, *Refined scales of spaces and elliptic boundary-value problems. III*, Ukrainian Math. J. **59** (2007), no. 5, 744–765.
20. V. A. Mikhailets, A. A. Murach, *An elliptic boundary-value problem in a two-sided refined scale of spaces*, Ukrainian Math. J. **60** (2008), no. 4, 574–597.
21. V. A. Mikhailets, A. A. Murach, *Interpolation with a function parameter and refined scale of spaces*, Methods Funct. Anal. Topology **14** (2008), no. 1, 81–100.
22. V. A. Mikhailets, A. A. Murach, *Hörmander Spaces, Interpolation, and Elliptic Problems*, Proceedings of the Institute of Mathematics of NAS of Ukraine, Kyiv, vol. 84, 2010. (Russian); available at arXiv:1106.3214.
23. V. A. Mikhailets, A. A. Murach, *The refined Sobolev scale, interpolation, and elliptic problems*, Banach J. Math. Anal. **6** (2012), no. 2, 211–281.
24. V. A. Mikhailets, A. A. Murach, *Hörmander Spaces, Interpolation, and Elliptic Problems*, De Gruyter, Berlin, 2014.
25. S. Nazarov, K. Pileckas, *On noncompact free boundary problems for the plane stationary Navier–Stokes equations*, J. Reine Angew. Math. **438** (1993), 103–141.
26. J. Peetre, *On interpolation functions. II*, Acta Sci. Math. (Szeged) **29** (1968), no. 1–2, 91–92.
27. Ya. A. Roitberg, *Elliptic problems with nonhomogeneous boundary conditions and local increase of smoothness up to the boundary for generalized solutions*, Dokl. Math. **5** (1964), 1034–1038.
28. Ya. A. Roitberg, *A theorem on the homeomorphisms induced in L_p by elliptic operators and the local smoothing of generalized solutions*, Ukrain. Mat. Zh. **17** (1965), no. 5, 122–129. (Russian)

29. Ya. A. Roitberg, *Elliptic Boundary Value Problems in the Spaces of Distributions*, Kluwer Academic Publishers, Dordrecht, 1996.
30. Ya. A. Roitberg, *Boundary Value Problems in the Spaces of Distributions*, Kluwer Academic Publishers, Dordrecht, 1999.
31. E. Seneta, *Regularly Varying Functions*, Springer, Berlin, 1976.
32. G. Slenzak, *Elliptic problems in a refined scale of spaces*, Moscow Univ. Math. Bull. **29** (1974), no. 3–4, 80–88.
33. L. R. Volevich, B. P. Paneah, *Certain spaces of generalized functions and embedding theorems*, Uspekhi Mat. Nauk **20** (1965), no. 1, 3–74. (Russian)

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS'KA,
KYIV, 01601, UKRAINE

E-mail address: `Chepuruhina@mail.ru`

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS'KA,
KYIV, 01601, UKRAINE

E-mail address: `murach@imath.kiev.ua`

Received 25/11/2014