

FRACTIONAL CONTACT MODEL IN THE CONTINUUM

ANATOLY N. KOCHUBEI AND YURI G. KONDRATIEV

To Yu. M. Berezansky with deep respect and gratitude

ABSTRACT. We consider the evolution of correlation functions in a non-Markov version of the contact model in the continuum. The memory effects are introduced by assuming the fractional evolution equation for the statistical dynamics. This leads to a behavior of time-dependent correlation functions, essentially different from the one known for the standard contact model.

1. STATISTICAL DYNAMICS OF CONTINUOUS SYSTEMS

A traditional interpretation of a dynamics is given in terms of trajectories in a phase space of the considered system. This may be formulated as an evolution of microscopic states of the system. However such microscopic dynamics creates naturally an associated evolution of functions on the phase space (observables) and the dual evolution of probability measures on the phase space (macroscopic states). This is a classical triple of evolution equations in the Hamiltonian systems theory where the Newton description of particles dynamics may be reformulated by means of Liouville equations for observables or states.

If we are dealing with Markov stochastic processes, then we have to use the language of random trajectories for the stochastic dynamics in the phase space and there are two related evolution equations. Namely, we have corresponding dynamics of functions given by the Kolmogorov backward equation and related dynamics of measures given by the Kolmogorov forward or Fokker-Planck equation. In all cases, we have evolution equations for probability measures on phase spaces. These measures, or macroscopic states, describe statistical properties of considered systems and we will call corresponding evolution the statistical dynamics associated with the initial (random) dynamical system. This concept has sense even in the case when dynamics in terms of trajectories can not be constructed due a complicated character of considered models. We illustrate this situations in the framework of Markov evolutions of interacting particle systems in the continuum, see e.g. [4].

The phase space of our systems is the configuration space

$$\Gamma := \{ \gamma \subset \mathbb{R}^d \mid |\gamma \cap \Lambda| < \infty \text{ for any compact } \Lambda \subset \mathbb{R}^d \},$$

where $|A|$ denotes the cardinality of a set $A \subset \mathbb{R}^d$. This space has natural interpretation as a space of counting measures on \mathbb{R}^d and is endowed with the vague topology induced from the space of Radon measures on \mathbb{R}^d .

2010 *Mathematics Subject Classification.* Primary 82C21; Secondary 34A08.

Key words and phrases. Contact model in the continuum, correlation functions, Caputo-Djrbashian fractional derivative, Mittag-Leffler function.

The financial support of DFG through the SFB 701 (Bielefeld University) is gratefully acknowledged.

Let L be a Markov pre-generator defined on some set of functions $\mathcal{F}(\Gamma)$ given on the configuration space Γ . The backward Kolmogorov equation on observables reads as

$$(1.1) \quad \begin{aligned} \frac{\partial F_t}{\partial t} &= LF_t, \\ F_t|_{t=0} &= F_0 \in \mathcal{F}(\Gamma). \end{aligned}$$

The duality between observables and states on Γ is given by

$$\langle\langle F, \mu \rangle\rangle := \int_{\Gamma} F d\mu.$$

Then the forward Kolmogorov, or Fokker-Planck, equation is the dual one for (1.1):

$$(1.2) \quad \begin{aligned} \frac{\partial \mu_t}{\partial t} &= L^* \mu_t, \\ \mu_t|_{t=0} &= \mu_0. \end{aligned}$$

To be able to construct a Markov process for the generator L we need to define transition probabilities $P_t(\gamma, d\gamma')$ for this process which correspond to the solution to (1.2) for the initial Dirac measures, i.e., $\mu_0(d\gamma') = \delta_{\gamma}(d\gamma')$, $\gamma \in \Gamma$. There appears a principal difference comparing with the classical theory of Markov processes. Namely, we can hope to have the existence of measure evolutions in (1.2) only for special classes of initial states. The latter means that we may have a statistical dynamics for the considered system but a possibility to construct the stochastic evolution remains an open problem.

Moreover, even in such a restricted approach an analysis of Fokker-Planck equations for measures on Γ is a difficult task and at the present time direct methods for the study of such kind evolution equations are absent. There exists a useful technical possibility to transfer evolution problems for measures to their characteristics such as correlation functions.

Let $f^{(n)}(x_1, \dots, x_n)$ be a compactly supported continuous symmetric function on $(\mathbb{R}^d)^n$ and μ be a given probability measure on Γ . The correlation function $k_{\mu}^{(n)}$ of order $n \in \mathbb{N}$ for the measure μ is defined by the following relation:

$$\begin{aligned} \int_{\Gamma} \sum_{\{x_{i_1}, \dots, x_{i_n}\} \subset \gamma} f^{(n)}(x_{i_1}, \dots, x_{i_n}) d\mu(\gamma) \\ = \frac{1}{n!} \int_{(\mathbb{R}^d)^n} f^{(n)}(y_1, \dots, y_n) k_{\mu}^{(n)}(y_1, \dots, y_n) dy_1 \cdots dy_n \end{aligned}$$

and we put $k_{\mu}^{(0)} = 1$. Of course, the existence of correlation functions needs certain a priori properties of the measure μ . On the other hand, correlation functions reconstruct a measure under proper condition of their positive definiteness and a bound on growth, see [9].

The Fokker-Planck equation (1.2) in terms of time dependent correlation functions may be rewritten as an infinite system of evolution equations (hierarchical chain)

$$\frac{\partial k_t^{(n)}}{\partial t} = (L^{\Delta} k_t)^{(n)}, \quad n \geq 0.$$

Here L^{Δ} is the image of L^* in a Fock-type space of vector-functions $k_t = \left(k_t^{(n)}\right)_{n=0}^{\infty}$. In applications to concrete models, the expression for operator L^{Δ} is typically easy to obtain from L using combinatoric calculations, c.f. [5].

The approach to the construction and study of statistical Markov dynamics described above was successfully applied in a number of interacting particle models, see, e.g., [4].

But there exists a possibility to apply similar approach to the case of non-Markov evolutions. Namely, in the Fokker-Planck evolution equation (1.2) we may use the Caputo–Djrbashian (CD) fractional time derivative \mathbb{D}_t^α instead of the usual time derivative. For functions $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ the CD fractional derivative provides a fractional generalization of the first derivative through the following formula in the Laplace transform domain:

$$(\mathcal{L}\mathbb{D}_t^\alpha f)(s) = s^\alpha(\mathcal{L}f)(s) - s^{\alpha-1}f(0), \quad s > 0, \quad \alpha \in (0, 1],$$

where

$$(\mathcal{L}f)(s) = \int_0^\infty e^{-st} f(t) dt.$$

Another representation of the CD derivative is

$$\mathbb{D}_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t \frac{f(\tau) - f(0)}{(t-\tau)^\alpha} d\tau, \quad 0 < \alpha < 1.$$

If f is absolutely continuous, then

$$\mathbb{D}_t^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{f'(\tau)}{(t-\tau)^\alpha} d\tau, \quad 0 < \alpha < 1.$$

In this form the CD fractional derivative is often used in physical literature.

The definition above have natural extensions for vector-valued or measure-valued functions on \mathbb{R}_+ . In the case $0 < \alpha < 1$ we shall expect that the fractional Fokker-Planck dynamics (if exists) will act in the space of states on Γ , i.e., will preserve probability measures on Γ . Of course, this property shall be stated rigorously for each particular model under consideration. From the analytic point of view, the fractional Fokker-Planck equation

$$(1.3) \quad \mathbb{D}_t^\alpha \mu_t = L^* \mu_t,$$

$$\mu_t|_{t=0} = \mu_0.$$

describes a dynamical system with memory in the space of measures on Γ . The corresponding evolution has no longer the semigroup property but we still can try to construct the solution μ_t by means of a related hierarchical chain

$$\mathbb{D}_t^\alpha k_t^{(n)} = (L^\Delta k_t)^{(n)}, \quad n \geq 0.$$

In the present paper we illustrate this approach in the case of so-called contact model in the continuum. This model was introduced in [11] as a Markov birth-and-death process in Γ and studied in detail from the point of view of statistical dynamics in [10]. The Markov generator for this process on a proper set of observables $F : \Gamma \rightarrow \mathbb{R}$ is given by

$$(LF)(\gamma) = \sum_{x \in \gamma} [F(\gamma \setminus \{x\}) - F(\gamma)] + \varkappa \sum_{x \in \gamma} \int_{\mathbb{R}^d} a(x-y) [F(\gamma \cup \{y\}) - F(\gamma)] dy,$$

where $0 \leq a \in L^1(\mathbb{R}^d)$ is even and $\varkappa > 0$. Roughly speaking, the contact process describes branching of points of a configuration in \mathbb{R}^d in the presence of given constant mortality rate for these points which is independent on the existing configuration. We will see that the use of fractional time derivative will not change the equilibrium states of the model but will lead to essentially different asymptotic for the time dependent correlation functions.

2. FRACTIONAL CAUCHY PROBLEM

Let us consider the following Cauchy problem:

$$(2.1) \quad \mathbb{D}^{(\alpha)}u = Au + f, \quad t > 0, \quad u(0) = x,$$

where $\mathbb{D}^{(\alpha)}$ is the CD fractional derivative, $0 < \alpha < 1$, A is a generator of a C_0 -semigroup $T(t)$ on a Banach space X . If $x \in X$, and the vector-function

$$t \ni t \mapsto \int_0^t f(s) ds$$

is exponentially bounded, then a mild solution of (2.1) is given [1], [7] by

$$(2.2) \quad u(t) = S_\alpha(t)x + \int_0^t P_\alpha(t-s)f(s) ds,$$

where

$$(2.3) \quad S_\alpha(t)x = \int_0^\infty \Phi_\alpha(\tau)T(\tau t^\alpha)x d\tau, \quad t \geq 0,$$

$$P_\alpha(t)y = \alpha t^{\alpha-1} \int_0^\infty \tau \Phi_\alpha(\tau)T(\tau t^\alpha)y d\tau, \quad t \geq 0, \quad x, y \in X.$$

Here $\Phi_\alpha(z)$ is the Wright function

$$\Phi_\alpha(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(-\alpha n + 1 - \alpha)}.$$

It is known (see [6]) that

$$\Phi_\alpha(t) \geq 0, \quad t > 0; \quad \int_0^\infty \Phi_\alpha(t) dt = 1, \quad \int_0^\infty t \Phi_\alpha(t) dt = \frac{1}{\alpha \Gamma(\alpha)}.$$

If T is a contraction semigroup, then

$$\|S_\alpha(t)x\| \leq \|x\|, \quad \|P_\alpha(t)y\| \leq \frac{t^{\alpha-1}}{\Gamma(\alpha)} \|y\|.$$

More generally, we have [6]

$$\int_0^\infty t^n \Phi_\alpha(t) dt = \frac{n!}{\Gamma(1 + \alpha n)}, \quad n = 0, 1, 2, \dots$$

If

$$\|T(t)\| \leq e^{\lambda t},$$

then

$$\begin{aligned} \|S_\alpha(t)\| &\leq \int_0^\infty \Phi_\alpha(\tau) e^{\lambda t^\alpha \tau} d\tau = \sum_{n=0}^{\infty} \frac{(\lambda t^\alpha)^n}{\Gamma(1 + \alpha n)} = E_\alpha(\lambda t^\alpha), \\ \|P_\alpha(t)\| &\leq \alpha t^{\alpha-1} \int_0^\infty \tau \Phi_\alpha(\tau) e^{\lambda t^\alpha \tau} d\tau = \alpha t^{\alpha-1} \sum_{n=0}^{\infty} \frac{(\lambda t^\alpha)^n}{n!} \int_0^\infty \tau^{n+1} \Phi_\alpha(\tau) d\tau \\ &= t^{\alpha-1} \sum_{n=0}^{\infty} \frac{(\lambda t^\alpha)^n}{\Gamma(\alpha n + \alpha)} = t^{\alpha-1} E_{\alpha, \alpha}(\lambda t^\alpha). \end{aligned}$$

Here E_α and $E_{\alpha, \alpha}$ are the Mittag-Leffler function and generalized Mittag-Leffler function respectively

$$E_\alpha(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(1 + \alpha n)},$$

$$E_{\alpha,\alpha}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha + \alpha n)}.$$

Here we follow notations from [7], [8].

3. FRACTIONAL CONTACT MODEL

Let us consider the hierarchical chain of evolution equations for correlation functions of the contact model in which we use the CD derivative in time instead of usual time derivative as in [10]. It leads to the following Cauchy problem:

$$(3.1) \quad \mathbb{D}_t^{(\alpha)} k_t^{(n)}(x_1, \dots, x_n) = \widehat{L}_n^* k_t^{(n)}(x_1, \dots, x_n) + f_t^{(n)}(x_1, \dots, x_n), \quad n \geq 1,$$

$$(3.2) \quad \begin{aligned} k_t^{(n)}(x_1, \dots, x_n)|_{t=0} &= k_0^{(n)}(x_1, \dots, x_n), \\ 0 \leq k_0^{(n)}(x_1, \dots, x_n) &\leq C^n n!, \quad C \geq 1. \end{aligned}$$

Here

$$\begin{aligned} \widehat{L}_n^* k_t^{(n)}(x_1, \dots, x_n) &:= -n k_t^{(n)}(x_1, \dots, x_n) \\ &+ \varkappa \sum_{i=1}^n \int_{\mathbb{R}^d} a(x_i - y) k_t^{(n)}(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n) dy, \quad n \geq 1 \end{aligned}$$

and

$$\begin{aligned} f_t^{(n)}(x_1, \dots, x_n) &:= \varkappa \sum_{i=1}^n k_t^{(n-1)}(x_1, \dots, \check{x}_i, \dots, x_n) \sum_{j:j \neq i} a(x_i - x_j), \quad n \geq 2, \\ f_t^{(1)} &\equiv 0. \end{aligned}$$

We assume that $0 \leq a \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ is even and $\varkappa > 0$.

Let $n \in \mathbb{N}$ be arbitrary and fixed. We consider the linear Cauchy problem

$$(3.3) \quad \begin{aligned} \mathbb{D}_t^{(\alpha)} k_t^{(n)}(x_1, \dots, x_n) &= \widehat{L}_n^* k_t^{(n)}(x_1, \dots, x_n) + f_t^{(n)}(x_1, \dots, x_n), \quad t \geq 0, \\ k_t^{(n)}(x_1, \dots, x_n) \Big|_{t=0} &:= k_0^{(n)}(x_1, \dots, x_n) \end{aligned}$$

in the Banach space X_n of bounded measurable functions on $(\mathbb{R}^d)^n$.

The operator \widehat{L}_n^* in X_n can be written also in another way

$$\widehat{L}_n^* k^{(n)}(x_1, \dots, x_n) = n(\varkappa - 1) k^{(n)}(x_1, \dots, x_n) + \sum_{i=1}^n L_a^i k^{(n)}(x_1, \dots, x_n),$$

where for each $1 \leq i \leq n$

$$\begin{aligned} L_a^i k^{(n)}(x_1, \dots, x_n) &= \varkappa \int_{\mathbb{R}^d} a(x_i - y) \left[k^{(n)}(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n) - k^{(n)}(x_1, \dots, x_n) \right] dy \end{aligned}$$

is a generator of a Markov process on $(\mathbb{R}^d)^n$ which describes the jump of the particle placed at the point $(x_1, \dots, x_i, \dots, x_n) \in (\mathbb{R}^d)^n$ into the point $(x_1, \dots, y, \dots, x_n) \in (\mathbb{R}^d)^n$ with intensity equal to $a(x_i - y)$.

Because $a \in L^1(\mathbb{R}^d)$, for any $n \geq 1$ the operator \widehat{L}_n^* is a bounded linear operator on X_n (as well as on $L^1((\mathbb{R}^d)^n)$). Moreover, for each $1 \leq i \leq n$, the operator L_a^i acting in variable x_i is a generator of a contraction semigroup on the space of bounded measurable functions (and on $L^1(\mathbb{R}^d)$); see [10]. Then $\bigotimes_{i=1}^n e^{tL_a^i}$ is a contraction semigroup on X_n (and on $L^1((\mathbb{R}^d)^n)$).

Let

$$T^{(n)}(t) = e^{n(\varkappa-1)t} \bigotimes_{i=1}^n e^{tL_a^i}.$$

We have $\|T^{(n)}(t)\| \leq e^{n(\varkappa-1)t}$. By (2.2), the solution to (3.1), (3.2) is

$$\begin{aligned} k_t^{(n)}(x_1, \dots, x_n) &= (S_\alpha^{(n)}(t)k_0^{(n)})(x_1, \dots, x_n) \\ &+ \int_0^t P_\alpha^{(n)}(t-s) \left[\varkappa \sum_{i=1}^n k_s^{(n-1)}(\dots, \check{x}_i, \dots) \sum_{j \neq i} a(\cdot_i - \cdot_j) \right] ds \end{aligned}$$

(for $n = 1$ the second summand is absent). Here $S_\alpha^{(n)}$ and $P_\alpha^{(n)}$ correspond as above to the semigroup $T^{(n)}$.

4. CORRELATION FUNCTIONS BOUNDS

Our estimates will be based on the identity ([2], page 2)

$$(4.1) \quad \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(z(t-\tau)^\alpha) E_\alpha(\lambda\tau^\alpha) d\tau = \frac{E_\alpha(zt^\alpha) - E_\alpha(\lambda t^\alpha)}{z - \lambda}.$$

Denote $A = \max\{1, \sup_{x \in \mathbb{R}^d} a(x)\}$.

Proposition 4.1. *If $\varkappa > 1$, then*

$$(4.2) \quad \begin{aligned} &k_t^{(n)}(x_1, \dots, x_n) \\ &\leq E_\alpha(n(\varkappa-1)t^\alpha) \left[C^n n! + C^{n-1} n! \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{\varkappa-1} \right)^{j+1} (n-1) \dots (n-j-1) \right]. \end{aligned}$$

The second term is absent for $n = 1$.

Proof. We show (4.2) by induction using the fact that due to (4.1)

$$\begin{aligned} &\int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}((n+1)(\varkappa-1)(t-\tau)^\alpha) E_\alpha(n(\varkappa-1)\tau^\alpha) d\tau \\ &= \frac{E_\alpha((n+1)(\varkappa-1)t^\alpha) - E_\alpha(n(\varkappa-1)t^\alpha)}{\varkappa-1} \leq \frac{1}{\varkappa-1} E_\alpha((n+1)(\varkappa-1)t^\alpha). \end{aligned}$$

Indeed, the inequality (4.2) is obvious for $n = 1$. Suppose it has been proved for some value of n . Then it follows from the estimates for $S_\alpha^{(n)}$ and $P_\alpha^{(n)}$ that

$$\begin{aligned} &k_t^{(n+1)}(x_1, \dots, x_{n+1}) \leq C^{n+1}(n+1)! E_\alpha((n+1)(\varkappa-1)t^\alpha) \\ &+ C^n(n+1)! \varkappa A n \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}((n+1)(\varkappa-1)(t-\tau)^\alpha) E_\alpha(n(\varkappa-1)\tau^\alpha) d\tau \\ &+ C^n(n+1)! \varkappa A n \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{\varkappa-1} \right)^{j+1} (n-1)(n-2) \dots (n-j-1) \\ &\times \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}((n+1)(\varkappa-1)(t-\tau)^\alpha) E_\alpha(n(\varkappa-1)\tau^\alpha) d\tau \\ &\leq E_\alpha((n+1)(\varkappa-1)t^\alpha) \left[C^{n+1}(n+1)! + C^n(n+1)! \frac{\varkappa A n}{\varkappa-1} \right. \\ &\left. + C^n(n+1)! \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{\varkappa-1} \right)^{j+2} n(n-1)(n-2) \dots (n-j-1) \right]. \end{aligned}$$

Making the change $i = j + 1$ of the summation index we see that

$$\begin{aligned} & \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{\varkappa - 1} \right)^{j+2} n(n-1)(n-2) \cdots (n-j-1) \\ &= \sum_{i=1}^{(n+1)-2} \left(\frac{\varkappa A}{\varkappa - 1} \right)^{i+1} [(n+1)-1] \cdots [(n+1)-i-1] \end{aligned}$$

while the missing summand corresponding to $i = 0$ coincides with the expression $\frac{\varkappa An}{\varkappa - 1}$ available in the above estimate. This results in (4.2) with $n + 1$ substituted for n , thus in the estimate (4.2) for any n . \square

It is known that [2], [8]

$$(4.3) \quad E_\alpha(s) \sim \frac{1}{\alpha} e^{s \frac{1}{\alpha}}, \quad s \rightarrow \infty.$$

By (4.2) and (4.3),

$$(4.4) \quad k_t^{(n)}(x_1, \dots, x_n) \leq M C^n n!(n-1)! \frac{q^n}{q-1} e^{[n(\varkappa-1)] \frac{1}{\alpha} t},$$

$$q = \frac{\varkappa A}{\varkappa - 1}, \quad M > 0.$$

Remark 4.2. The case $\varkappa > 1$ corresponds to the super-critical regime in the contact model. As in the case of Markov contact model, we may expect that bound (4.4) is exact, i.e., a similar kind of the lower estimate will be valid; see [3]. For a given $n \in \mathbb{N}$ this bound means an exponential growth of correlation functions for $t \rightarrow \infty$ as in the Markov case. But for a given $t > 0$ we observe much stronger growth of correlations w.r.t. $n \in \mathbb{N}$.

Proposition 4.3. *If $\varkappa < 1$, then*

$$(4.5) \quad \begin{aligned} & k_t^{(n)}(x_1, \dots, x_n) \leq C^n n! E_\alpha(-n(1-\varkappa)t^\alpha) + C^{n-1} n! \\ & \times \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{1-\varkappa} \right)^{j+1} \frac{(n-1)(n-2) \cdots (n-j-1)}{(j+1)!} E_\alpha(-(n-j-1)(1-\varkappa)t^\alpha). \end{aligned}$$

The second term is absent for $n = 1$.

Proof. We proceed by induction, using the fact that by (4.1) for $m < n$,

$$\begin{aligned} & \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(-n(1-\varkappa)(t-\tau)^\alpha) E_\alpha(-m(1-\varkappa)\tau^\alpha) d\tau \\ &= \frac{E_\alpha(-m(1-\varkappa)t^\alpha) - E_\alpha(-n(1-\varkappa)t^\alpha)}{(n-m)(1-\varkappa)} \leq \frac{1}{(n-m)(1-\varkappa)} E_\alpha(-m(1-\varkappa)t^\alpha) \end{aligned}$$

(note that the function $s \mapsto E_\alpha(-s)$ is monotonically decreasing, see [2], page 6).

The inequality (4.5) is obvious for $n = 1$. Suppose it has been proved for some value of n . Then

$$\begin{aligned}
 k_t^{(n+1)}(x_1, \dots, x_{n+1}) &\leq C^{n+1}(n+1)!E_\alpha(-(n+1)(1-\varkappa)t^\alpha) \\
 &+ C^n(n+1)!\varkappa An \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(-(n+1)(1-\varkappa)(t-\tau)^\alpha) E_\alpha(-n(1-\varkappa)\tau^\alpha) d\tau \\
 &+ C^{n-1}(n+1)!\varkappa An \sum_{j=0}^{n-2} \left(\frac{\varkappa A}{1-\varkappa}\right)^{j+1} \frac{(n-1)(n-2)\dots(n-j-1)}{(j+1)!} \\
 &\times \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(-(n+1)(1-\varkappa)(t-\tau)^\alpha) E_\alpha(-(n-j-1)(1-\varkappa)\tau^\alpha) d\tau \\
 &\leq C^{n+1}(n+1)!E_\alpha(-(n+1)(1-\varkappa)t^\alpha) + C^n(n+1)! \frac{\varkappa An}{1-\varkappa} E_\alpha(-n(1-\varkappa)t^\alpha) \\
 &+ C^n(n+1)! \frac{\varkappa An}{1-\varkappa} \sum_{j=0}^{n-2} \left[\left(\frac{\varkappa A}{1-\varkappa}\right)^{j+1} \frac{(n-1)(n-2)\dots(n-j-1)}{(j+2)!} \right. \\
 &\left. \times E_\alpha(-(n-j-1)(1-\varkappa)t^\alpha) \right].
 \end{aligned}$$

As in the proof of Proposition 4.1, we make the change of index $i = j + 1$ and note that the "missing" summand corresponding to $i = 0$ is available in our estimate of $k_t^{(n+1)}$. This proves (4.5) for any n . □

It is known that [2], [8]

$$(4.6) \quad E_\alpha(-s) \sim \frac{1}{\Gamma(1-\alpha)} s^{-1}, \quad s \rightarrow +\infty.$$

By (4.5) and (4.6), for $t \geq 1$,

$$(4.7) \quad k_t^{(n)}(x_1, \dots, x_n) \leq MC^n n!(n-1)! \left(\frac{\varkappa A}{1-\varkappa}\right)^n t^{-\alpha}, \quad M > 0.$$

Remark 4.4. For $\varkappa < 1$ we have the sub-critical regime and in the Markov model correlation functions have exponential decay to zero for $t \rightarrow \infty$ [10]. For $\alpha \in (0, 1)$ and for given $n \in \mathbb{N}$ the bound (4.7) is quite different: we have only $t^{-\alpha}$ decay that demonstrates an essential effect of the memory on the rate of asymptotic degeneration of the system.

Proposition 4.5. *If $\varkappa = 1$, then*

$$(4.8) \quad k_t^{(n)}(x_1, \dots, x_n) \leq C^n n! + \alpha^{-1} C^{n-1} n! \sum_{j=0}^{n-2} \frac{A^{j+1}}{(j+1)\Gamma((j+1)\alpha)} (n-1)(n-2)\dots(n-j-1)t^{(j+1)\alpha}.$$

The second term is absent for $n = 1$.

Proof. As in the previous propositions, we proceed by induction, this time based on the identity

$$\int_0^t (t-\tau)^{\alpha-1} \tau^{\beta-1} d\tau = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} t^{\alpha+\beta-1}.$$

Assuming (4.8) we find that

$$\begin{aligned}
 k_t^{(n+1)}(x_1, \dots, x_{n+1}) &\leq C^{n+1}(n+1)! + C^n(n+1)! \frac{An}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} d\tau \\
 &+ C^{n-1}(n+1)! \frac{An}{\alpha} \sum_{j=0}^{n-2} \frac{A^{j+1}}{(j+1)\Gamma((j+1)\alpha)} (n-1)(n-2)\dots(n-j-1) \\
 &\times \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} \tau^{(j+1)\alpha} d\tau.
 \end{aligned}$$

Computing the integrals, using the identity $\Gamma(1+z) = z\Gamma(z)$ and making a change of the summation index, we obtain the required estimate of $k_t^{(n+1)}$. \square

REFERENCES

1. E. G. Bazhlekova, *Subordination principle for fractional evolution equations*, *Frac. Calc. Appl. Anal.* **3** (2000), no. 3, 213–230.
2. M. M. Djrbashian, *Harmonic Analysis and Boundary Value Problems in Complex Domain*, Birkhäuser, Basel, 1993.
3. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Individual based model with competition in spatial ecology*, *SIAM J. Math. Anal.* **41** (2009), no. 1, 297–317.
4. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Semigroup approach to birth-and-death stochastic dynamics in continuum*, *J. Funct. Anal.* **262** (2012), no. 3, 1274–1308.
5. D. Finkelshtein, Y. Kondratiev, and M. J. Oliveira, *Markov evolutions and hierarchical equations in the continuum. I. One-component systems*, *J. Evol. Equ.* **9** (2009), no. 2, 197–233.
6. R. Gorenflo, Yu. Luchko, and F. Mainardi, *Analytic properties and applications of the Wright function*, *Frac. Calc. Appl. Anal.* **2** (1999), no. 4, 383–414.
7. V. Keyantuo, C. Lizama, and M. Warma, *Spectral criteria for solvability of boundary value problems and positivity of solutions of time-fractional differential equations*, *Abstract and Applied Analysis*, Article ID 614328, 2013.
8. A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
9. Y. Kondratiev and T. Kuna, *Harmonic analysis on configuration space. I. General theory*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **5** (2002), no. 2, 201–233.
10. Y. Kondratiev, O. Kutoviy, and S. Pirogov, *Correlation functions and invariant measures in continuous contact model*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **11** (2008), no. 2, 231–258.
11. Yu. G. Kondratiev and A. Skorokhod, *On contact processes in continuum*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **9** (2006), no. 2, 187–198.

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS’KA, KYIV, 01601, UKRAINE

E-mail address: kochubei@imath.kiev.ua

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, BIELEFELD, 33615, GERMANY

E-mail address: kondrat@math.uni-bielefeld.de

Received 04/09/2014; Revised 30/11/2014