

AROUND OVSYANNIKOV'S METHOD

DMITRI FINKELSHTEIN

Dedicated to Yuriĭ Makarovich Berezanskiĭ on the occasion of his 90th birthday

ABSTRACT. We study existence, uniqueness, and a limiting behavior of solutions to an abstract linear evolution equation in a scale of Banach spaces. The generator of the equation is a perturbation of the operator which satisfies the classical assumptions of Ovsyannikov's method by a generator of a C_0 -semigroup acting in each of the spaces of the scale. The results are (slightly modified) abstract version of those considered in [10] for a particular equation. An application to a birth-and-death stochastic dynamics in the continuum is considered.

1. INTRODUCTION

The study of Markov evolutions for distributions of points in a Euclidean space may be reduced to the study of the evolution equations

$$(1.1) \quad \frac{d}{dt}u(t) = Zu(t), \quad u(0) = u_0,$$

in the Fock-type spaces with weighted L^1 - or L^∞ -norms with an (unbounded) linear operator Z , see, e.g., [23, 18, 13] and the references therein. The well-posedness of the initial value problem (1.1) in a Banach space (existence, uniqueness, and continuous dependence on the initial value) requires the operator Z to be a generator of a strongly continuous (C_0 -) semigroup of linear operators acting in this space. The semigroup approach to the evolution of the so-called birth-and-death dynamics of the distributions of points mentioned above was realized in, e.g., [24, 23, 11, 13]. In particular, the technical restrictions on the birth and death rates were introduced. However, for some important models these restrictions either were not satisfied, cf. [17], or required more strong assumptions on the birth and death rates (that is less interesting for applications), compare [24] and [9] or [11] and [10]. Moreover, the dynamics with jumps were not covered by the semigroup approach at all, see, e.g., [4, 5, 15].

To overcome these restrictions, the evolution (1.1) was allowed to be considered in a scale of Banach spaces. In the latter approach the dynamics was constructed on a finite time interval $[0, T)$ only, and, for any $t \in (0, T)$, the solution $u(t)$ to (1.1) belonged to a proper space of the scale. This was realized in [9, 4, 19, 20, 16] using the so-called Ovsyannikov's method: it requires Z to be considered as a bounded operator between each two spaces $B_{\alpha'} \subset B_{\alpha''}$, of a scale $\{B_\alpha\}_{\alpha \in I}$, $I \subset \mathbb{R}$, of Banach spaces with the operator norm proportional to $(\alpha'' - \alpha')^{-1}$. Originally this method was found by G. E. Shilov and A. G. Kostyuchenko, and it was firstly published in 1958 in the book [21] by I. M. Gelfand and G. E. Shilov. In particular, it was applied to the so-called Kovalevskya's system of first-order PDE. In 1960, T. Yamanaka generalized this approach for the case of the time dependent operator $Z(t)$. In 1965, the method was rediscovered

2000 *Mathematics Subject Classification.* Primary 35K90, 47D06; Secondary 82C22, 60J80.

Key words and phrases. Ovsyannikov's method, scale of spaces, evolution equations, birth-and-death dynamics, Vlasov scaling, kinetic equation.

in [29] by L. V. Ovsjannikov and was named in the book [32] written by F. Trèves, who has realized a detailed analysis of the problem

$$(1.2) \quad \frac{d}{dt}u(t) = Z(t)u(t) + f(t), \quad u(s) = u_s, \quad s \geq 0,$$

with numerous applications. After that, the initial value problems (1.1), (1.2) with the operator norm estimate as above were named the ‘abstract Kovalevskya’s systems’ in the literature. (Note that the method allowed the immediate generalization for the complex values of t .) The non-linear generalization was introduced by F. Trèves [33] under assumptions which were essentially simplified by L. Nirenberg [27] and T. Nishida [28]. K. Deimling generalized the linear case to the equation

$$(1.3) \quad \frac{d}{dt}u(t) = Z(t)u(t) + f(t, u(t)), \quad u(s) = u_s, \quad s \geq 0,$$

for a some class of functions f . Being quite general, the conditions on f guaranteed the existence of a solution to (1.2) only, cf., e.g., the survey [25]. For further generalizations see, e.g., [2, 35, 6, 31, 36, 1].

On the other hand, in [10], there were considered the birth-and-death dynamics of complex (point) systems in the continuum whose generator of the evolution of correlation functions (see Section 3 below for the definitions) did not allow the singularity $(\alpha'' - \alpha')^{-1}$ for the norm of Z in (1.1). Namely, the singularity was of the order $(\alpha'' - \alpha')^{-2}$ which can not be realized in the scheme of Ovsyannikov’s method. Fortunately, the corresponding evolution equation could be rewritten in the form

$$(1.4) \quad \frac{d}{dt}u(t) = Zu(t) + Au(t), \quad u(0) = u_0,$$

where Z was suitable for Ovsyannikov’s method and A was a generator of a contraction semigroup acting in *each* of the spaces of the scale. The important point is that the evolution of correlation functions was considered in the scale of L^∞ -type spaces (see, e.g., [13, 14] for the explanation of the reasons). By [26], there is not a C_0 -semigroup in a space isomorphic to L^∞ with an unbounded generator (see also the proof of Proposition 3.1 below). Therefore, the semigroups generated by A were considered in suitable subspaces of the spaces of the scale, using the technique of the so-called sun-dual semigroups, see e.g. [7, 34], which goes back to R. S. Phillips [30]. Another problem considered in [10] was the convergence of the solutions to the equation

$$(1.5) \quad \frac{d}{dt}u_\varepsilon(t) = Zu_\varepsilon(t) + Au_\varepsilon(t), \quad u(0) = u_0,$$

to the solution to the limiting equation, in the course of the so-called Vlasov-type scaling, see, e.g., [12, 13] for details. This allowed to derive the so-called kinetic equation which approximately describes the behavior of the density of the birth-and-death dynamics. All constructions in [10] were done for the particular model, however, in a form useful for further generalization.

The aims of the present paper are the following. First, we consider the abstract equation (1.4) in an (increasing) scale of Banach spaces. The assumptions on A and Z will be abstract and slightly generalized versions of those in [10]. For this equation we prove the existence result (Theorem 2.1) and show the uniqueness of the ‘integral curves’ of the differential equation (Theorem 2.4). As was mentioned above, for any $t \in (0, T)$, the solution to (1.4) belongs to a space of the scale, more precisely, $u(t) \in B_\alpha$, for all $\alpha > \alpha_t$, for some ‘minimal’ α_t . In general, one can not check whether $u(t) \in B_{\alpha_t}$. Therefore, it is quite natural to consider (1.4) in the scale of projective limits $\{\bigcap_{\beta > \alpha} B_\beta\}_{\alpha \in I}$, $I \subset \mathbb{R}$ (Proposition 2.6). This allows to prove rigorously that the flow $t \mapsto u(t)$ is indeed unique and satisfies the semigroup property; surely, on a finite time interval only (Proposition 2.21). The problem here was that, in contrast to the classical existence

and uniqueness Picard–Lindelöf theorem for ordinary differential equations, where the finite time interval was defined by an estimate on the right hand side of the equation, in Ovsyannikov’s methods we should *choose* the initial and the terminal spaces, and they generate the time interval. As a result, we need to be sure that by choosing more wider terminal space we will have the same solution as in the smaller terminal space (with the same initial space), see also Remark 2.5 below. Also one should verify that, by choosing a properly small intermediate moment of time, we could start a new initial value problem at that time and get the same solution thereafter as if we would consider the first problem on a bigger time interval.

The second aim is to consider the convergence of the solutions to the abstract equations (1.5). For the case $A_\varepsilon = 0$, $\varepsilon > 0$, the corresponding abstract statement was proved in [19]. In Theorem 2.10, we generalize this scheme by a simple modification of the statements presented in [10], for the particular model discussed there.

Finally, in Section 3, we consider birth-and-death dynamics using the technique above. The dynamics are a ‘combination’ of those studied in [8] and [17]; it describes the evolution in the course of which the elements disappear (die) more intensively in that regions of the space, where the amounts of their ‘neighbors’ are too big or too small. The birth rate of the dynamics is assumed to be constant in the space: the elements appear from the outside ‘reservoir’ of the system. We prove that the corresponding dynamics exist on a finite time interval and realize the Vlasov-type scaling. This yields a nonlocal nonlinear kinetic equation of a new type. In particular, this equation may have one or three positive stationary points depending on the values of parameters of the system, see Remark 3.3 below. The detailed analysis of this equation will be done in a subsequent paper.

The author is grateful to Prof. Dr. Yuri Kondratiev, Prof. Dr. Yuri Kozitsky, and Dr. Oleksandr Kutoviy for fruitful discussions.

2. MAIN RESULTS

Let us fix several arrangements. We will use notations like $(B, \|\cdot\|)$ to say that B is a Banach space with a norm $\|\cdot\|$. Let A be an (unbounded) linear operator on B with a domain $D \subset B$, we will denote this by (A, D) . Note that here and below any inclusion allows equality. A strongly continuous semigroup of linear bounded operators $S(t)$, $t \geq 0$, on B will be called a C_0 -semigroup. By e.g. [7, Proposition I.5.5], there exist $\omega \in \mathbb{R}$ and $\nu \geq 1$ such that $\|S(t)\| \leq \nu e^{\omega t}$, $t \geq 0$, i.e. $S(t)$ is exponentially bounded. If C is a linear subset of B , we will denote the closure of C with respect to the norm of B by \overline{C} . Note that if C is closed, i.e. $\overline{C} = C$, then $(C, \|\cdot\|_B)$ is a Banach space as well.

Our first assumption is about a scale of Banach spaces where the dynamics will exist.

Assumption 1. Let $\underline{\alpha} > 0$, $\overline{\alpha} \in (0, \infty]$ be fixed; set $I = (\underline{\alpha}, \overline{\alpha})$. Let $\mathbf{B} := (B_\alpha)_{\alpha \in I}$ be a family of Banach spaces $(B_\alpha, \|\cdot\|_\alpha)$ which is supposed to be increasing, i.e.

$$(2.1) \quad B_{\alpha'} \subset B_{\alpha''}, \quad \|\cdot\|_{\alpha'} \geq \|\cdot\|_{\alpha''}, \quad \alpha' \leq \alpha'', \quad \alpha', \alpha'' \in I.$$

Suppose, additionally, that, $u \in B_{\alpha'} \subset B_{\alpha''}$, $u = 0$ in $B_{\alpha''}$ yields $u = 0$ in $B_{\alpha'}$, for any α', α'' as above.

Next, we consider a linear mapping A on $B_I := \bigcup_{\alpha \in I} B_\alpha$ which acts in each of spaces of the scale \mathbf{B} and satisfies the following assumption.

Assumption 2. Let, for any $\alpha \in I$, in the space B_α , there exist a closed linear subset C_α and its dense linear subset D_α , i.e. $D_\alpha \subset C_\alpha \subset B_\alpha$ and $\overline{D_\alpha} = C_\alpha = \overline{C_\alpha}$. Let $A : B_I \rightarrow B_I$ be linear operators and such that, for any $\alpha \in I$, the operator (A, D_α) is a generator of a C_0 -semigroup $S_\alpha(t)$ on the Banach space $(C_\alpha, \|\cdot\|_\alpha)$. Assume that, for any $\alpha' \in I$ with $\alpha' < \alpha$, $B_{\alpha'} \subset D_\alpha$; $B_{\alpha'}$ is $S_\alpha(t)$ -invariant, and $S_\alpha(t) \upharpoonright_{B_{\alpha'}} = S_{\alpha'}(t)$. Suppose

also that the constants $\nu \geq 1$, $\omega \in \mathbb{R}$, in the definition of quasi-boundedness for $S_\alpha(t)$ are independent in α , i.e.

$$(2.2) \quad \|S_\alpha(t)\| \leq \nu e^{\omega t}, \quad t \geq 0, \quad \alpha \in I.$$

Finally, we will deal with a linear mapping Z on B_I which may be considered as a bounded operator between each two spaces of the scale \mathbf{B} .

Assumption 3. Let $M, N : I \rightarrow (0, \infty)$ be increasing continuous functions. Let, for any $\alpha^* \in I$ and for any $\alpha', \alpha'' \in (\underline{\alpha}, \alpha^*]$ with $\alpha' < \alpha''$, Z be a bounded linear operator from $B_{\alpha'}$ to $B_{\alpha''}$, such that the following estimate holds:

$$(2.3) \quad \|Zu\|_{\alpha''} \leq \left(\frac{M(\alpha^*)}{\alpha'' - \alpha'} + N(\alpha^*) \right) \|u\|_{\alpha'}, \quad u \in B_{\alpha'}.$$

(Note that M, N may depend on $\underline{\alpha}$.)

Under Assumptions above, consider the following function

$$(2.4) \quad T(\alpha, \beta) := \frac{\beta - \alpha}{e\nu M(\beta)}, \quad \beta \geq \alpha > \underline{\alpha}.$$

Theorem 2.1. *Let Assumptions 1–3 hold. Let $\alpha^* \in I$ and $s \geq 0$. Take an arbitrary $\alpha_s \in (\underline{\alpha}, \alpha^*)$ and set $T := T(\alpha_s, \alpha^*)$. Then, for any $u_s \in B_{\alpha_s}$, there exists a function $u : [s, s+T) \rightarrow B_{\alpha^*}$ such that*

- (1) u is continuous on $[s, s+T)$ and continuously differentiable on $(s, s+T)$;
- (2) for any $t \in (s, s+T)$, $Au(t) \in B_{\alpha^*}$ and $Zu(t) \in B_{\alpha^*}$;
- (3) u solves the following differential equation:

$$(2.5) \quad \frac{d}{dt}u(t) = Au(t) + Zu(t), \quad t \in (s, s+T);$$

$$(4) \quad u(s) = u_s.$$

Proof. We will follow the scheme from [10]. Take arbitrary $\Upsilon \in (0, T)$. By the continuity of M , there exists $\alpha \in (\alpha_s, \alpha^*)$ such that $\Upsilon < T(\alpha_s, \alpha) =: T'$. Let $q = q(\Upsilon, T, T') > 1$ be such that $q\Upsilon < \min\{T, T'\}$. For any $n \in \mathbb{N}$, consider the following partition of the interval $[\alpha_s, \alpha]$ on $2n+2$ parts:

$$(2.6) \quad \alpha^{(2j,n)} = \alpha_s + j(\delta_1 + \delta_2), \quad \alpha^{(2j+1,n)} = \alpha^{(2j,n)} + \delta_1,$$

where $j = 0, \dots, n$ and

$$(2.7) \quad \delta_1 = \frac{(q-1)(\alpha - \alpha_s)}{q(n+1)}, \quad \delta_2 = \frac{\alpha - \alpha_s}{qn}.$$

In particular, $\alpha^{(0,n)} = \alpha_s$, $\alpha^{(2n+1,n)} = \alpha$, and

$$(2.8) \quad \alpha^{(2j+2,n)} - \alpha^{(2j+1,n)} = \delta_2, \quad j = 0, \dots, n-1.$$

For an $n \in \mathbb{N}$, consider a mapping on B_I

$$(2.9) \quad U_\alpha^{(n)}(t, t_1, \dots, t_n) = S_\alpha(t - t_1)ZS_\alpha(t_1 - t_2)Z \dots S_\alpha(t_{n-1} - t_n)ZS_\alpha(t_n),$$

where we set $t_0 := t$. Then, for any $\tau > 0$ and $u \in B_{\alpha^{(2j,n)}} \subset D_{\alpha^{(2j+1,n)}} \subset C_{\alpha^{(2j+1,n)}}$, one has $S_\alpha(\tau)u = S_{\alpha^{(2j,n)}}(\tau)u \in B_{\alpha^{(2j+1,n)}}$ and hence $ZS_\alpha(\tau)u \in B_{\alpha^{(2j+2,n)}}$, $j = 0, \dots, n-1$, with

$$\|ZS_\alpha(\tau)u\|_{\alpha^{(2j+2,n)}} \leq \nu e^{\omega\tau} \left(\frac{M(\alpha)}{\delta_2} + N(\alpha) \right) \|u\|_{\alpha^{(2j,n)}}.$$

As a result, $U_\alpha^{(n)}(t, t_1, \dots, t_n)u_s \in B_\alpha$ with

$$(2.10) \quad \begin{aligned} \|U_\alpha^{(n)}(t, t_1, \dots, t_n)u_s\|_\alpha &\leq \nu^{n+1} e^{\omega t} \left(\frac{M(\alpha)}{\delta_2} + N(\alpha) \right)^n \|u\|_{\alpha_s} \\ &= \nu e^{\omega t} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^n \|u\|_{\alpha_s}. \end{aligned}$$

Set now $V_\alpha^{(1)}(s, t)u_s := \int_s^t U^{(1)}(t, t_1)u_s dt_1$ and

$$V_\alpha^{(n)}(s, t)u_s := \int_s^t \int_s^{t_1} \dots \int_s^{t_{n-1}} U_\alpha^{(n)}(t, t_1, \dots, t_n)u_s dt_n \dots dt_1.$$

Therefore, the series

$$(2.11) \quad S_\alpha(t-s)u_s + \sum_{n=1}^{\infty} V_\alpha^{(n)}(s, t)u_s$$

is majorized in B_α by the series

$$(2.12) \quad \nu e^{\omega t} \|u_s\|_{\alpha_s} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^n (t-s)^n,$$

which converges uniformly on $t \in [s, s + \Upsilon]$, as

$$(2.13) \quad \left(\frac{1}{n!} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^n (t-s)^n \right)^{\frac{1}{n}} \\ \sim \frac{e}{n} \left(\frac{qn}{eT'} + \nu N(\alpha) \right) (t-s) \sim \frac{q(t-s)}{T'} \leq \frac{q\Upsilon}{T'} < 1.$$

Therefore, the series (2.11) converges uniformly on $t \in [s, s + \Upsilon]$ in B_α to a function $u(t) \in B_\alpha$. Evidently, each term of (2.11) is continuous as a mapping $[s, s + \Upsilon] \rightarrow B_\alpha$ thus $u(t)$ is continuous as well. Since the norm in B_α is stronger than in B_{α^*} one has that $[s, s + \Upsilon] \ni t \mapsto u(t) \rightarrow B_{\alpha^*}$ is also continuous.

Consider now the series of derivatives of the terms from (2.11). Each of them belongs to $B_\alpha \subset D_{\alpha^*}$, thus

$$(2.14) \quad \frac{d}{dt} V_\alpha^{(n)}(s, t) = \int_s^t \int_s^{t_2} \dots \int_s^{t_{n-1}} U^{(n)}(t, t, t_2, \dots, t_n)u_s dt_n \dots dt_1 \\ + \int_s^t \int_s^{t_1} \dots \int_s^{t_{n-1}} AU_\alpha^{(n)}(t, t_1, \dots, t_n)u_s dt_n \dots dt_1 \\ = ZV_\alpha^{(n-1)}(s, t) + AV_\alpha^{(n)}(s, t)$$

is well-defined and belongs to B_{α^*} . By the same arguments as above the series of derivatives converges uniformly on $t \in [s, s + \Upsilon]$ in B_{α^*} and hence its sum is equal to $\frac{d}{dt}u(t)$. Note also that $u(t) \in B_\alpha \subset D_{\alpha^*}$ and hence $Au, Zu \in B_{\alpha^*}$. Thus, by (2.14), $u(t)$ satisfies (2.5). Since $\Upsilon \in (0, T)$ was arbitrary, the statement is proved. \square

Remark 2.2. It is easy to see that the summand $N(\alpha^*)$ in (2.3) might be changed on $\frac{N(\alpha^*)}{(\alpha'' - \alpha')^\delta}$, for an arbitrary $\delta \in (0, 1)$, without any changes in (2.4).

Corollary 2.3. *Let conditions and notations of Theorem 2.1 hold. Set*

$$N_* := \sup_{\alpha \in [\alpha_s, \alpha^*]} N(\alpha) < \infty, \quad T_* := \sup_{\alpha \in [\alpha_s, \alpha^*]} T(\alpha_s, \alpha) < \infty.$$

Then, for any $\Upsilon \in (0, T)$ and $\alpha \in (\alpha_s, \alpha^)$ such that $\Upsilon < T(\alpha_s, \alpha) =: T'$, and for any $q \in (1, \frac{T'}{\Upsilon})$,*

$$(2.15) \quad \|u(t)\|_\alpha \leq \frac{C}{T' - q\Upsilon} e^{\omega t} \|u_s\|_{\alpha_s}, \quad t \in [s, s + \Upsilon],$$

where $C = C(\nu, T_*, N_*) > 0$.

Proof. By (2.12), we have, for $t \in [s, s + \Upsilon]$,

$$\begin{aligned} \|u(t)\|_\alpha &\leq \nu e^{\omega t} \|u_s\|_{\alpha_s} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{n}{e}\right)^n \left(\frac{q\Upsilon}{T'}\right)^n \left(1 + \frac{e\nu T' N(\alpha)}{qn}\right)^n, \\ &\leq \nu e^{\omega t} \|u_s\|_{\alpha_s} \sum_{n=0}^{\infty} \frac{1}{e} \left(\frac{q\Upsilon}{T'}\right)^n \left(1 + \frac{e\nu T_* N_*}{n}\right)^n \\ &\leq \nu e^{\omega t + e\nu T_* N_* - 1} \|u_s\|_{\alpha_s} \frac{T_*}{T' - q\Upsilon}, \end{aligned}$$

where we have used the following elementary inequalities

$$n! \geq e\left(\frac{n}{e}\right)^n, \quad \left(1 + \frac{x}{n}\right)^n \leq e^x, \quad n \in \mathbb{N}, \quad x > 0. \quad \square$$

Now we are ready to formulate a uniqueness result.

Theorem 2.4. *Let Assumptions 1–3 hold. Let $\alpha^* \in I$ and $T > 0$. Let, for some $s \geq 0$ and $\tau \in [s, s + T)$ continuous functions $[\tau, s + T) \rightarrow u_i(t) \in B_{\alpha^*}$, $i = 1, 2$ satisfy to the differential equation (2.5) on $(\tau, s + T)$ in B_{α^*} . Suppose that there exists $\alpha_\tau \in (\underline{\alpha}, \alpha^*)$ such that $\tau + T(\alpha_\tau, \alpha^*) \geq s + T$ and $u_1(\tau) = u_2(\tau) =: u_\tau \in B_{\alpha_\tau}$. Then $u_1(t) = u_2(t)$ in B_{α^*} , for any $t \in (\tau, s + T)$.*

In particular, the function u in Theorem 2.1 is unique.

Proof. Take an arbitrary $\Upsilon \in (\tau - s, T)$ thus $\tau < s + \Upsilon < s + T < \tau + T(\alpha_\tau, \alpha^*)$. Since $\alpha^* \in I$, I is an open interval and M is continuous on I , there exists $\alpha^\circ \in I$ such that $\alpha^\circ > \alpha^*$ and $s + \Upsilon < \tau + T(\alpha_\tau, \alpha^\circ)$. Let $u(t) := u_1(t) - u_2(t)$, $t \in [\tau, s + T)$. Then u solves (2.5) on $(\tau, s + T)$ with $u(\tau) = 0 \in B_{\alpha_\tau}$. It is enough to prove that $u(t) = 0 \in B_{\alpha^\circ}$ (and thus $u(t) = 0 \in B_{\alpha^*}$). Since the norm in B_{α^*} is stronger than the norm in B_{α° , $u(t)$ solves (2.5) in B_{α° as well. Then, one has the following equality in B_{α°

$$(2.16) \quad u(t) = \int_\tau^t S_{\alpha^\circ}(t - t') Zu(t') dt', \quad t \in [\tau, s + \Upsilon].$$

However, $u(t) \in B_{\alpha^*}$ hence, for any $\alpha' \in (\alpha^*, \alpha^\circ)$, one can take any $\alpha'' \in (\alpha^*, \alpha')$ and consider the right hand side of (2.16) as follows: $u(t') \in B_{\alpha^*}$, $Zu(t') \in B_{\alpha''} \subset D_{\alpha'}$, $S_{\alpha^\circ}(t - t') Zu(\tau) = S_{\alpha'}(t - t') Zu(\tau) \in B_{\alpha'} \subset B_{\alpha^\circ}$, and all the mappings are continuous. Therefore, one can iterate (2.16) n times and consider partition (2.6)–(2.7) with α°, α^* in place of α^*, α_s respectively. As a result, one gets, cf. (2.10),

$$(2.17) \quad \|u(t)\|_{\alpha^\circ} \leq e^{\omega t} \left(\frac{q'n}{eT(\alpha^*, \alpha^\circ)} + \nu N(\alpha^\circ) \right)^n \frac{(t - \tau)^n}{n!} \|u(t)\|_{\alpha^*}, \quad t \in [\tau, s + \Upsilon],$$

with a properly chosen $q' > 1$. Let now $N \in \mathbb{N}$ be big enough to guarantee that (2.17) implies $\|u(t)\|_{\alpha^\circ} = 0$, for $t \in [\tau, \tau + \sigma]$, $\sigma := \frac{s + \Upsilon - \tau}{N}$, i.e. $N > \frac{s + \Upsilon - \tau}{T(\alpha^*, \alpha^\circ)} > 0$. Thus $u(t) = 0$ in B_{α° and hence in B_{α^*} , for $t \in [\tau, \tau + \sigma]$. Repeat now the same arguments with initial zero value at $t = \tau + \sigma$, it will lead to the zero solution in B_{α^*} on $[\tau + \sigma, \tau + 2\sigma]$ and so on. As a result, we will get that $u(t) = 0$ in B_{α^*} on the whole $[\tau, s + \Upsilon]$, and since Υ was arbitrary we will have the uniqueness on the $[\tau, s + T)$. \square

Remark 2.5. In applications, we often have an estimate like (2.3) with $\tilde{M}(\alpha'', \alpha')$ in place of $\mathcal{M}(\alpha^*)$, with a function \tilde{M} which is increasing in the first variable and decreasing in the second one thus $\tilde{M}(\alpha'', \alpha') \leq \tilde{M}(\alpha^*, \underline{\alpha}) =: M(\alpha^*)$, and M is an increasing function, cf. [9, 4, 16, 17]. Note also, that the function $T(\alpha, \beta)$ is not typically increasing in β , see (2.4) and the references above; therefore, the bigger terminal space B_{α^*} does not necessarily lead to a wider time interval. Note that, in the cited references, the function $T(\alpha, \cdot)$ had a unique maximum point.

For any $\alpha \in I$, consider the set

$$(2.18) \quad B_{\alpha+} := \bigcap_{\beta > \alpha} B_{\beta},$$

which may be endowed by the sequential topology of a projective space, see e.g. [3], i.e. $u_n \rightarrow u$ in $B_{\alpha+}$ if and only if $u_n \rightarrow u$ in all B_{β} , $\beta > \alpha$ (of course, by (2.1), it is sufficient to take $\beta \in (\alpha, \alpha + \delta)$ only, for some $\delta > 0$). Stress that, under Assumption 3, Z is continuous on $B_{\alpha+}$.

Proposition 2.6. *In conditions and notations of Theorem 2.1,*

- (1) *the mapping $B_{\alpha_s} \ni u_s \mapsto u(t) \in B_{\alpha^*}$ is continuous, uniformly in $t \in [s, s + \Upsilon] \subset [s, s + T]$;*
- (2) *for any $t \in (s, s + T)$, there exist*

$$(2.19) \quad \alpha(t, s, \alpha_s) := \inf \{ \alpha \in [\alpha_s, \alpha^*) \mid u(t) \in B_{\alpha} \} < \alpha^*,$$

such that $u(t) \in B_{\alpha(t, s, \alpha_s)+}$ and the mapping $B_{\alpha_s} \ni u_s \mapsto u(t) \in B_{\alpha(t, s, \alpha_s)+}$ is also continuous, uniformly in $t \in [s, s + \Upsilon] \subset [s, s + T]$;

- (3) *one can take $u_s \in B_{\alpha_s+}$; then all previous statements remain true with B_{α_s+} in place of B_{α_s} only.*

Proof. We will use details of the proof of Theorem 2.1.

(1) Let $v(t)$ solve (2.5) on $[s, s + T)$ with $v(0) = v_s \in B_{\alpha_s}$. Then we will have that, for any $t \in [s, s + \Upsilon] \subset [s, s + T)$, and for the same $\alpha \in (\alpha_s, \alpha^*)$, $q > 1$,

$$(2.20) \quad \begin{aligned} \|u(t) - v(t)\|_{\alpha^*} &\leq \|u(t) - v(t)\|_{\alpha} \\ &\leq \max\{e^{\omega(s+\Upsilon)}, 1\} \|u_s - v_s\|_{\alpha_s} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{qn}{\epsilon T'} + \nu N(\alpha) \right)^n \Upsilon^n, \end{aligned}$$

that implies the needed continuity, as, recall, α depends on Υ , $q = q(T, T') = q(\alpha, \alpha_s)$ and thus the estimate is uniform in $t \in [s, s + \Upsilon]$.

(2) Recall that the solution $u(t)$ in B_{α^*} to (2.5) is given on $[s, s + T)$ by (2.11) and, for a chosen $t \in [s, s + T)$, the value $u(t)$, as a matter of fact, belongs to B_{α} , for any $\alpha \in [\alpha_s, \alpha^*)$ such that $s < t < s + T(\alpha_s, \alpha)$. Since $T(\alpha_s, \alpha_s) = 0$ we have by the continuity arguments that there exists $\alpha^\circ = \alpha(t, s, \alpha_s)$ such that $T(\alpha_s, \alpha^\circ) = t - s$ and there exists an open subinterval of (α_s, α^*) where $T(\alpha_s, \alpha) > t - s$ (or a union of such subintervals). Thus, the set in (2.19) is non-empty, the infimum does exist, that yields the first statement. Next, by (2.20), the mapping $B_{\alpha_s} \ni u_s \mapsto u(t) \in B_{\alpha}$ will be continuous, for any $\alpha > \alpha(t, s, \alpha_s)$, uniformly in $t \in [s, s + \Upsilon]$. This fulfilled the statement.

(3) Let $u_s \in B_{\alpha_s+}$. For any $\Upsilon \in (0, T)$, one can choose $\alpha'_s \in (\alpha_s, \alpha^*)$ with $\alpha'_s - \alpha_s$ small enough to guarantee that $\Upsilon < T(\alpha'_s, \alpha^*) < T$. Then one can repeat all arguments above and get that there exists solution to (2.5) on $[s, s + \Upsilon]$ in B_{α^*} with $u(0) = u_s \in B_{\alpha'_s}$ such that $B_{\alpha'_s} \ni u_s \mapsto u(t) \in B_{\alpha^*}$ (and B_{α^*} may be replaced on $B_{\alpha(t, s, \alpha'_s)+}$). If we take now $\alpha''_s \in (\alpha_s, \alpha^*)$, $\alpha''_s \neq \alpha'_s$, then the solution will be given by *the same* series (2.11) which converges *in the same* space B_{α^*} (or even in a smaller space). The difference will be in the denominator of (2.15) only: namely, $\|u(t)\|_{\beta} \leq \bar{C}(\beta) \|u_s\|_{\gamma}$, $\gamma > \alpha_s$, $\beta \in (\alpha(t, s, \alpha'_s), \alpha^*)$, $t \in [s, s + \Upsilon]$. This, naturally, implies the continuity of the mapping $B_{\alpha_s+} \ni u_s \mapsto u(t) \in B_{\alpha(t, s, \alpha_s)+}$. \square

Remark 2.7. Note that, by the comparison series criterion, we have from (2.13) that the majorized series for (2.11) diverges for $\alpha = \alpha(t, s, \alpha_s)$. However, one can not state that $u(t) \notin B_{\alpha(t, s, \alpha_s)}$.

According to Theorem 2.6, for any $s \geq 0$, $\alpha^* \in I$, $\alpha_s \in (\underline{\alpha}, \alpha^*)$, $t \in [s, s + T(\alpha_s, \alpha^*)]$ one can define the mapping $U(s, t) : B_{\alpha_s} \rightarrow B_{\alpha(t, s, \alpha_s)}$ given by

$$(2.21) \quad U(s, t)u_s = u(t).$$

Proposition 2.8. *Let $s \geq 0$, $\alpha^* \in I$, $\alpha_s \in B_{\alpha_s}$ be arbitrary. Let $t > \tau > s$ be such that $\tau < s + T(\alpha_s, \alpha^*)$, $t < \min\{\tau + T(\alpha(\tau, s, \alpha_s), \alpha^*), s + T(\alpha_s, \alpha^*)\}$. Then*

$$(2.22) \quad U(s, t)u_s = U(\tau, t)U(s, \tau)u_s.$$

Proof. The statements follows from the uniqueness Theorem 2.4. Indeed, $\sigma := \min\{\tau + T(\alpha(\tau, s, \alpha_s), \alpha^*), s + T(\alpha_s, \alpha^*)\} > s$, therefore, $T := \sigma - s > 0$. Then, by the construction of the mapping (2.21), both functions $U(s, t)u_s$ and $U(\tau, t)U(s, \tau)u_s$ solves (2.5) on $(\tau, s + T)$ and they are both equal to $U(s, \tau)u_s$ at $t = \tau$. Hence, by Theorem 2.4 they coincide on $(\tau, \tau + T)$ as well. \square

Remark 2.9. In the same manner as before, one prove the following statement. In conditions and notations of Theorem 2.1, suppose, additionally that there exists $\alpha^{**} \in I$ such that $\alpha^* < \alpha^{**}$ and Assumption 3 holds for α^{**} in place of α^* . Set $\tilde{T} := T(\alpha_s, \alpha^{**})$ and $T_0 := \min\{T, \tilde{T}\}$. Let $\tilde{u} : [s, s + \tilde{T}] \rightarrow B_{\alpha^{**}}$ be the solution to (2.5) according to Theorem 2.1. Then, for any $t \in [s, s + T_0]$, $\tilde{u}(t) = u(t) \in B_{\alpha^*}$.

Assumption 4. Let $\{D_\alpha, C_\alpha\}_{\alpha \in I}$ be such as in Assumption 2. Let $A_\varepsilon : B_I \rightarrow B_I$, $\varepsilon \geq 0$ be linear operators, such that, for any $\varepsilon \geq 0$ and for any $\alpha \in I$, the operator $(A_\varepsilon, D_\alpha)$ is a generator of a C_0 -semigroup $S_{\alpha, \varepsilon}(t)$ on the Banach space $(C_\alpha, \|\cdot\|_\alpha)$. Assume that, for any $\alpha' \in I$ with $\alpha' < \alpha$, $B_{\alpha'} \subset D_\alpha$; $B_{\alpha'}$ is $S_{\alpha, \varepsilon}(t)$ -invariant, and $S_{\alpha, \varepsilon}(t) \upharpoonright_{B_{\alpha'}} = S_{\alpha', \varepsilon}(t)$. Suppose also that the constants $\nu \geq 1$, $\omega \in \mathbb{R}$ are such that

$$(2.23) \quad \|S_{\alpha, \varepsilon}(t)\| \leq \nu e^{\omega t}, \quad t \geq 0, \quad \alpha \in I, \quad \varepsilon \geq 0.$$

Assumption 5. Let $M, N : I \rightarrow (0, \infty)$ be increasing continuous functions. Let, for any $\alpha^* \in I$ and for any $\alpha', \alpha'' \in (\underline{\alpha}, \alpha^*]$ with $\alpha' < \alpha''$, Z_ε , $\varepsilon \geq 0$, be bounded linear operators from $B_{\alpha'}$ to $B_{\alpha''}$, such that the following estimate holds:

$$(2.24) \quad \|Z_\varepsilon u\|_{\alpha''} \leq \left(\frac{M(\alpha^*)}{\alpha'' - \alpha'} + N(\alpha^*) \right) \|u\|_{\alpha'}, \quad u \in B_{\alpha'}.$$

Theorem 2.10. *Let Assumption 1, 4, 5 hold. Let $P_\varepsilon, p_\varepsilon : I \rightarrow (0, \infty)$, $\varepsilon > 0$ be increasing continuous functions, such that*

$$(2.25) \quad \lim_{\varepsilon \rightarrow 0} p_\varepsilon(\alpha) = \lim_{\varepsilon \rightarrow 0} P_\varepsilon(\alpha) = 0, \quad \alpha \in I,$$

and let $r \in \mathbb{N}$. Let $\alpha^* \in I$ and $\alpha', \alpha'' \in (\underline{\alpha}, \alpha^*)$, $\alpha' < \alpha''$ be arbitrary, and suppose that

$$(2.26) \quad \|S_{\alpha'', \varepsilon}(t)u - S_{\alpha'', 0}(t)u\|_{\alpha''} \leq p_\varepsilon(\alpha^*)e^{\omega t} \|u\|_{\alpha'}, \quad t \in (s, s + T), \quad \varepsilon > 0, \quad u \in C_{\alpha'},$$

and

$$(2.27) \quad \|Z_\varepsilon u - Z_0 u\|_{\alpha''} \leq \sum_{j=1}^r \frac{P_\varepsilon(\alpha^*)}{(\alpha'' - \alpha')^j} \|u\|_{\alpha'}, \quad \varepsilon > 0, \quad u \in B_{\alpha'}.$$

Let $s \geq 0$, $\alpha_s \in (\underline{\alpha}, \alpha^*)$, $u_{s, \varepsilon}, u_{s, 0} \in B_{\alpha_s}$ be arbitrary, and suppose that

$$(2.28) \quad \lim_{\varepsilon \rightarrow 0} \|u_{s, \varepsilon} - u_{s, 0}\|_{\alpha_s} = 0.$$

Then, for any $\varepsilon \geq 0$, there exist a unique solution to the differential equation

$$(2.29) \quad \begin{cases} \frac{d}{dt} u_\varepsilon(t) = (A_\varepsilon + Z_\varepsilon)u_\varepsilon(t), & t \in (s, s + T), \\ u_\varepsilon(s) = u_{s, \varepsilon} \end{cases}$$

in B_{α^*} , where $T = T(\alpha_s, \alpha^*)$; and, moreover, for any $\Upsilon \in (s, s + T)$,

$$(2.30) \quad \lim_{\varepsilon \rightarrow 0} \sup_{t \in [s, s + \Upsilon]} \|u_\varepsilon(t) - u_0(t)\|_{\alpha^*} = 0.$$

Proof. The existence and uniqueness of solutions to (2.29) follow directly from Theorems 2.1 and 2.4. By the proof of Theorem 2.1, it is easy to see that there exists $\alpha = \alpha(\Upsilon) \in (\alpha_s, \alpha^*)$, which does not depend on ε , such that

$$(2.31) \quad u_\varepsilon(t) = \sum_{n=0}^{\infty} V_{\alpha, \varepsilon}^{(n)}(s, t) u_{s, \varepsilon}, \quad \varepsilon \geq 0,$$

where $V_{\alpha, \varepsilon}^{(0)}(s, t) := S_{\alpha, \varepsilon}(t - s)$ and

$$V_{\alpha, \varepsilon}^{(n)}(s, t) := \int_s^t \int_s^{t_1} \dots \int_s^{t_{n-1}} U_{\alpha, \varepsilon}^{(n)}(t, t_1, \dots, t_n) dt_n \dots dt_1,$$

$$U_{\alpha, \varepsilon}^{(n)}(t, t_1, \dots, t_n) := S_{\alpha, \varepsilon}(t - t_1) Z_\varepsilon S_{\alpha, \varepsilon}(t_1 - t_2) Z_\varepsilon \dots S_{\alpha, \varepsilon}(t_{n-1} - t_n) Z_\varepsilon S_{\alpha, \varepsilon}(t_n),$$

and the series (2.31) converges in B_α . Recall that $\Upsilon < T' = T(\alpha_s, \alpha)$ and let, as before, $q \in (1, \frac{T'}{\Upsilon})$.

Therefore, by the proof of Corollary 2.3,

$$\begin{aligned} & \|u_\varepsilon(t) - u_0(t)\|_\alpha \\ & \leq \sum_{n=0}^{\infty} \|(V_{\alpha, \varepsilon}^{(n)}(s, t) - V_{\alpha, 0}^{(n)}(s, t)) u_{s, 0}\|_\alpha + \sum_{n=0}^{\infty} \|V_{\alpha, 0}^{(n)}(s, t) (u_{s, \varepsilon} - u_{s, 0})\|_\alpha \\ & \leq \|(S_{\alpha, \varepsilon}(t - s) - S_{\alpha, 0}(t - s)) u_{s, 0}\|_\alpha \\ & \quad + \sum_{n=1}^{\infty} \int_s^t \int_s^{t_1} \dots \int_s^{t_{n-1}} \|(U_{\alpha, \varepsilon}^{(n)}(t, t_1, \dots, t_n) - U_{\alpha, 0}^{(n)}(t, t_1, \dots, t_n)) u_{s, 0}\|_\alpha dt_n \dots dt_1 \\ & \quad + \frac{C e^{\omega(s + \Upsilon)}}{T' - q\Upsilon} \|u_{s, \varepsilon} - u_{s, 0}\|_{\alpha_s}. \end{aligned}$$

Denote, for simplicity of notations, $Q_{\alpha, \varepsilon}(t) := S_{\alpha, \varepsilon}(t) - S_{\alpha, 0}(t)$, $t \geq 0$, $R_\varepsilon := Z_\varepsilon - Z_0$. Then, for $n \geq 1$,

$$\begin{aligned} & U_{\alpha, \varepsilon}^{(n)}(t, t_1, \dots, t_n) - U_{\alpha, 0}^{(n)}(t, t_1, \dots, t_n) \\ & = Q_{\alpha, \varepsilon}(t - t_1) Z_\varepsilon S_{\alpha, \varepsilon}(t_1 - t_2) Z_\varepsilon \dots S_{\alpha, \varepsilon}(t_{n-1} - t_n) Z_\varepsilon S_{\alpha, \varepsilon}(t_n) \\ & \quad + S_{\alpha, 0}(t - t_1) R_\varepsilon S_{\alpha, \varepsilon}(t_1 - t_2) Z_\varepsilon \dots S_{\alpha, \varepsilon}(t_{n-1} - t_n) Z_\varepsilon S_{\alpha, \varepsilon}(t_n) \\ & \quad + S_{\alpha, 0}(t - t_1) Z_0 Q_{\alpha, \varepsilon}(t_1 - t_2) Z_\varepsilon \dots S_{\alpha, \varepsilon}(t_{n-1} - t_n) Z_\varepsilon S_{\alpha, \varepsilon}(t_n) \\ & \quad + \dots \\ & \quad + S_{\alpha, 0}(t - t_1) Z_0 S_{\alpha, 0}(t_1 - t_2) Z_0 \dots Q_{\alpha, \varepsilon}(t_{n-1} - t_n) Z_\varepsilon S_{\alpha, \varepsilon}(t_n) \\ & \quad + S_{\alpha, 0}(t - t_1) Z_0 S_{\alpha, 0}(t_1 - t_2) Z_0 \dots S_{\alpha, 0}(t_{n-1} - t_n) R_\varepsilon S_{\alpha, \varepsilon}(t_n) \\ & \quad + S_{\alpha, 0}(t - t_1) Z_0 S_{\alpha, 0}(t_1 - t_2) Z_0 \dots S_{\alpha, 0}(t_{n-1} - t_n) Z_0 Q_{\alpha, \varepsilon}(t_n). \end{aligned}$$

By using the partition (2.6)–(2.7), one gets, cf. (2.10),

$$\begin{aligned} & \|U_{\alpha, \varepsilon}^{(n)}(t, t_1, \dots, t_n) - U_{\alpha, 0}^{(n)}(t, t_1, \dots, t_n)\|_\alpha \\ & \leq \nu e^{\omega t} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^{n-1} \left(np_\varepsilon(\alpha) + n\nu \sum_{j=1}^r \frac{P_\varepsilon(\alpha)}{\delta_2^j} \right) \|u\|_{\alpha_s} \\ & \leq \nu e^{\omega t} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^{n-1} \left(n\nu p_\varepsilon(\alpha) + n\nu \sum_{j=1}^r \frac{(qn)^j}{(eT')^j} P_\varepsilon(\alpha) \right) \|u\|_{\alpha_s}. \end{aligned}$$

As a result,

$$\begin{aligned}
 & \|u_\varepsilon(t) - u_0(t)\|_\alpha \\
 & \leq p_\varepsilon(\alpha^*) \nu e^{\omega\Upsilon} \|u_{s,0}\|_{\alpha,s} \\
 & \quad + P_\varepsilon(\alpha) \sum_{n=1}^{\infty} \frac{\Upsilon^n}{n!} \nu e^{\omega(s+\Upsilon)} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^n \left(n\nu \sum_{j=1}^r \frac{(qn)^{j-1}}{(eT')^{j-1}} \right) \|u_{s,0}\|_{\alpha,s} \\
 & \quad + p_\varepsilon(\alpha) \sum_{n=1}^{\infty} \frac{\Upsilon^n}{n!} \nu e^{\omega(s+\Upsilon)} \left(\frac{qn}{eT'} + \nu N(\alpha) \right)^{n-1} n\nu \|u_{s,0}\|_{\alpha,s} \\
 & \quad + \frac{C e^{\omega(s+\Upsilon)}}{T' - q\Upsilon} \|u_{s,\varepsilon} - u_{s,0}\|_{\alpha,s},
 \end{aligned}$$

that fulfills the statement, by (2.28), (2.25); note that convergence of two latter series holds by (2.13). (As a matter of fact, we have proved the convergence (2.30) with a stronger norm $\|\cdot\|_{\alpha,s}$) \square

3. AN APPLICATION TO BIRTH-AND-DEATH DYNAMICS

We will start with a brief introduction to the configuration space analysis. More detailed explanation may be found in, e.g., [22, 13, 14].

Let $\mathcal{B}_b(\mathbb{R}^d)$ be the set of all bounded Borel subsets of \mathbb{R}^d . The configuration space over space \mathbb{R}^d consists of all locally finite subsets (configurations) of \mathbb{R}^d , i.e.

$$(3.1) \quad \Gamma := \{ \gamma \subset \mathbb{R}^d \mid |\gamma_\Lambda| < \infty, \text{ for all } \Lambda \in \mathcal{B}_b(\mathbb{R}^d) \}.$$

Here $|\cdot|$ means the cardinality of a set, and $\gamma_\Lambda := \gamma \cap \Lambda$. The Borel σ -algebra $\mathcal{B}(\Gamma)$ is generated by all mappings $\Gamma \ni \gamma \mapsto |\gamma_\Lambda| \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$. Let $\mathcal{M}_{\text{fm}}^1(\Gamma)$ be the set of all probability measures μ on $(\Gamma, \mathcal{B}(\Gamma))$ such that $\int_\Gamma |\gamma_\Lambda|^n d\mu(\gamma) < \infty$, for any $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ and $n \in \mathbb{N}$.

Let Γ_0 be the space of all finite configurations from \mathbb{R}^d , i.e.

$$(3.2) \quad \Gamma_0 := \{ \eta \subset \mathbb{R}^d \mid |\eta| < \infty \}.$$

Then $\Gamma_0 = \bigsqcup_{n \in \mathbb{N}_0} \Gamma^{(n)}$, where $\Gamma^{(n)} := \{ \eta \subset \mathbb{R}^d \mid |\eta| = n \}$, $n \in \mathbb{N}_0$. Clearly, $\Gamma^{(n)} \sim \widetilde{(\mathbb{R}^d)^n} / S_n$, where the tilde denotes the product set without diagonals and S_n is the permutation group. This isomorphism provides the natural σ -algebra $\mathcal{B}(\Gamma_0)$ on Γ_0 . The Lebesgue–Poisson measure on $(\Gamma_0, \mathcal{B}(\Gamma_0))$ is defined via the following equality:

$$(3.3) \quad \int_{\Gamma_0} G(\eta) d\eta = G^{(0)} + \sum_{n=1}^{\infty} \int_{(\mathbb{R}^d)^n} G^{(n)}(x_1, \dots, x_n) dx_1 \dots dx_n,$$

where G is a measurable nonnegative function on Γ_0 , which may be identified with the sequence of symmetric functions $G^{(n)}$, namely, $G(\{x_1, \dots, x_n\}) = G^{(n)}(x_1, \dots, x_n)$, $G(\emptyset) = G^{(0)} \in \mathbb{R}$.

Let $B_{\text{bs}}(\Gamma_0)$ be the set of all measurable bounded functions $G : \Gamma_0 \rightarrow \mathbb{R}$ such that there exist $N \in \mathbb{N}$ and $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ such that, for $n > N$, $G^{(n)} \equiv 0$, and, for $n \leq N$, $G^{(n)}(x_1, \dots, x_n) = 0$ if only $x_i \notin \Lambda$, for some $1 \leq i \leq n$. Then, for any $G \in B_{\text{bs}}(\Gamma_0)$, one can define the following function on Γ :

$$(3.4) \quad (KG)(\gamma) := \sum_{\eta \in \gamma} G(\eta), \quad \gamma \in \Gamma,$$

where the summation is taken over all finite subconfigurations $\eta \in \Gamma_0$ of the (infinite) configuration $\gamma \in \Gamma$; we denote this by the symbol, $\eta \in \gamma$. The mapping K is linear,

positivity preserving, and invertible, with

$$(3.5) \quad (K^{-1}F)(\eta) := \sum_{\xi \subset \eta} (-1)^{|\eta \setminus \xi|} F(\xi), \quad \eta \in \Gamma_0.$$

It can be shown that, for $G \in B_{\text{bs}}(\Gamma_0)$ with $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$, $N \in \mathbb{N}$ as above, $KG(\gamma) = KG(\gamma_\Lambda)$ and $|KG(\gamma)| \leq C(1 + |\gamma_\Lambda|)^N$, $\gamma \in \Gamma$. In particular, $KG \in L^1(\Gamma, \mu)$, for any $\mu \in \mathcal{M}_{\text{fm}}^1(\Gamma)$. The correlation function of a measure $\mu \in \mathcal{M}_{\text{fm}}^1(\Gamma)$ is the function $k_\mu : \Gamma_0 \rightarrow \mathbb{R}_+$ which satisfies the identity

$$(3.6) \quad \int_{\Gamma} (KG)(\gamma) d\mu(\gamma) = \int_{\Gamma_0} G(\eta) k_\mu(\eta) d\eta,$$

for any $0 \leq G \in B_{\text{bs}}(\Gamma_0)$, provided such k_μ does exist.

We consider a model which is a combination of models discussed in [8] and [17].

Let $a, \phi : \mathbb{R}^d \rightarrow \mathbb{R}_+ := [0, \infty)$ be measurable nonnegative symmetric functions, i.e. $a(-x) = a(x)$, $\phi(-x) = \phi(x)$, $x \in \mathbb{R}^d$. Assume that $a, \phi \in L^1(\mathbb{R}^d, dx) \cap L^\infty(\mathbb{R}^d, dx)$. Set

$$(3.7) \quad \langle a \rangle := \int_{\mathbb{R}^d} a(x) dx, \quad \bar{a} := \text{ess sup}_{x \in \mathbb{R}^d} a(x), \quad \langle \phi \rangle := \int_{\mathbb{R}^d} \phi(x) dx, \quad \bar{\phi} := \text{ess sup}_{x \in \mathbb{R}^d} \phi(x).$$

Let $m, \lambda > 0$ be constants. For any $F \in K(B_{\text{bs}}(\Gamma_0))$, we define the mapping

$$(3.8) \quad \begin{aligned} (LF)(\gamma) &= \sum_{x \in \gamma} \left(\sum_{y \in \gamma \setminus x} a(x-y) + m \exp\left(-\sum_{y \in \gamma \setminus x} \phi(x-y)\right) \right) (F(\gamma \setminus x) - F(\gamma)) \\ &+ \lambda \int_{\mathbb{R}^d} (F(\gamma \cup x) - F(\gamma)) dx. \end{aligned}$$

Here and below we use the notations $\setminus x$ and $\cup x$ instead of more precise $\setminus \{x\}$ and $\cup \{x\}$, respectively. Heuristically, L describes the following evolution of configurations: during a (small) time t in an arbitrary domain $\Lambda \in \mathcal{B}_b(\mathbb{R}^d)$ a new elements may appear with the probability $\lambda \text{vol}(\Lambda)t + o(t)$, whereas the probability for the existing point $x \in \gamma$ disappears is equal to $\sum_{y \in \gamma \setminus x} a(x-y)t + m \exp(-\sum_{y \in \gamma \setminus x} \phi(x-y))t + o(t)$. Thus this probability will be close to 1 in very dense regions of the space as well as in the almost ‘uninhabited’ places.

Since $F(\gamma \cup x) = F((\gamma \cup x)_\Lambda)$, the integrand in (3.8) equals to 0 outside of Λ , thus the integral is well-defined. By the same arguments, the first (outer) sum in (3.8) is taken over $x \in \gamma_\Lambda$ only. The other sums (in y) are, however, infinite. In particular, (3.8) is well-defined for all $\gamma \in \Gamma$, if, say, a has a bounded support. It is worth noting, that, regardless of a , (3.8) is defined pointwise, for $\gamma \in \Gamma_0$. This is sufficient to consider

$$(3.9) \quad (\widehat{L}G)(\eta) := (K^{-1}LKG)(\eta), \quad \eta \in \Gamma_0, \quad G \in B_{\text{bs}}(\Gamma_0).$$

By results of [18] and [17, Proposition 3.1], one has that, for any $G \in B_{\text{bs}}(\Gamma_0)$, $\eta \in \Gamma_0$,

$$(3.10) \quad \begin{aligned} (\widehat{L}G)(\eta) &= -E^a(\eta)G(\eta) - \sum_{x \in \eta} \left(\sum_{y \in \eta \setminus x} a(x-y) \right) G(\eta \setminus x) \\ &- m \sum_{\xi \subset \eta} G(\xi) \sum_{x \in \xi} e^{-E^\phi(x, \xi \setminus x)} e_\lambda(e^{-\phi(x-\cdot)} - 1, \eta \setminus \xi) \\ &+ \lambda \int_{\mathbb{R}^d} G(\eta \cup x) dx, \end{aligned}$$

where

$$(3.11) \quad E^a(\eta) := \sum_{x \in \eta} \sum_{y \in \gamma \setminus x} a(x - y), \quad \eta \in \Gamma_0,$$

$$(3.12) \quad E^\phi(x, \eta \setminus x) := \sum_{y \in \gamma \setminus x} \phi(x - y), \quad \eta \in \Gamma_0, \quad x \in \eta,$$

and, for any measurable $f : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$(3.13) \quad e_\lambda(f, \eta) := \prod_{x \in \eta} f(x), \quad \eta \in \Gamma_0 \setminus \{\emptyset\}, \quad e_\lambda(f, \emptyset) := 1.$$

The mapping L defines an evolution of measures in $\mathcal{M}_{\text{fin}}^1(\Gamma)$. Namely, for a given $\mu_0 \in \mathcal{M}_{\text{fin}}^1(\Gamma)$, consider the initial value problem

$$(3.14) \quad \begin{cases} \frac{d}{dt} \int_\Gamma F(\gamma) d\mu_t(\gamma) = \int_\Gamma (LF)(\gamma) d\mu_t(\gamma), & t > 0, \\ \mu_t|_{t=0} = \mu_0, \end{cases}$$

which should hold for any $F \in K(B_{\text{bs}}(\Gamma_0))$ such that the right hand side of (3.14) is well-defined. The equation (3.14) may be rewritten in terms of the correlation functions $k_t := k_{\mu_t}$ of measures $\mu_t \in \mathcal{M}_{\text{fin}}^1(\Gamma)$, provided that they all do exist. Namely, one has

$$(3.15) \quad \begin{cases} \frac{d}{dt} \int_{\Gamma_0} G(\eta) k_t(\eta) d\eta = \int_{\Gamma_0} (\widehat{L}G)(\eta) k_t(\eta) d\eta, & t > 0, \\ k_t|_{t=0} = k_0 = k_{\mu_0}. \end{cases}$$

The latter equation will be the main object of our interest. For relations between solutions to (3.15) and (3.14) see, e.g., [13]. One can rewrite (3.15) in the ‘‘strong’’ form

$$(3.16) \quad \frac{\partial}{\partial t} k_t(\eta) = (L^\Delta k_t)(\eta), \quad \eta \in \Gamma_0, \quad t > 0,$$

where the linear mapping L^Δ is defined via the duality

$$(3.17) \quad \int_{\Gamma_0} (\widehat{L}G)(\eta) k(\eta) d\eta = \int_{\Gamma_0} G(\eta) (L^\Delta k)(\eta) d\eta,$$

for $G, k \in B_{\text{bs}}(\Gamma_0)$, and it is extended to the linear operator in (a scale of) Banach spaces by the constructions below. By, e.g., [18] and [17], one has that

$$(3.18) \quad \begin{aligned} (L^\Delta k)(\eta) &= -E^a(\eta)k(\eta) - \sum_{y \in \eta} \int_{\mathbb{R}^d} a(x - y)k(\eta \cup x) dx \\ &\quad - m \sum_{x \in \eta} e^{-E^\phi(x, \eta \setminus x)} \int_{\Gamma_0} k(\eta \cup \xi) e_\lambda(e^{-\phi(x-\cdot)} - 1, \xi) d\xi \\ &\quad + \lambda \sum_{x \in \eta} k(\eta \setminus x), \quad \eta \in \Gamma_0. \end{aligned}$$

We consider the following scale of Banach spaces:

$$(3.19) \quad \mathcal{K}_\alpha := \{k : \Gamma_0 \rightarrow \mathbb{R} \mid k(\eta)\alpha^{-|\eta|} \in L^\infty(\Gamma_0, d\eta)\}, \quad \alpha > 1,$$

with the norms given by

$$(3.20) \quad \|k\|_\alpha := \text{ess sup}_{\eta \in \Gamma_0} |k(\eta)|\alpha^{-|\eta|}.$$

For the motivation, see, e.g., [13, 14]. It is easy to see, that $\{\mathcal{K}_\alpha\}_{\alpha > 1}$ satisfies to Assumption 1 with $\underline{\alpha} = 1$, $I = (1, \infty)$.

We set, for $\eta \in \Gamma_0$,

$$(Ak)(\eta) = -E^a(\eta)k(\eta), \quad (Zk)(\eta) = (L^\Delta k) - (Ak)(\eta).$$

Proposition 3.1. *The linear mappings A and Z satisfy Assumptions 2 and 3, correspondingly.*

Proof. The operator A with the maximal domain

$$(3.21) \quad \mathcal{D}_\alpha := \{k \in \mathcal{K}_\alpha \mid Ek \in \mathcal{K}_\alpha\},$$

naturally, generates the semigroup

$$(3.22) \quad (S(t)k)(\eta) = e^{-tE(\eta)}k(\eta), \quad \eta \in \Gamma_0,$$

in any \mathcal{K}_α , $\alpha > 1$. However, this semigroup is not a C_0 one. Indeed, for $k_\alpha(\eta) := \alpha^{-|\eta|}$, one has

$$\|S(t)k_\alpha - k_\alpha\|_\alpha = \operatorname{ess\,sup}_{\eta \in \Gamma_0} |e^{-tE(\eta)} - 1| = 1 \not\rightarrow 0, \quad t \rightarrow 0.$$

Therefore, one should use the technique of the \odot -dual semigroups; for details see, e.g., [34, 7]. Namely, we consider the C_0 -semigroup given by the same expression (3.22), but considered in the space $\mathcal{L}_\alpha := L^1(\Gamma_0, \alpha^{|\eta|} d\eta)$. Then $S(t)$ is dual to that semigroup in the dual space \mathcal{K}_α (where duality is realized by (3.17)). Then, the space $C_\alpha := \overline{\mathcal{D}_\alpha}$ (the closure is in the norm of \mathcal{K}_α) is $S(t)$ -invariant and the restriction $S_\alpha(t) := S(t) \upharpoonright_{C_\alpha}$ consists a C_0 -semigroup there. The generator of $S_\alpha(t)$ will be the part of A , i.e. (A, D_α) , where $D_\alpha = \{k \in C_\alpha \mid Ak \in C_\alpha\}$, cf. [13]. Hence D_α is $S(t)$ -invariant as well. It should be stressed also that $\mathcal{K}_{\alpha'} \subset D_\alpha$, for any $1 < \alpha' < \alpha$. Indeed, for a $k \in \mathcal{K}_{\alpha'}$,

$$(3.23) \quad \alpha^{-|\eta|} E^\alpha(\eta) |k(\eta)| \leq \|k\|_{\alpha'} \bar{a} |\eta|^2 \left(\frac{\alpha'}{\alpha}\right)^{|\eta|} \leq \frac{4\|k\|_{\alpha'} \bar{a}}{e^2 \ln^2 \frac{\alpha'}{\alpha}},$$

where we used that $\sup_{r>0} r^2 q^r = 4/(e \ln q)^2$, for $q \in (0, 1)$. Since $|S(t)k| \leq |k|$ pointwise, the space $\mathcal{K}_{\alpha'}$ is also $S(t)$ -invariant. From these arguments we easily get that A satisfies Assumption 2 with $\nu = 1$, $\omega = 0$.

Next, let us denote

$$(Z^{(1)}k)(\eta) := - \sum_{y \in \eta} \int_{\mathbb{R}^d} a(x-y)k(\eta \cup x) dx.$$

Then, for any $1 < \alpha' < \alpha'' < \alpha^*$ and for any $k \in \mathcal{K}_{\alpha'}$,

$$(\alpha'')^{-|\eta|} |(Z^{(1)}k)(\eta)| \leq \langle a \rangle \alpha' \|k\|_{\alpha'} |\eta| \left(\frac{\alpha'}{\alpha''}\right)^{|\eta|} \leq \langle a \rangle \alpha' \|k\|_{\alpha'} \frac{1}{-e \ln \frac{\alpha'}{\alpha''}} \leq \frac{\langle a \rangle (\alpha^*)^2 \|k\|_{\alpha'}}{e(\alpha'' - \alpha')},$$

where we used that $\sup_{r>0} r q^r = 1/(e \ln q)$, for $q \in (0, 1)$, and $\ln \alpha'' - \ln \alpha' = \frac{1}{\tilde{\alpha}}(c'' - c')$, for some $\tilde{\alpha} \in (\alpha', \alpha'')$.

The similar estimate for $(Z^{(2)}k)(\eta) := (Zk)(\eta) - (Z^{(1)}k)(\eta)$ was obtained in [17, Proposition 3.2]. Combining these results, one gets that Z satisfies Assumption 3, with

$$M(\alpha^*) = \frac{1}{e} (\langle a \rangle (\alpha^*)^2 + \alpha^* m e^{\langle \phi \rangle \alpha^*} + \alpha^* \lambda).$$

(To be more precise, we used here the estimate $\int_{\mathbb{R}^d} (1 - e^{-\phi(x)}) dx \leq \langle \phi \rangle$, to simplify the expression from [17, Proposition 3.2].) \square

Consider now the so-called Vlasov scaling of the dynamics above, see, e.g., [12, 13]. Namely, for an $\varepsilon > 0$, we denote by L_ε the operator (3.8) with $\varepsilon a(\cdot)$, $\varepsilon \phi(\cdot)$, $\varepsilon^{-1} \lambda$ in place of $a(\cdot)$, $\phi(\cdot)$, λ , respectively. Then, one can construct L_ε^Δ in the same way as above. We set also

$$(3.24) \quad (L_{\varepsilon, \text{ren}}^\Delta k)(\eta) := \varepsilon^{|\eta|} L_\varepsilon^\Delta \varepsilon^{-|\eta|} k(\eta).$$

Directly from (3.18), one gets

$$(3.25) \quad \begin{aligned} (L_{\varepsilon, \text{ren}}^{\Delta} k)(\eta) &= -\varepsilon E^a(\eta)k(\eta) - \sum_{y \in \eta} \int_{\mathbb{R}^d} a(x-y)k(\eta \cup x) dx \\ &- m \sum_{x \in \eta} e^{-\varepsilon E^{\phi}(x, \eta \setminus x)} \int_{\Gamma_0} k(\eta \cup \xi) e_{\lambda} \left(\frac{e^{-\varepsilon \phi(x-\cdot)} - 1}{\varepsilon}, \xi \right) d\xi \\ &+ \lambda \sum_{x \in \eta} k(\eta \setminus x), \quad \eta \in \Gamma_0. \end{aligned}$$

We denote, for $\varepsilon > 0$,

$$(A_{\varepsilon} k)(\eta) := -\varepsilon E^a(\eta)k(\eta), \quad (Z_{\varepsilon} k)(\eta) := (L_{\varepsilon, \text{ren}}^{\Delta} k)(\eta) - (A_{\varepsilon} k)(\eta),$$

and we set, naturally, $(A_0 k)(\eta) := 0$ and

$$\begin{aligned} (Z_0 k)(\eta) &:= - \sum_{y \in \eta} \int_{\mathbb{R}^d} a(x-y)k(\eta \cup x) dx \\ &- m \sum_{x \in \eta} \int_{\Gamma_0} k(\eta \cup \xi) e_{\lambda}(-\phi(x-\cdot), \xi) d\xi + \lambda \sum_{x \in \eta} k(\eta \setminus x). \end{aligned}$$

Proposition 3.2. *The linear mappings A_{ε} and Z_{ε} , $\varepsilon \geq 0$ satisfy Assumptions 4 and 5, correspondingly. Moreover, the conditions (2.26), (2.27), (2.25) of Theorem 2.10 hold.*

Proof. The operators A_{ε} satisfy Assumption 4 by the same arguments as in the proof of Proposition 3.1 (independently on $\varepsilon \geq 0$). The operators Z_{ε} satisfy Assumption 5 by the estimation for $Z^{(1)}$ in the proof of Proposition 3.1 and by [17, Proposition 4.2] for $Z_{\varepsilon} - Z^{(1)}$, $\varepsilon \geq 0$.

Next, in the notations of Theorem 2.10, for any $1 < \alpha' < \alpha'' < \alpha^*$ and for any $k \in \alpha'$,

$$\begin{aligned} \|S_{\varepsilon}(t)k - S_0(t)k\|_{\alpha''} &= \text{ess sup}_{\eta \in \Gamma_0} (\alpha'')^{-|\eta|} |e^{-t\varepsilon E^a(\eta)} - 1| \|k(\eta)\| \\ &\leq \|k\|_{\alpha'}(s+T)\varepsilon \text{ess sup}_{\eta \in \Gamma_0} E^a(\eta) \left(\frac{\alpha'}{\alpha}\right)^{|\eta|}, \end{aligned}$$

that implies (2.26), by using the same estimate as in (3.23).

Finally,

$$\begin{aligned} (Z_{\varepsilon} k)(\eta) - (Z_0 k)(\eta) &= -m \sum_{x \in \eta} e^{-\varepsilon E^{\phi}(x, \eta \setminus x)} \int_{\Gamma_0} k(\eta \cup \xi) e_{\lambda} \left(\frac{e^{-\varepsilon \phi(x-\cdot)} - 1}{\varepsilon}, \xi \right) d\xi \\ &+ m \sum_{x \in \eta} \int_{\Gamma_0} k(\eta \cup \xi) e_{\lambda}(-\phi(x-\cdot), \xi) d\xi, \end{aligned}$$

and the estimate (2.27) (with $r = 2$) was proved in [17, Proposition 4.6]. \square

Suppose now, for simplicity, that $k_{0, \varepsilon} := k_0 \in \mathcal{K}_{\alpha_0}$, $\alpha_0 > 1$, $\varepsilon > 0$. Then, by Theorem 2.10, the solutions to the equation

$$(3.26) \quad \frac{\partial}{\partial t} k_{t, \varepsilon}(\eta) = (L_{\varepsilon, \text{ren}}^{\Delta} k_{t, \varepsilon})(\eta), \quad \eta \in \Gamma_0, \quad t \in (0, T),$$

converges in any \mathcal{K}_{α^*} with $\alpha^* > \alpha_0$ to the solution to the equation

$$(3.27) \quad \frac{\partial}{\partial t} k_t(\eta) = (Z_0 k_t)(\eta), \quad \eta \in \Gamma_0, \quad t \in (0, T),$$

uniformly on any $[0, \Upsilon] \subset (0, T)$, where $T = T(\alpha_0, \alpha^*)$.

The limiting equation (3.27) has the following key-property: if $k_0(\eta) = e_\lambda(\rho_0, \eta)$, for a function $\rho_0 \in L^\infty(\mathbb{R}^d, dx)$ then one can find a (unique) solution to (3.27) of the same form: $k_t(\eta) = e_\lambda(\rho_t, \eta)$. To show this, note that

$$\frac{\partial}{\partial t} e_\lambda(\rho_t, \eta) = \sum_{x \in \eta} \rho_t(x) e_\lambda(\rho_t, \eta \setminus x)$$

and

$$\begin{aligned} (Z_0 e_\lambda(\rho_t))(\eta) &:= - \sum_{x \in \eta} \rho_t(x) \int_{\mathbb{R}^d} a(x-y) \rho_t(y) dy e_\lambda(\rho_t, \eta \setminus x) \\ &\quad - m \sum_{x \in \eta} \rho_t(x) e_\lambda(\rho_t, \eta \setminus x) \int_{\Gamma_0} e_\lambda(\rho_t, \xi) e_\lambda(-\phi(x-\cdot), \xi) d\xi \\ &\quad + \lambda \sum_{x \in \eta} e_\lambda(\rho_t, \eta \setminus x). \end{aligned}$$

By (3.3), we have, for any $f \in L^1(\mathbb{R}^d, dx)$,

$$\int_{\Gamma_0} e_\lambda(f, \eta) d\eta = \exp\left\{ \int_{\mathbb{R}^d} f(x) dx \right\}.$$

Therefore, $k_t(\eta) = e_\lambda(\rho_t, \eta)$ indeed solves (3.27), provided that ρ_t is a unique solution to the following equation

$$(3.28) \quad \frac{\partial}{\partial t} \rho_t(x) = -\rho_t(x)(a * \rho_t)(x) - m\rho_t(x)e^{-(\phi * \rho_t)(x)} + \lambda$$

in the space $L^\infty(\mathbb{R}^d, dx)$ (at least on $(0, T)$).

The existence and uniqueness of nonnegative solutions to (3.28) may be done using the same approaches as in [17, 10]. We will realize this in a sequel paper.

Remark 3.3. It is worth noting that the equation (3.28) may have one or three positive stationary solutions depending on values of the parameters. Indeed, if $\rho_t(x) \equiv \rho > 0$ is a stationary solution to (3.28), then $\lambda = \langle a \rangle \rho^2 + m\rho \exp(-\langle \phi \rangle \rho)$. Denote $x = \langle \phi \rangle \rho > 0$, $c = \frac{\lambda \langle \phi \rangle}{m}$, $b = \frac{\langle a \rangle}{m \langle \phi \rangle}$; then we will get $xe^{-x} + bx^2 = c$. The function $f(x) = xe^{-x} + bx^2$, $x \geq 0$, may have zero or two points of local extremum. Indeed, $f'(x) = 0$ yields $2bx = (x-1)e^{-x}$. The function $g(x) = (x-1)e^{-x}$, $x \geq 0$, has the derivative $g'(x) = (2-x)e^{-x}$ and hence g increases from -1 to e^{-2} on $(0, 2)$ and decreases for $x > 2$. The tangent line to the graph of $y = g(x)$ at a point $(x_0, g(x_0))$ which passes through the origin has the equation $y - g(x_0) = g'(x_0)(x - x_0)$, and thus $x = y = 0$ yields

$$-(x_0 - 1)e^{-x_0} = (2 - x_0)e^{-x_0}(-x_0), \quad x_0^2 - x_0 - 1 = 0, \quad x_0 = \frac{1 + \sqrt{5}}{2} > 0.$$

As a result, if $2b < g'(x_0)$, i.e. if

$$\frac{\langle a \rangle}{m \langle \phi \rangle} < \frac{3 - \sqrt{5}}{4} \exp\left(-\frac{1 + \sqrt{5}}{2}\right),$$

then the function $f(x)$ has two points of local extremum and, therefore, there exists λ (and thus c) such that the equation $f(x) = c$ has three solutions. For $2b \geq g'(x_0)$, it will have one solution only, for any $\lambda > 0$.

REFERENCES

1. E. A. Barkova, P. P. Zabreiko, *The Cauchy problem for differential equations of fractional order with deteriorating right-hand sides*, *Differencial'nye Uravneniya* **42** (2006), 1132–1134. (Russian); English transl. *Differential Equations* **42** (2006), 1199–1202.
2. T. D. Benavides, *Generic existence of a solution for a differential equation in a scale of Banach spaces*, *Proc. Amer. Math. Soc.* **86** (1982), no. 3, 477–484.

3. Yu. M. Berezanskii, *Selfadjoint Operators in Spaces of Functions of Infinitely Many Variables*, Amer. Math. Soc., Providence, RI, 1986. (Russian edition: Naukova Dumka, Kiev, 1978)
4. C. Berns, Y. Kondratiev, Y. Kozitsky, and O. Kutoviy, *Kawasaki dynamics in continuum: micro- and mesoscopic descriptions*, J. Dyn. Diff. Equat. **25** (2013), no. 4, 1027–1056.
5. C. Berns, Y. Kondratiev, and O. Kutoviy, *Construction of a state evolution for Kawasaki dynamics in continuum*, Analysis and Mathematical Physics **3** (2013), no. 2, 97–117.
6. R. E. Caflisch, *A simplified version of the abstract Cauchy-Kowalewski theorem with weak singularities*, Bull. Amer. Math. Soc. (N.S.) **23** (1990), no. 2, 495–500.
7. K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*, vol. 194 of *Graduate Texts in Mathematics*, Springer-Verlag, New York, 2000.
8. D. Finkelshtein and Y. Kondratiev, *Regulation mechanisms in spatial stochastic development models*, J. Stat. Phys. **136** (2009), no. 1, 103–115.
9. D. Finkelshtein, Y. Kondratiev, and Y. Kozitsky, *Glauber dynamics in continuum: a constructive approach to evolution of states*, Discrete and Cont. Dynam. Syst., Ser A **33** (2013), no. 4, 1431–1450.
10. D. Finkelshtein, Y. Kondratiev, Y. Kozitsky, and O. Kutoviy, *The statistical dynamics of a spatial logistic model and the related kinetic equation*, Mathematical Models and Methods in Applied Sciences **25** (2015), no. 2, 343–370.
11. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Individual based model with competition in spatial ecology*, SIAM J. Math. Anal. **41** (2009), no. 1, 297–317.
12. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Vlasov scaling for stochastic dynamics of continuous systems*, J. Stat. Phys. **141** (2010), no. 1, 158–178.
13. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Semigroup approach to birth-and-death stochastic dynamics in continuum*, J. Funct. Anal. **262** (2012), no. 3, 1274–1308.
14. D. Finkelshtein, Y. Kondratiev, and O. Kutoviy, *Statistical dynamics of continuous systems: perturbative and approximative approaches*, Arabian Journal of Mathematics, 2014.
15. D. Finkelshtein, Y. Kondratiev, O. Kutoviy, and E. Lytvynov, *Binary jumps in continuum. II. Non-equilibrium process and a Vlasov-type scaling limit*, J. Math. Phys. **52** (2011), 113301:1–27.
16. D. Finkelshtein, Y. Kondratiev, O. Kutoviy, and M. J. Oliveira, *Dynamical Widom–Rowlinson model and its mesoscopic limit*, J. Stat. Phys. **158** (2015), no. 1, 57–86.
17. D. Finkelshtein, Y. Kondratiev, O. Kutoviy, and E. Zhizhina, *On an aggregation in birth-and-death stochastic dynamics*, Nonlinearity **27** (2014), no. 6, 1105–1133.
18. D. Finkelshtein, Y. Kondratiev, and M. J. Oliveira, *Markov evolutions and hierarchical equations in the continuum. I. One-component systems*, Journal of Evolution Equations **9** (2009), no. 2, 197–233.
19. D. Finkelshtein, Y. Kondratiev, and M. J. Oliveira, *Glauber dynamics in the continuum via generating functionals evolution*, Complex Analysis and Operator Theory **6** (2012), no. 4, 923–945.
20. D. Finkelshtein, Y. Kondratiev, and M. J. Oliveira, *Kawasaki dynamics in the continuum via generating functionals evolution*, Methods Funct. Anal. Topology **18** (2012), no. 1, 55–67.
21. I. M. Gel'fand and G. E. Šilov, *Nekotorye voprosy teorii differentsialnykh uravnenii*, Obobščennye funkcii, Vypusk 3, Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow, 1958; English transl. I. M. Gelfand and G. E. Shilov, *Generalized Functions Vol. 3 Theory of Differential Equations*, Academic Press, New York—London, 1967.
22. Y. Kondratiev and T. Kuna, *Harmonic analysis on configuration space. I. General theory*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **5** (2002), no. 2, 201–233.
23. Y. Kondratiev, O. Kutoviy, and R. Minlos, *On non-equilibrium stochastic dynamics for interacting particle systems in continuum*, J. Funct. Anal. **255** (2008), no. 1, 200–227.
24. Y. Kondratiev, O. Kutoviy, and E. Zhizhina, *Nonequilibrium Glauber-type dynamics in continuum*, J. Math. Phys. **47** (2006), no. 11, 113501:1–17.
25. S. G. Lobanov and O. G. Smolyanov, *Ordinary differential equations in locally convex spaces*, Uspekhi Mat. Nauk **49** (1994), no. 3, 93–168. (Russian); English transl. Russian Math. Surveys **49** (1994), no. 3, 97–175.
26. H. P. Lotz, *Uniform convergence of operators on L^∞ and similar spaces*, Math. Z. **190** (1985), no. 2, 207–220.
27. L. Nirenberg, *An abstract form of the nonlinear Cauchy-Kowalewski theorem*, Journal of Differential Geometry **6** (1972), no. 4, 561–576.
28. T. Nishida, *A note on a theorem of Nirenberg*, Journal of Differential Geometry **12** (1977), no. 4, 629–633.
29. L. V. Ovsjannikov, *Singular operator in the scale of Banach spaces*, Dokl. Akad. Nauk SSSR **163** (1965), 819–822. (Russian); English transl. Soviet Math. Dokl. **6** (1965), 1025–1028.

30. R. S. Phillips, *The adjoint semi-group*, Pacific J. Math. **5** (1955), 269–283.
31. M. V. Safonov, *The abstract Cauchy-Kovalevskaya theorem in a weighted Banach space*, Comm. Pure Appl. Math. **48** (1995), no. 6, 629–637.
32. F. Trèves, *Ovcyannikov Theorem and Hyperdifferential Operators*, Instituto de Matemática Pura e Aplicada, Conselho Nacional de Pesquisas, 1968.
33. F. Trèves, *An abstract nonlinear Cauchy-Kovalevska theorem*, Trans. Amer. Math. Soc. **150** (1970), 77–92.
34. J. van Neerven, *The Adjoint of a Semigroup of Linear Operators*, vol. 1529 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1992.
35. P. P. Zabreiko, *Existence and uniqueness theorems for solutions of the Cauchy problem for differential equations with worsening operators*, Dokl. Akad. Nauk BSSR **33** (1989), no. 12, 1068–1071. (Russian)
36. O. Zubelevich, *Abstract version of the Cauchy-Kowalewski problem*, Cent. Eur. J. Math. **2** (2004), no. 3, 382–387. (electronic)

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3 TERESHCHENKIVS'KA, KYIV, 01601, UKRAINE

Current address: Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, U.K.

E-mail address: d.l.finkelshtein@swansea.ac.uk

Received 20/01/2015